



Original article

The effectiveness of a 17-week lifestyle intervention on health behaviors among airline pilots during COVID-19

Daniel Wilson^{a,b,*}, Matthew Driller^c, Ben Johnston^d, Nicholas Gill^{a,e}

^a Te Huataki Waiora School of Health, The University of Waikato, Hamilton 3216, New Zealand

^b Faculty of Health, Education and Environment, Toi Ohomai Institute of Technology, Tauranga 3112, New Zealand

^c Sport and Exercise Science, Human Services and Sport, La Trobe University, Melbourne 3086, Australia

^d Aviation and Occupational Medicine Unit, Wellington School of Medicine, Otago University, Wellington 6242, New Zealand

^e New Zealand Rugby, Wellington 6011, New Zealand

Received 14 August 2020; revised 24 September 2020; accepted 15 October 2020

2095-2546/© 2020 Published by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Abstract

Purpose: The aim of this study was to evaluate the efficacy of a 17-week, 3-component lifestyle intervention for enhancing health behaviors during the coronavirus disease-2019 (COVID-19) pandemic.

Methods: A parallel-group (intervention and control) study was conducted amongst 79 airline pilots over a 17-week period during the COVID-19 pandemic. The intervention group ($n = 38$) received a personalized sleep, dietary, and physical activity (PA) program. The control group ($n = 41$) received no intervention. Outcome measures for sleep, fruit and vegetable intake, PA, and subjective health were measured through an online survey before and after the 17-week period. The changes in outcome measures were used to determine the efficacy of the intervention.

Results: Significant main effects for Time \times Group were found for International Physical Activity Questionnaire-Walk ($p = 0.02$) and for all other outcome measures ($p < 0.01$). The intervention group significantly improved in sleep duration ($p < 0.01$; $d = 1.02$), Pittsburgh Sleep Quality Index score ($p < 0.01$; $d = -1.01$), moderate-to-vigorous PA ($p < 0.01$; $d = 1.32$), fruit and vegetable intake ($p < 0.01$; $d = 3.11$), Short-Form-12v2 physical score ($p < 0.01$; $d = 1.84$), and Short-Form-12v2 mental score ($p < 0.01$; $d = 2.69$). The control group showed significant negative change for sleep duration ($p < 0.01$; $d = -0.47$), Pittsburgh Sleep Quality Index score ($p < 0.01$; $d = 0.28$), and Short-Form-12v2 mental score ($p < 0.01$; $d = -0.64$).

Conclusion: Results provide preliminary evidence that a 3-component healthy sleep, eating and PA intervention elicit improvements in health behaviors and perceived subjective health in pilots and may improve quality of life during an unprecedented global pandemic.

Keywords: COVID-19; Healthy eating; Lifestyle health; Moderate-to-vigorous physical activity; Sleep

1. Introduction

The global coronavirus disease-2019 (COVID-19) pandemic has rapidly spread, showing capability to infect the world's population.¹ Widespread infectious diseases such as COVID-19 are associated with adverse mental health consequences² and perturbations in physical activity (PA) behaviors due to environmental factors such as forced self-isolation.³ The confinement of individuals to their homes may increase sedentary behavior⁴ and has a direct impact on lifestyle, and, consequently, on sleeping, eating, and PA patterns.⁵ Furthermore, psychological and emotional responses to the pandemic⁶

may lead to dysfunctional dietary and sleep behaviors.^{7,8} Numerous industries have experienced substantial operational perturbations emanating from COVID-19, including civil aviation.⁹ Consequently, these conditions may have unexplored impacts on the health behaviors of airline pilots. Vocational requirements of airline pilots present health risks, such as circadian disruption due to shift work and flight schedules,¹⁰ fatigue induced by flight schedules,¹¹ irregular meal times, mental stress demands associated with flight safety,¹² and the sedentary nature of the job.¹³ Circadian disruption is detrimental to acute physiological¹⁴ and psychological¹⁵ health metrics and is associated with elevated risk for some chronic conditions such as cardiovascular disease.¹⁶ Despite reduced workloads for airline pilots during COVID-19,⁹ substantial industry disruption, uncertainty,¹⁷ and financial concerns confounded

Peer review under responsibility of Shanghai University of Sport.

*Corresponding author.

E-mail address: Daniel.Wilson@toiohomai.ac.nz (D. Wilson).

<https://doi.org/10.1016/j.jshs.2020.11.007>

by lockdown conditions may present adverse effects on the physical^{3,18} and mental health¹⁹ of pilots.

Obtaining adequate sleep, consuming enough fruits and vegetables, and engaging in sufficient PA are 3 lifestyle behaviors that significantly reduce all-cause mortality^{20–22} and have a positive effect on physical and mental health.^{23, 24} Good sleep health facilitates the ability to maintain attentive wakefulness and is characterized by duration, quality, timing, and efficiency.²⁵ Sleep duration guidelines proposed for adults from 18 to 64 years is 7–9 h per night.²⁶ Fruit and vegetables supply dietary fiber, vitamins and minerals, phytochemicals, and anti-inflammatory agents.²⁷ Consumption of 400 g or more of fruit and vegetables per day, excluding starchy vegetables, is associated with protective effects against cardiovascular disease, some cancers,²⁸ depression,²⁹ and total mortality.²⁰ An inverse association between fruit and vegetable intake and mortality has been reported, with benefits observed in up to 7 daily portions.³⁰ Sufficient PA is defined as the achievement of 150 min/week or more of moderate-to-vigorous PA (MVPA), or 75 min/week or more of vigorous-intensity PA, or an accumulative equivalent combination of both, with added health benefits of 300 or more total MVPA minutes per week.³¹

Adequate sleep, healthy dietary behaviors, and sufficient PA also play an important role in strengthening the immune system and its antiviral defenses.^{32,33} Lack of sleep, poor dietary habits, and physical inactivity are all independently associated with immunocompromising effects,^{32,34,35} which impair host defenses against viral infection and may lead to individuals being at higher risk of more severe and complicated outcomes than those which are non-immunocompromised.³⁶ A lack of sleep,³⁷ dietary characteristics such as consuming a Western diet,³⁸ and insufficient PA are each associated with obesity,³⁹ which is suggested to be a profound risk factor for adverse health outcomes from COVID-19.⁴⁰

Avoidance of health behaviors during a pandemic outbreak may lead to immunocompromise, increased susceptibility to viral propagation and elevated risk of severe symptoms.^{36,41} Behavioral countermeasures for individuals are vital determinants to health resilience amongst exposure to unprecedented environmental events such as the COVID-19 pandemic.⁵ Thus, evidence-driven interventions targeting the promotion of behaviors that enable individuals to protect themselves physically and psychologically during a pandemic are of public health importance.^{5,42} No previous studies have examined change in health behaviors in airline pilots before and after a pandemic event like COVID-19, nor have any studies evaluated the effectiveness of a controlled lifestyle-based health intervention during such times. The aim of this study was to evaluate the effectiveness of a 3-component healthy eating, sleeping and PA program in airline pilots during COVID-19 lockdown in New Zealand.

2. Methods

2.1. Design

A between-group, parallel controlled study with pre and post-testing was performed to evaluate the effectiveness of a

17-week, 3-component lifestyle intervention for enhancing health behaviors during a national COVID-19 pandemic response in New Zealand.

During the 17-week intervention period, the first 5 weeks preceded the New Zealand government's enforcement of a 4-tier response system to COVID-19.¹ Thereafter, 5 weeks were at alert Level 4 (March 25 to April 27, 2020), where all non-essential workers were instructed to stay home, and safe recreational activity was permitted only in the local area whilst maintaining social distancing of 2 m or more. Subsequently, 2.5 weeks were at alert Level 3 (April 27 to May 14, 2020), where some businesses could reopen; however, people were instructed to work from home unless that was not possible. Finally, 2 weeks were at alert Level 2 (May 14 to June 8, 2020), where gatherings of up to 100 people were allowed and sport and recreation activities were also allowed. On June 8, New Zealand returned to alert Level 1,⁴³ where within-community restrictions were removed yet international border restrictions remained. Around the globe in the weeks following the World Health Organization's characterization of COVID-19 as a pandemic on March 11, 2020, the airline industry experienced a decrease of approximately 60%–80% in capacity at major carriers.⁹ Consequently, pilots experienced limited flying duties during the time when the intervention was conducted (Table 1). Pre-testing occurred between February 14 and March 9, 2020, and post-testing was conducted between June 8 and June 19, 2020. At pre- and post-testing, participants completed an electronic survey measuring the following outcome variables: self-report PA levels, dietary behaviors, quality and quantity of sleep, and subjective health.

2.2. Participants

The study population consisted of commercial pilots from a large international airline. Seventy-nine pilots (aged 42 ± 12 years, mean ± SD, age, 65 male, 14 female) participated in the study (Table 1). Participants were from a combination of short haul, long haul, and mixed fleet rosters ($n = 32, 35,$ and $12,$ respectively). Inclusion criteria were pilots who had a valid commercial flying license and worked on a full-time basis. Control group participants consisted of volunteers who were pilots recruited at the time they completed their routine aviation medical examinations at the airline medical unit during the pre-test period (between February 14 and March 9, 2020). The intervention group volunteered to participate in the lifestyle intervention via self-selection. All volunteers participating in the study had responded to an invitation delivered to all pilots within the company via internal organization communication channels. In the control group, mixed-fleet pilots had the most flights, followed by long-haul and short-haul pilots ($10 \pm 9, 7 \pm 7,$ and $6 \pm 5;$ respectively). In the intervention group, mixed-fleet pilots had the most flights, followed by short-haul and long-haul pilots ($14 \pm 1, 9 \pm 9,$ and $6 \pm 4,$ respectively). All participants provided written informed consent prior to participation in the study and were made aware that they could withdraw from the study at any time should they wished to do so. In order to support anonymity and

Table 1
Baseline demographic characteristics of participants (mean \pm SD; *n*).

Parameters	All subjects (<i>n</i> = 79)	Intervention (<i>n</i> = 38)	Control (<i>n</i> = 41)
Sex (F/M)	14/65	8/30	6/35
Age (year)	42 \pm 12	39 \pm 10	44 \pm 13
Short haul (<i>n</i>)	32	22	10
Long haul (<i>n</i>)	35	14	21
Mixed fleet (<i>n</i>)	12	2	10
Flights during lockdown (<i>n</i>)	7.8 \pm 7.3	5.9 \pm 7.5	7.2 \pm 7.1

Abbreviations: F = female; M = male.

dataset blinding during data analysis, participants were provided with a unique identification code on their informed consent form and were instructed to input it into their survey instead of their name. This study was approved by the Human Research Ethics Committee of the University of Waikato in New Zealand (reference number 2020#07).

2.3. Intervention group

Participants who registered their interest in participating in the intervention and agreed to attend a face-to-face consultation session at the airline medical unit were included in the intervention group. The intervention group received an initial one-on-one 60-min consultation session with an experienced health coach, followed by provision of an individualized health program, weekly educational content emails during the intervention and a mid-intervention phone call. The health advice provided was evidence based and derived from experts in the areas of PA, nutrition and chronobiology.

Personalized goal setting was carried out for each participant in the intervention group, with relevant outcome, performance and process goals⁴⁴ discussed pertaining to each of the 3 intervention components: (1) dietary behaviors, (2) PA, and (3) sleep habits. Moreover, individual perceived barriers to health change were assessed with methods outlined elsewhere⁴⁵ and were factored into the individualized program.

Sleep hygiene as an educational strategy has demonstrated improvements in self-report sleep quality in blue collar employees.⁴⁶ Stimulus control as a behavior technique has demonstrated positive outcomes on sleep parameters in insomniacs.⁴⁷ An evidence-based sleep health checklist was developed for utilization in this study (Appendix). The items in the checklist consisted of standard recommendations derived from previous sleep hygiene and stimulus control studies.^{48,49} Application of the items in the sleep health checklist involved collaborative identification of the strategies that participants were achieving at baseline. Pilots completed the Pittsburgh Sleep Quality Index (PSQI) prior to attending their consultation. During the face-to-face consultation, the health coach discussed suboptimal PSQI component scores with the participant and identified individual sleep priorities. Thereafter, participants collaboratively set personalized sleep practice goals with support from the health coach. Personalized collaborative goal setting was implemented as a behavioral technique to support development of sleep habits that support restorative sleep.⁴⁴

PA prescription was individualized based on participant perceived barriers and facilitators to exercise;⁴⁵ application of the frequency, intensity, time, and type principles;⁵⁰ and progression to attainment of sufficient MVPA to meet guidelines³¹ in congruence with individual capabilities. The frequency and time of PA sessions were tailored to participant time availability. Intensity was tailored based on participant exercise experience, physical health, and goal orientation.⁵¹ The type of PA was determined by the individual's modality preferences for cardiovascular (such as walking, running, or cycling) and strengthening (e.g., resistance equipment and/or bodyweight exercises) PA. The PA progression self-monitoring was indicated, and participants were advised to implement small, progressive changes in PA during the intervention (such as increased session duration, more repetitions, greater exercise intensity or more weekly bouts).⁵² Healthy eating principles were emphasized through individualized advice and educational materials focused on adding color to the diet via consumption of fruit and vegetables;²⁷ choosing nutrient-dense foods;⁵³ limiting processed foods; enhancing whole-food consumption;⁵⁴ and reducing white carbohydrates, refined carbohydrates, and added sugar (e.g., energy-dense food).⁵⁵ Relative to participant baseline behaviors, collaborative individualized process behavior goals were established, for example, adding color to meals and replacing high glycemic index foods with low glycemic index options. The prescribed dose of fruit was two or more servings per day, and the prescribed dose of vegetables was three or more servings per day. A mid-intervention (during Week 8 \pm 1) follow-up phone call lasting approximately 10 min consisted of a semi-structured interview emphasizing discussion of progress and compliance pertaining to individualized sleep, PA, and dietary goals established during the pre-test consultation. Advice was provided when necessary and was consistent with the advice given at pre-test. Phone calls were used because they have been shown to support adherence to health interventions.⁵⁶ Congruent with evidence-based methods previously outlined, weekly emails were sent consisting of educational blog posts on varying topics related to sleep health, PA, nutrition, and support for a healthy immune system. Content was derived from health authorities via publicly available information from the World Health Organization and the Centers for Disease Control and Prevention. During the COVID-19 lockdown, content was tailored to pandemic conditions, including strategies for PA at home, healthy recipes and immune system health information.

2.4. Control group

The participants in the control group were blind to the intervention and received no intervention or instruction regarding health behaviors between pre- and post-tests. Control group pilots were invited to voluntarily complete an electronic survey, and those who completed it during the previously defined pre-testing period prior to COVID-19 lockdown were sent an invitation via email to voluntarily complete the survey again during the post-testing period in order to provide insight into the effects of COVID-19 lockdown on their health behaviors.

2.5. Instruments

The PSQI was utilized to evaluate subjective sleep quality, and scores were obtained before the start of the study at the pre-test and were compared to scores obtained at the post-test stage. The PSQI is a self-rated, 19-item questionnaire designed to measure sleep quality and disturbances over a 1-month retrospective period.⁵⁷ Sleep quality component scores are derived for subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medications, and daytime dysfunction, and collectively produce a global sleep score.⁵⁷ Lower scores denote a healthier sleep quality and range from 0 (no difficulties) to 21 (severe sleep difficulties).⁵⁷ The PSQI has demonstrated good test-retest reliability and validity and has been implemented in many population groups.^{57,58} The outcome measurements were the change in global score between each group at post-testing.

PA levels were assessed using the International Physical Activity Questionnaire (IPAQ) Short Form, a validated self-report measurement tool for MVPA that has been widely utilized in large cohort studies, including New Zealand population surveys.⁵⁹ The IPAQ Short Form estimates PA achievement by quantifying weekly walking, as well as moderate and vigorous PA duration and frequency. Responses were used to compare participants' PA levels with the health guidelines of 150 min of moderate-intensity PA or 75 min of vigorous-intensity PA per week, or an equivalent combination of MVPA per week.³¹ IPAQ Short Form outcome measures derived were (a) total weekly minutes of moderate + vigorous PA in bouts of 10 min or more, excluding walking (IPAQ–MVPA), and (b) total weekly minutes of walking in bouts of 10 min or more (IPAQ–walk). Responses were capped at 3 h per day and 21 h per week, as recommended by the IPAQ guidelines.⁶⁰ Fruit and vegetable intake were measured using 2 questions with acceptable validity and reliability derived from the New Zealand Health Survey.⁵⁹ The questions asked participants to report, on average, over the last week, how many servings of fruit and vegetables they ate per day. Responses to these questions were combined to determine total daily fruit and vegetable intake.

Subjective self-report physical and mental health was determined using the Short Health Form 12v2 (SF-12v2), a short version of the SF-36, which reduces the burden on participants and has demonstrated a high correlation with SF-36 physical and mental component summary scale scores.⁶¹ The 12-item

survey produces a physical component summary scale (PCS-12) and a mental component summary scale (MCS-12), both of which have shown good test-retest reliability and convergent validity in detecting changes in mental and physical health over time in adults.⁶² Scoring of the SF-12v2 was carried out in accordance with standard summary scoring methods.⁶¹ The summary scores are on a scale of 0–100, with clinical significance change scores suggested to be 5–10 points.⁶³

2.6. Statistical analyses

For all statistical analyses, raw data were extracted from the Qualtrics online survey software (Qualtrics, Provo, UT, USA), entered into an Excel spreadsheet (Microsoft, Seattle, WA, USA) and then imported into the Statistical Package for the Social Sciences (SPSS) (Version 26; IBM, New York, NY, USA). Listwise deletion was applied for individual datasets with missing values or participants who did not complete post-testing. Stem and leaf plots were inspected to ascertain whether there were any outliers in the data for each variable. A Shapiro-Wilk's test ($p > 0.05$) and its histograms, Q–Q plots and box plots were inspected for normality of data distribution for all variables. Levene's test was used to test homogeneity of variance. Time-related effects within and between groups on pre-test and post-test were assessed using t tests and repeated measures mixed-design analysis of variance. Age, sex, and number of flights were included as covariates in the analysis of variance. Effect sizes were calculated using Cohen's d in order to quantify between-group effects from pre-testing to post-testing. Effect size thresholds were set at greater than 1.2, greater than 0.6, greater than 0.2, and greater than 0.2, which were classified as large, moderate, small, and trivial, respectively.⁶⁴ The α level was set at a p value of less than 0.05.

3. Results

The demographic and baseline health characteristics between the intervention and control groups are given in Table 1. The attrition rates were 16% and 30% for the intervention and control group, respectively. Intervention group attrition was influenced by employment layoffs, whereas control group attrition was due to non-response to the online invitation to voluntarily complete the survey at the post-test period. In order to measure compliance, average sleep hours, exercise sessions per week and daily fruit and vegetable consumption was collected at the time of the mid-intervention phone call. In the intervention group, 35 participants (92%) were achieving 7.0 h or more of sleep per night and 3 (8%) were achieving 6.9 h or less per night. For fruit and vegetable servings per day, 36 participants (95%) were eating 5 or more servings of fruit and vegetables per day, whereas 2 (5%) were eating 2–4 servings per day. For MVPA, 33 participants (87%) were completing 150 min or more of MVPA per week and 5 (13%) were completing 149 min or less of MVPA per week.

At baseline, the control group achieved significantly ($p < 0.05$) more sleep per night (36 min), a lower global PSQI score

(−1.8) (denoting better health), higher consumption of daily fruit and vegetables (1.2 servings), a greater amount of weekly walking (94 min), and higher SF-12v2 physical and mental component scores (8.4 and 6.7, respectively). At baseline, the control group exhibited an MVPA of 171 ± 78 min per week, surpassing the MVPA recommended in the health guidelines of 150 or more min per week,³¹ whereas the intervention group exhibited 116 ± 51 min per week of MVPA, which is beneath the health guidelines threshold. However, the between-group difference in MVPA was not statistically significant ($p \geq 0.05$).

Group changes from pre-test to post-test are presented in Table 2. Significant main effects for time \times group were found for IPAQ-Walk ($p = 0.02$) and for all other outcome measures ($p < 0.01$), which is associated with small to large effect size differences between groups (Table 2). Significant main effects for time were found for PSQI, fruit and vegetable intake, PCS-12 and MCS-12 ($p < 0.01$). Additional significant time effects were found for hours slept and MVPA ($p < 0.05$). The within-group analysis revealed that the intervention elicited significant improvements ($p < 0.01$) in all health metrics at post-test and IPAQ-walk ($p = 0.02$). The control group reported a significantly higher PSQI score ($p < 0.01$), decreased hours of sleep and MCS-12 score ($p < 0.01$), yet no significant change was reported in other health metrics.

4. Discussion

To our knowledge, this study is the first controlled experiment that has explored the effects of a lifestyle intervention on health outcomes among pilots during a national pandemic lockdown. This study aimed to improve health-related behaviors and promote positive change in subjective health of pilots through personalized advice on healthy eating, sleep hygiene, and PA. For most outcome measures, the controlled trial demonstrated significant improvements in the intervention group compared to the control group. These results are important in order for researchers and health care professionals to provide insight into potential health and quality-of-life perturbations resulting from COVID-19 that may have potential implications to flight safety. Furthermore, given the dearth of published

data pertaining to health behavior interventions during a pandemic and the limited availability of preventive lifestyle-based interventions for pilots, our findings provide novel contributions to this field.

The average PSQI score for the intervention group decreased (−2.25), compared to a significant increase for the control group (0.76). These results support previous studies that used non-pilot populations, which have reported PSQI score decreases of −1.54 to −1.8 following a sleep hygiene education intervention^{46,65} and −2.5 after a PA and sleep education intervention based on a mobile health app.⁶⁶ An app-based intervention to reduce fatigue in a pilot population reported a smaller effect on PSQI score (−0.59) after a 6-month intervention that focused on advice regarding daylight exposure and sleep duration. None of these studies were conducted under global pandemic conditions. A potential confounding factor to sleep quality and quantity improvements during COVID-19 due to social isolation and lockdown constraints has been proposed,⁶⁷ where more time at home and flexible sleep–wake schedules may promote enhancements in sleep. Furthermore, sleep disruption is an inherent risk for pilots, and they are likely to have better sleep quality and quantity when not at work.

Curiously, in our study the control group significantly increased its PSQI score. Similar findings have been reported elsewhere in a general-population-based study in Australia ($n = 1491$), which reported that 40.1% of the participants indicated a negative change in sleep quantity during lockdown.⁸ In another study, sleep quantity improved while sleep quality was degraded in Austrian adults ($n = 435$).⁶⁸ In that study, an increase in subject burden and decreased physical and mental wellbeing were also observed. Cognitive states of elevated distress and emotional disturbances have been associated with unhealthy dietary patterns and poor diet quality and may impair health behavior motivation.⁵ Researchers have advocated for implementation of strategies to mitigate the effects of lockdown on sleep quality, including obtaining sufficient PA, exposure to natural daylight,⁶⁸ and well-balanced meals rich in vitamins and minerals.⁵ Our study findings support these messages in that we observed significant improvements in PCS-12 and MCS-12 scores among the intervention group

Table 2
Changes in health behaviors from baseline to post-test at 17 weeks (mean \pm SD).

	Intervention group			Control group			ANOVA	Effect size
	Pre	Post	Change	Pre	Post	Change	(time \times group interaction) <i>p</i>	<i>d</i>
Hours slept (h:mm)	7:00 \pm 0:54	7:48 \pm 0:48**	0:48 \pm 0:54	7:36 \pm 0:48	7:12 \pm 1:00**	−0:24 \pm 0:42	<0.01	1.35, Large
PSQI global	6.4 \pm 2.7	4.1 \pm 1.8**	−2.3 \pm 2.4	4.5 \pm 2.5	5.3 \pm 2.9**	0.8 \pm 0.9	<0.01	1.14, Large
IPAQ-walk	61 \pm 69	93 \pm 96	32 \pm 105	155 \pm 135	136 \pm 111	−19 \pm 79	0.02	0.45, Small
IPAQ-MVPA	116 \pm 51	201 \pm 77**	85 \pm 100	171 \pm 78	146 \pm 75	−26 \pm 82	<0.01	1.44, Large
Fruit and vegetable intake	3.0 \pm 0.6	6.1 \pm 1.7**	3.1 \pm 1.5	4.2 \pm 0.9	4.1 \pm 1.5	−0.1 \pm 1.3	<0.01	2.09, Large
PCS-12	43.3 \pm 5.2	52.6 \pm 4.9**	9.4 \pm 5.5	51.7 \pm 5.7	51.5 \pm 6.0	−0.2 \pm 2.2	<0.01	1.52, Large
MCS-12	48.0 \pm 4.0	54.8 \pm 1.0**	6.8 \pm 4.1	54.7 \pm 4.3	51.9 \pm 4.5**	−2.8 \pm 4.4	<0.01	2.09, Large

d = Cohen's *d* effect size, effect threshold. * $p < 0.05$, ** $p < 0.01$.

Abbreviations: ANOVA = analysis of variance; IPAQ = International Physical Activity Questionnaire; MCS-12 = SF-12v2 Mental Component Summary Scale; MVPA = moderate-to-vigorous physical activity; PCS-12 = SF-12v2 Physical Component Scale; PSQI = Pittsburgh Sleep Quality Index; SF-12v2 = Short Health Form 12v2.

compared to the control group, after implementation of a 3-component healthy sleep, eating and PA intervention during an unprecedented global pandemic.

The average MVPA and fruit and vegetable consumption significantly increased in our intervention group, compared to no significant change in the control group. The intervention group increased MVPA from 116 min to 201 min at post-testing, which crossed the MVPA guideline threshold of 150 min or more per week.³¹ Conversely, the control group decreased MVPA from 171 min to 146 min per week at post-testing, decreasing to below the guideline threshold. Both the control and intervention groups did not achieve the recommendation of 5 or more servings per day for fruit and vegetable intake at baseline.⁶⁹ The intervention group elevated its intake per day, achieving the guideline threshold at post-testing. Our study findings suggest that the 3-component intervention supported achievement of PA and fruit and vegetable guideline thresholds and significantly improved PCS-12 and MCS-12 scores in the intervention group. Furthermore, the intervention appeared to mitigate decay in both SF-12v2 summary scales, which were observed in the control group.

Research exploring the relationships among COVID-19, PA and dietary behaviors have yielded mixed outcomes. Within an Italian population ($n = 3533$) during COVID-19 lockdown, similar proportions of the participants stated that they ate less or had better diets (35.8% and 37.4%, respectively) in regard to intake of fruit, vegetables, nuts, and legumes.⁷ Furthermore, their PA behaviors did not significantly change; however, a greater amount of exercise was completed at home.⁷ In a study with Australians participants, 48.9% expressed a negative change in PA during lockdown and also reported that negative changes in PA and sleep were associated with expression of higher anxiety, depression and stress levels.⁸ In a Canadian sample ($n = 1098$), those who engaged in more outdoor PA time during lockdown had lower anxiety levels than those who did not. The COVID-19 lockdown in New Zealand at Levels 3–4 presented barriers for engaging in PA, such as social distancing, travel restrictions and inaccessibility to parks, gyms, and other recreational facilities, which may have promoted sedentary behavior and contributed to the decline in PA observed in the control group.^{4,17} Other studies found that life stressors, including job uncertainty,⁹ anxiety and psychological stress,² financial loss and disconnection with community and nature may have impacted health behaviors during this time.¹⁷ In contrast, the reduced time at work has been suggested as a potential facilitative factor in enhancing health behaviors during lockdown,⁶⁷ because this naturally alleviates one of the most commonly expressed barriers to engagement in PA: a lack of time.

A limited number of studies have investigated the efficacy of interventions targeting healthy dietary and PA behaviors among pilots. A study of an app-based, 6-month intervention using a pilot population reported significant improvements in their weekly moderate and strenuous activity (0.21 and 0.19 days per week, respectively), as well as a reduction in their snacking behavior (-0.88 servings per duty).⁷⁰ Consultations with pilots on their diet and physical exercise behaviors yielded a positive change in their blood lipids and body mass

index.¹² Our findings pertaining to MVPA, fruit and vegetable intake, PSQI and SF-12v2 among pilots provide promising preliminary outcomes regarding the effects of a 3-component intervention and warrant further investigation with objective measures of outcomes.

The differential recruitment strategies and limited exclusion criteria for the intervention and control groups are limitations that may have contributed to the significant differences observed at baseline for sleep quantity and PSQI score, IPAQ-walk, MVPA, fruit and vegetable consumption and SF-12v2 subjective health scores, with superior results in favor of the control group. Thus, it is recommended that future research studies have more robust randomization assignment conditions for participant allocation to groups in order to increase the probability of capturing the true population average and enhance the generalizability of findings. Future research should also increase the sample size, given the apparent variances in health behaviors amongst the pilot population, which in itself warrants further investigation in order to characterize these variances.

Future research with pilot populations that compare single-behavior to multiple-behavior interventions would provide valuable insight into the magnitude of the effects when these differing intervention approaches are used. Given the bidirectional relationship between sleep, nutrition and PA and some evidence to suggest multiple-component interventions may elicit stronger participation and adherence,⁷¹ we suggest future research examine more time-efficient and scalable strategies for implementation of a 3-component sleep, nutrition and PA program. For feasibility reasons, self-report methods were utilized in this study, and the limitations of this approach have been discussed elsewhere.⁷² To enhance outcome measure validity and reliability, it would be preferable to use objective measurement methods such as actigraphy to monitor sleep and PA and photo logging of dietary behaviors to quantify health behavior metrics. Comparisons of our study to related studies are limited because most other studies were conducted under normal societal conditions while ours was conducted under pandemic conditions.

5. Conclusion

Behavioral countermeasures for individuals are vital determinants of health resilience during exposure to unprecedented environmental events such as the COVID-19 pandemic.⁵ The attainment of sleep, fruit and vegetable intake and PA guidelines is associated with increased physical and mental health,^{24,29} enhanced immune defenses,³³ and a reduction in the risk of obesity.³⁹ Evidence-based interventions targeting the promotion of these behaviors will enable individuals to protect themselves physically and psychologically during a pandemic, and therefore are of immense public health importance.^{5,42} The 3-component healthy sleep, eating, and PA intervention implemented in this study elicited significant improvements in sleep quality and quantity, fruit and vegetable intake, and MVPA and suggests that achieving these 3 guideline thresholds promotes mental and physical health and

improves quality of life among pilots during a global pandemic. Our study of this intervention provides preliminary evidence that a low-intensity, multi-behavior intervention may be efficacious during a pandemic and that similar outcomes may be transferrable to other populations. However, more robust recruitment methods are required to confirm our findings and increase their generalizability.

Acknowledgments

The authors wish to thank the pilots for participating in this study.

Authors' contributions

DW and NG participated in conceptualization of the study and data collection; DW, MD, BJ, and NG contributed to the design of the study, data analysis, interpretation of the results and manuscript writing. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

The authors declare they have no competing interests.

Appendix

Sleep hygiene strategies for enhancing sleep	YES Achieving	NOT Achieving
1. Sleep at least 7 h		
2. Sleep routine or depower hour		
3. Regular sleep and wake time		
4. Dim lights near bedtime and turn off electronics >30 min before bed		
5. Avoid sleep disruptors 4–6 h before bed, e.g., caffeine, large meals, alcohol		
6. Have a dark, cool, quiet sleep environment		
7. Exercise every day, not too close to bedtime		
8. Use the bedroom only for sleeping and intimacy		
9. Do a brain dump on paper before bed		
10. Early morning light exposure		

References

- Baker M, Kvalsvig A, Verrall AJ, Telfar-Barnard L, Wilson N. New Zealand's elimination strategy for the COVID-19 pandemic and what is required to make it work. *N Z Med J* 2020;**133**:10–4.
- Bao Y, Sun Y, Meng S, Shi J, Lu L. 2019-nCoV epidemic: address mental health care to empower society. *The Lancet* 2020;**395**:e37–8. doi:10.1016/S0140-6736(20)30309-3.
- Hammami A, Harrabi B, Mohr M, Krstrup P. Physical activity and coronavirus disease 2019 (COVID-19): specific recommendations for home-based physical training. *Manag Sport Leis* 2020. doi:10.1080/23750472.2020.1757494.
- Hobbs M, Pearson N, Foster PJ, Biddle SJ. Sedentary behaviour and diet across the lifespan: an updated systematic review. *Br J Sports Med* 2015;**49**:1179–88.
- Naja F, Hamadeh R. Nutrition amid the COVID-19 pandemic: a multi-level framework for action. *Eur J Clin Nutr* 2020;**74**:1117–21.
- Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate psychological responses and associated factors during the initial stage of the 2019 Coronavirus Disease (COVID-19) epidemic among the general population in China. *Int J Environ Res Public Health* 2020;**17**:1729. doi:10.3390/ijerph17051729.
- Di Renzo L, Gualtieri P, Pivari F, Soldati L, Attinà A, Cinelli G, et al. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J Transl Med* 2020;**18**:229. doi:10.1186/s12967-020-02399-5.
- Stanton R, To QG, Khalesi S, Williams SL, Alley SJ, Thwaite TL, et al. Depression, anxiety and stress during COVID-19: associations with changes in physical activity, sleep, tobacco and alcohol use in Australian adults. *Int J Environ Res Public Health* 2020;**17**:4065. doi:10.3390/ijerph17114065.
- Sobieralski JB. COVID-19 and airline employment: insights from historical uncertainty shocks to the industry. *TRIP* 2020;**5**: 100123. doi:10.1016/j.trip.2020.100123.
- Reis C, Mestre C, Canhão H, Gradwell D, Paiva T. Sleep complaints and fatigue of airline pilots. *Sleep Sci* 2016;**9**:73–7.
- Petrilli RM, Roach GD, Dawson D, Lamond N. The sleep, subjective fatigue, and sustained attention of commercial airline pilots during an international pattern. *Chronobiol Int* 2006;**23**:1347–62.
- Choi YY, Kim KY. Effects of physical examination and diet consultation on serum cholesterol and health-behavior in the Korean pilots employed in commercial airline. *Ind Health* 2013;**51**:603–11.
- Sykes AJ, Larsen PD, Griffiths RF, Aldington S. A study of airline pilot morbidity. *Aviat Space Environ Med* 2012;**83**:1001–5.
- Buxton OM, Cain SW, O'Connor SP, Porter JH, Duffy JF, Wang W, et al. Adverse metabolic consequences in humans of prolonged sleep restriction combined with circadian disruption. *Sci Transl Med* 2012;**4**:129ra43. doi:10.1126/scitranslmed.3003200.
- Walker 2nd WH, Walton JC, DeVries AC, Nelson RJ. Circadian rhythm disruption and mental health. *Transl Psychiatry* 2020;**10**:28. doi:10.1038/s41398-020-0694-0.
- Puttonen S, Härmä M, Hublin C. Shift work and cardiovascular disease — pathways from circadian stress to morbidity. *Scand J Work Environ Health* 2010;**36**:96–108.
- Matias T, Dominski FH, Marks DF. Human needs in COVID-19 isolation. *J Health Psychol* 2020;**25**:871–82.
- Woods J, Hutchinson NT, Powers SK, Roberts WO, Gomez-Cabrera MC, Radak Z, et al. The COVID-19 pandemic and physical activity. *Sports Med Health Sci* 2020;**2**:55–64.
- Wilson JM, Lee J, Fitzgerald HN, Oosterhoff B, Sevi B, Shook NJ. Job insecurity and financial concern during the COVID-19 pandemic are associated with worse mental health. *J Occup Environ Med* 2020;**62**:686–91.
- Bellavia A, Larsson SC, Bottai M, Wolk A, Orsini N. Fruit and vegetable consumption and all-cause mortality: a dose-response analysis. *Am J Clin Nutr* 2013;**98**:454–9.
- Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Sleep duration and all-cause mortality: a systematic review and meta-analysis of prospective studies. *Sleep* 2010;**33**:585–92.
- Lear SA, Hu W, Rangarajan S, Gasevic D, Leong D, Iqbal R, et al. The effect of physical activity on mortality and cardiovascular disease in 130 000 people from 17 high-income, middle-income, and low-income countries: the PURE study. *The Lancet* 2017;**390**:2643–54.
- Mozaffarian D, Wilson PW, Kannel WB. Beyond established and novel risk factors: lifestyle risk factors for cardiovascular disease. *Circulation* 2008;**117**:3031–8.
- Mandolesi L, Polverino A, Montuori S, Foti F, Ferraioli G, Sorrentino P, et al. Effects of physical exercise on cognitive functioning and wellbeing: biological and psychological benefits. *Front Psychol* 2018;**9**:509. doi:10.3389/fpsyg.2018.00509.
- Buysse DJ. Sleep health: can we define it? Does it matter? *Sleep* 2014;**37**:9–17.
- Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's updated sleep duration recommendations: final report. *Sleep Health* 2015;**1**:233–43.
- Slavin JL, Lloyd B. Health benefits of fruits and vegetables. *Adv Nutr* 2012;**3**:506–16.
- Aune D, Giovannucci E, Boffetta P, Fadnes LT, Keum N, Norat T, et al. Fruit and vegetable intake and the risk of cardiovascular disease, total cancer and all-cause mortality—a systematic review and dose-response meta-analysis of prospective studies. *Int J Epidemiol* 2017;**46**:1029–56.

29. McMartin SE, Jacka FN, Colman I. The association between fruit and vegetable consumption and mental health disorders: evidence from five waves of a national survey of Canadians. *Prev Med* 2013;**56**:225–30.
30. Oyebo O, Gordon-Dseagu V, Walker A, Mindell JS. Fruit and vegetable consumption and all-cause, cancer and CVD mortality: analysis of Health Survey for England data. *J Epidemiology Community Health* 2014;**68**:856–62.
31. World Health Organization. *Global recommendations on physical activity for health*. Geneva: World Health Organization; 2010.
32. Walsh NP, Gleeson M, Shephard RJ, Gleeson M, Woods JA, Bishop NC, et al. Position statement. Part one: immune function and exercise. *Exerc Immunol Rev* 2011;**17**:6–63.
33. Valdés-Ramos R, Martínez-Carrillo BE, Aranda-González II, Guadarrama AL, Pardo-Morales RV, Tlatempa P, et al. Diet, exercise and gut mucosal immunity. *Proc Nutr Soc* 2010;**69**:644–50.
34. Tobaldini E, Costantino G, Solbiati M, Cogliati C, Kara T, Nobili L, et al. Sleep, sleep deprivation, autonomic nervous system and cardiovascular diseases. *Neurosci Biobehav Rev* 2017;**74**:321–9.
35. Butler MJ, Barrientos RM. The impact of nutrition on COVID-19 susceptibility and long-term consequences. *Brain Behav Immun* 2020;**87**:53–4.
36. Memoli MJ, Athota R, Reed S, Czajkowski L, Bristol T, Proudfoot K, et al. The natural history of influenza infection in the severely immunocompromised vs. nonimmunocompromised hosts. *Clin Infect Dis* 2014;**58**:214–24.
37. Cappuccio FP, Taggart FM, Kandala NB, Currie A, Peile E, Stranges S, et al. Meta-analysis of short sleep duration and obesity in children and adults. *Sleep* 2008;**31**:619–26.
38. Kanoski SE, Davidson TL. Western diet consumption and cognitive impairment: links to hippocampal dysfunction and obesity. *Physiol Behav* 2011;**103**:59–68.
39. Cecchini M, Sassi F, Lauer JA, Lee YY, Guajardo-Barron V, Chisholm D. Tackling of unhealthy diets, physical inactivity, and obesity: health effects and cost-effectiveness. *The Lancet* 2010;**376**:1775–84.
40. Földi M, Farkas N, Kiss S, Zádori N, Váncsa S, Szakó L, et al. Obesity is a risk factor for developing critical condition in COVID-19 patients: a systematic review and meta-analysis. *Obes Rev* 2020;**21**:e13095. doi:10.1111/obr.13095.
41. Shi Y, Wang Y, Shao C, Huang J, Gan J, Huang X, et al. COVID-19 infection: the perspectives on immune responses. *Cell Death Differ* 2020;**27**:1451–4.
42. Reissman DB, Watson PJ, Klomp RW, Tanielian TL, Prior SD. Pandemic influenza preparedness: adaptive responses to an evolving challenge. *J Homel Secur Emerg Manag* 2006;**3**:1–28.
43. Ministry of Health. *COVID-19 (novel coronavirus)*. Available at: <https://www.health.govt.nz/our-work/diseases-and-conditions/covid-19-novel-coronavirus?mega=Our%20work&title=COVID-19>. [accessed 02.07.2020].
44. Weinberg R. Making goals effective: a primer for coaches. *J Sport Psychol Action* 2010;**1**:57–65.
45. Kulavic K, Hultquist CN, McLester JR. A comparison of motivational factors and barriers to physical activity among traditional versus nontraditional college students. *J Am Coll Health* 2013;**61**:60–6.
46. Nishinoue N, Takano T, Kaku A, Eto R, Kato N, Ono Y, et al. Effects of sleep hygiene education and behavioral therapy on sleep quality of white-collar workers: a randomized controlled trial. *Ind Health* 2012;**50**:123–31.
47. Riedel B, Lichstein K, Peterson BA, Epperson MT, Means MK, Aguillard RN. A comparison of the efficacy of stimulus control for medicated and nonmedicated insomniacs. *Behav Modif* 1998;**22**:3–28.
48. Morgenthaler T, Kramer M, Alessi C, Friedman L, Boehlecke B, Brown T, et al. Practice parameters for the psychological and behavioral treatment of insomnia: an update. An American Academy of Sleep Medicine report. *Sleep* 2006;**29**:1415–9.
49. Perlis M, Aloia M, Millikan A, Boehmler J, Smith M, Greenblatt D, et al. Behavioral treatment of insomnia: a clinical case series study. *J Behav Med* 2000;**23**:149–61.
50. Barisic A, Leatherdale ST, Kreiger N. Importance of Frequency, Intensity, Time and Type (FITT) in physical activity assessment for epidemiological research. *Can J Public Health* 2011;**102**:174–5.
51. Norton K, Norton L, Sadgrove D. Position statement on physical activity and exercise intensity terminology. *J Sci Med Sport* 2010;**13**:496–502.
52. Piercy KL, Troiano RP, Ballard RM, Carlson SA, Fulton JE, Galuska DA, et al. The physical activity guidelines for Americans. *JAMA* 2018;**320**:2020–8.
53. Di Noia J. Defining powerhouse fruits and vegetables: a nutrient density approach. *Prev Chronic Dis* 2014;**11**:E95. doi:10.5888/pcd11.130390.
54. Shahidi F. Nutraceuticals and functional foods: whole versus processed foods. *Trends Food Sci Technol* 2009;**20**:376–87.
55. Drewnowski A. Obesity and the food environment: dietary energy density and diet costs. *Am J Prev Med* 2004;**27**:154–62.
56. Nesari M, Zakerimoghdam M, Rajab A, Bassampour S, Faghihzadeh S. Effect of telephone follow-up on adherence to a diabetes therapeutic regimen. *Jpn J Nurs Sci* 2010;**7**:121–8.
57. Buysse DJ, Reynolds 3rd CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989;**28**:193–213.
58. Manzar MD, BaHammam AS, Hameed UA, Spence DW, Pandi-Perumal SR, Moscovitch A, et al. Dimensionality of the Pittsburgh Sleep Quality Index: a systematic review. *Health Qual Life Outcomes* 2018;**16**:89. doi:10.1186/s12955-018-0915-x.
59. Ministry of Health. *Methodology Report 2017/18: New Zealand Health Survey*. Available at: <https://www.health.govt.nz/publication/methodology-report-2017-18-new-zealand-health-survey>. [accessed 02.07.2020].
60. IPAQ Research Committee. *Guidelines for data processing and analysis of the International Physical Activity Questionnaire (IPAQ)-short and long forms*. Available at: <https://sites.google.com/site/theipaq/scoring-protocol>. [accessed 02.07.2020].
61. Ware JE, Keller SD, Kosinski M. *SF-12: How to score the SF-12 physical and mental health summary scales*. Boston, MA: Health Institute, New England Medical Center; 1995.
62. Cheak-Zamora NC, Wyrwich KW, McBride TD. Reliability and validity of the SF-12v2 in the medical expenditure panel survey. *Qual Life Res* 2009;**18**:727–35.
63. Samsa G, Edelman D, Rothman ML, Williams GR, Lipscomb J, Matchar D. Determining clinically important differences in health status measures: a general approach with illustration to the Health Utilities Index Mark II. *Pharmacoeconomics* 1999;**15**:141–55.
64. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Florence: Taylor & Francis Group; 1988.
65. Falloon K, Elley CR, Fernando 3rd A, Lee AC, Arroll B. Simplified sleep restriction for insomnia in general practice: a randomised controlled trial. *Br J Gen Pract* 2015;**65**:e508–15. doi:10.3399/bjgp15X686137.
66. Murawski B, Plotnikoff RC, Rayward AT, Oldmeadow C, Vandelanotte C, Brown WJ, et al. Efficacy of an m-Health physical activity and sleep health intervention for adults: a randomized waitlist-controlled trial. *Am J Prev Med* 2019;**57**:503–14.
67. Arora T, Grey I. Health behaviour changes during COVID-19 and the potential consequences: a mini-review. *J Health Psychol* 2020;**25**:1155–63.
68. Blume C, Schmidt MH, Cajochen C. Effects of the COVID-19 lockdown on human sleep and rest-activity rhythms. *Curr Biol* 2020;**30**:R795–7. doi:10.1016/j.cub.2020.06.021.
69. Moore LV, Dodd KW, Thompson FE, Grimm KA, Kim SA, Scanlon KS. Using behavioral risk factor surveillance system data to estimate the percentage of the population meeting US department of agriculture food patterns fruit and vegetable intake recommendations. *Am J Epidemiol* 2015;**181**:979–88.
70. van Drongelen A, Boot CR, Hlobil H, Twisk JW, Smid T, van der Beek AJ. Evaluation of an mHealth intervention aiming to improve health-related behavior and sleep and reduce fatigue among airline pilots. *Scand J Work Environ Health* 2014;**40**:557–68.
71. Robroek SJ, van Lenthe FJ, van Empelen P, Burdorf A. Determinants of participation in worksite health promotion programmes: a systematic review. *Int J Behav Nutr Phys Act* 2009;**6**:26. doi:10.1186/1479-5868-6-26.
72. Brutus S, Aguinis H, Wassmer U. Self-reported limitations and future directions in scholarly reports: analysis and recommendations. *J Manage* 2013;**39**:48–75.