

---

## Literature Cited

---

- Abdus, S.A. and Ross L.G. 2000. Optimizing site selection for development of shrimp (*Penaeus monodon*) and mud crab (*Scylla serrata*) culture in southwestern Bangladesh. *Proceedings of GIS'2000*, 14<sup>th</sup> Annual conference of Geographic Information Systems. Toronto, Canada, March 2000. no pagenumbers.
- Agostini, V. N., and Bakun, A. 2002. 'Ocean triads' in the Mediterranean Sea: physical mechanisms potentially structuring reproductive habitat suitability (with example application to European anchovy, *Engraulis encrasicolus*). *Fisheries Oceanography*, 11(3), 129–142.
- Aguilar-Manjarrez, J., 1996. Development and evaluation of GIS-based models for planning and management of coastal aquaculture: a case study in Sinaloa, Mexico. PhD Thesis. Institute of Aquaculture, University of Stirling, Scotland, UK, 375p.
- Aguilar-Manjarrez, J. and Ross, L.G. 1995a. GIS enhances aquaculture development. *GIS World* 8(3), 52-56.
- Aguilar-Manjarrez, J. and Ross, L.G. 1995b. Geographical information system (GIS) environmental models for aquaculture development in Sinaloa State, Mexico. *Aquaculture International*, 3, 103-115.
- Alfaro A.C., Jeffs, A.G., and Hooker, S.H. 2001. Reproductive behaviour of the green-lipped mussel, *Perna canaliculus*, in northern New Zealand. *Bulletin of Marine Science*, 69(3), 1095-1108.
- Ali, C.Q., Ross, R.G, and Beveridge, M.C.M., 1991. Microcomputer spreadsheets for the implementation of GIS in aquaculture: a case study on carp in Pakistan. *Aquaculture*, 92(2-3), 199-205.
- Allen, J.S., and Kundu, P.K. 1978. On the momentum, vorticity and mass balance on the Oregon shelf. *Journal of Physical Oceanography*, 8, 13-27.
- Allen, J.S., Newburger, P.A., and Federiuk, J. 1995. Upwelling circulation on the Oregon continental shelf. Part I: Response to idealized forcing. *Journal of Physical Oceanography*, 25, 1843-1866.
- American Public Health Association (APHA), 1980. Greenburg, A.E., Conners, J.J., and Jenkins, D. (eds), *Standard methods for the examination of water and wastewater*, 15<sup>th</sup> ed, American Water Works Association; Water Pollution Control Federation, Washington, D.C., 1134p.
- American Public Health Association (APHA), 1998. Eaton, A.D., Clesceri, L. S., and Greenberg, A.E (eds), *Standard methods for the examination of water and wastewater*, 20th ed, American Water Works Association; Water Pollution Control Federation, Washington, D.C., 1325p.
- American Public Health Association (APHA), 2005. Eaton, A.D., Clesceri, L. S., Rice, E.W., and Greenberg, A.E (eds), *Standard methods for the examination of water and wastewater*. 21<sup>st</sup> ed, American Water Works Association; Water Pollution Control Federation, Washington, D.C., 1368p.
- Andréfouët, S., Ouillon, S., Brinkman, R., Falter, J., Douillet, P., Wolk, F., Smith, R., Garen, P., Martinez, E., Laurent, V., Lo, C., Remoissenet, G., Scourizic, B., Gilbert, A., Deleersnijder, E., Steinberg, C., Choukron, S., Buestel, D. 2006. Review of solutions for 3D hydrodynamic modeling applied to aquaculture in South Pacific atoll lagoons. *Marine Pollution Bulletin*, 52, 1138-1155.
- Andreu-Burillo, I., Holt, J., Procter, R., Annan, J.D., James, I.D., and Prandle, D. 2007. Assimilation of sea surface temperature in the POL coastal ocean modeling system. *Journal of Marine Systems*, 65, 27-40.
- Arnold, W.S., White, M.W., Norris, H.A., and Berrigan, M.E. 2000. Hard clam (*Mercenaria* spp.) aquaculture in Florida, USA: geographic information system applications to lease site selection. *Aquacultural Engineering*, 23, 203-231.
- Aure, J., Strohmeier, T., and Strand, Ø. 2007. Modelling current speed and carrying capacity in long-line blue mussel (*Mytilus edulis*) farms. *Aquaculture Research*, 38, 304-312.
- Bacher, C., Duarte, P., Ferriera, J.G., Heral, M., and Raillard, O. 1998. Assessment and comparison of the Marennes-Oleron Bay (France) and Carlingford Lough (Ireland) carrying capacity with ecosystem models. *Aquatic Ecology*, 31, 379-394.
- Bacher, C., Grant, J., Hawkins, A.J.S., Fang, J., Zhu, M., and Besnard, M. 2003. Modelling the effect of food depletion on scallop growth in Sungo Bay (China). *Aquatic Living Resources*, 16, 10-24.
- Ballinger, M. M., 2003. *Developing methods for identifying cryptic tephra in peat cores from the Waikato region, North Island, New Zealand.*, Faculty of Science, University of Plymouth, 126p.
- Barber, R.T. and Smith, W.O. 1981. The role of circulation, sinking and vertical migration in physical sorting of phytoplankton in the upwelling center at 15S, *In: Richards, F.A. (ed), Coastal Upwelling*. American Geophysical Union, Washington, pp.366-371.

- Baxter, K.J., 2003. Broad scale classification and prediction of marine habitats: integrating GIS and rule based modelling. In: Woodroffe, C.D., and Furness, R.A. (eds.), *Coastal GIS 2003: an integrated approach to Australian coastal issues*. Wollongong, Australia, pp. 125-137.
- Bayne, B.L. 1993. Feeding physiology of bivalves: time dependence and compensation for changes in food availability. In: Dame, R.F. (ed). *Bivalve filter feeders in estuarine and coastal ecosystems processes*. NATO ASI series vol. G33, Springer Verlag, Berlin, 1-24.
- Bayne, B.L., Thompson, R.J., and Widdows, J. 1976. Physiology: I. In: Bayne, B. L. (ed.) *Marine Mussels: Their Ecology and Physiology*. Cambridge University Press, 121-207.
- Bayne, B.L. and Scullard, C. 1977. Rates of nitrogen excretion by species of *Mytilus* (Bivalvia: Mollusca). *Journal of the Marine Biological Association of the United Kingdom*, 57, 355-369.
- Beamsley, B., Black, K.P., Longdill, P.C., and McComb, P., 2005. Bay of Plenty Current and Temperature Measurements: Aquaculture Management Areas. Report for Environment Bay of Plenty, ASR Ltd, P.O. Box 67, Raglan, NZ, 65p.
- Bell, R.G. 1985. Harmonic analysis of tidal water heights at Moturiki, 1979-85, *Water Quality Centre internal report 85/6*, 13p.
- Bell, R.G., and Goring, D.G. 1996. Techniques for analyzing sea level records around New Zealand, *Marine Geodesy*, 19, 77-98.
- Bell, R.G., and Goring, D.G. 1998. Seasonal variability of sea level and sea-surface temperature on the north-east coast of New Zealand. *Estuarine, Coastal and Shelf Science*, 46, 307-318.
- Benitez-Nelson, C.R., Buesseler, K.O., and Crossin, G. 2000. Upper ocean export, horizontal transport, and vertical eddy diffusivity in the southwestern Gulf of Maine. *Continental Shelf Research*, 20, 707-736.
- Black, K.D. 2001. Sustainability of aquaculture. In: Black, K.D. (ed), *Environmental impacts of aquaculture*. Sheffield Academic Press, Sheffield, U.K., 119-212.
- Black, K.P. 1987. A numerical sediment transport model for application to natural estuaries, harbours and rivers. In: Noye, J. (ed). *Numerical modelling applications to marine systems*. North Holland, Elsevier, pp. 77-105.
- Black, K.P. 1989. Numerical simulation of steady and unsteady meso-scale eddies. Proceedings of the 9th Australasian Conference on Coastal and Ocean Engineering, Adelaide, pp. 204-208.
- Black, K.P. 1995. The hydrodynamic model 3DD and support software, Occasional Report 19, Department of Earth Sciences, University of Waikato, 53p.
- Black, K.P. 2006. Model POL3DD description and users guide, ASR Ltd, Marine and Freshwater Consultants, Raglan, New Zealand, 117p.
- Black, K.P., and Gay, S. L. 1987. Eddy formation in unsteady Flows. *Journal of Geophysical Research*, 92, 9514-9522.
- Black, K.P., Moran, P.J., and Hammond, L.S. 1991. Numerical models show coral reefs can be self seeding. *Marine Ecology Progress Series*, 74, 1-11.
- Black, K.P.; Hatton, D.; Rosenberg, M. 1993. Locally and externally-driven dynamics of a large semienclosed bay in southern Australia. *Journal of Coastal Research*, 9(2), 509-538.
- Black, K.P., Bell, R. G., Oldman, J. W., Carter, G. S., Hume, T. M. 2000. Features of 3-dimensional barotropic and baroclinic circulation in the Hauraki Gulf, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 34, 1-28.
- Blanton, J.O., Tenore, K.R., Castillejo, R., Atkinson, L.P., Schwing, F.B., and Lavin, A. 1987. The relationship of upwelling to mussel production in the rias on the western coast of Spain. *Journal of Marine Research*, 45, 497-511.
- Blatt, H., 1982. *Sedimentary Petrology*. W.H. Freeman and Company, New York, USA, 546p.
- Bowden, K.F. 1983. *Physical oceanography of coastal waters*, Ellis Horwood Ltd., Chichester, U.K., 302p.
- Bowie, G.L., Mills, W.B., Porchella, D.B., Campbell, C.L., Pagenkopf, J.R., Rupp, G.L., Johnson, K.M., Chan, P.W.H., and Gherini, S.A. 1985. Rates, Constants, and Kinetics Formulations in Surface Water Quality Modeling (2nd ed.). Tetra Tech Inc. Lafayette, California. EPA/600/3-85/040 Environmental Research Laboratory, Office of Research and Development, U.S. Environmental Protection Agency, Athens, Georgia. 455p.
- Boyd, A.J. and Heasman, K.G. 1998. Shellfish mariculture in the Benguela system: water flow patterns within a mussel farm in Saldanha Bay, South Africa. *Journal of Shellfish Research*, 17, 25-32.
- Boyd, C.E. and Clay, J.W. 1998. Shrimp aquaculture and the environment. *Scientific American*, 278, 58-65.
- Boyd, C.E. and Schmittou, H.R. 1999. Achievement of sustainable aquaculture through environmental management. *Aquaculture Economics and Management*, 3(1), 59-69.
- Boyer, T.P., Antonov, J.I., Garcia, H.E., Johnson, D.R., Locarnini, R.A., Mishonov, A.V., Pitcher, M.T., Baranova, O.K., and Smolyar, I.V., 2006. *World Ocean Database 2005*. Levitus, S. (ed), NOAA Atlas NESDIS 60, U.S. Government Printing Office, Washington, D.C., 190 pp.

- Bradbury, I.R., and Snelgrove, P.V.R. 2001. Contrasting larval transport in demersal fish and benthic invertebrates: the roles of behaviour and advective processes in determining spatial pattern. *Canadian Journal of Fisheries and Aquatic Sciences*, 58, 811-823.
- Bradford, J.M., Heath, R.A., Chang, F.H., and Hay, C.H. 1982. The effect of warm-core eddies on oceanic productivity off northeastern New Zealand. *Deep-Sea Research*, 29(12A), 1501-1516.
- Bradshaw, B.E., Healy, T.R., Nelson, C.S., Dell, P.M., and de Lange, W.P. 1994. Holocene sediment lithofacies and dispersal systems on a storm dominated, back-arc shelf margin: the east Coromandel coast, New Zealand. *Marine Geology*, 119, 75-98.
- Brando, V.E., Ceccarelli, R., Libralatos, S., Ravagnan, G., 2004. Assessment of environment management effects in a shallow water basin using mass balance models. *Ecological Modelling*, 172, 213-232.
- Briere, P.R., Scanlon, K.M., Gledhill, C.T., Koenig, C.C., and Fitzhugh, G., 1999. West Florida Shelf: Sidescan Sonar and Sediment Data from Shelf Edge Habitats in the Northeastern Gulf of Mexico., *U.S. Geological Survey Open-File Report 99-589*. U.S. Geological Survey, Woods Hole, MA 02543
- Brink, K.H., Jones, B.J., van Leen, J.C., Mooers, C.K.N., Stuart, D.W., Stevenson, M.R., Dugdale, R.C., and Heburn, G.W. 1981. Physical and biological structure and variability in an upwelling center off Peru near 15°S during March, 1977, *In: Richards, F.A. (ed), Coastal Upwelling*, American Geophysical Union, Washington, pp.473-495.
- Broekhuizen, N., Zeldis, J., Stephens, S.A., Oldman, J.A., Ross, A.H., Ren, J., and James, M.R. 2002. Factors relating to the sustainability of shellfish aquaculture operations in the Firth of Thames: a preliminary analysis, NIWA report, Hamilton, New Zealand, 106p.
- Buchwald, V.T., and Kachoyan, B.J. 1987. Shelf waves generated by a coastal flux. *Australian Journal of Marine and Freshwater Research*, 38, 429-437
- Burbidge, P.R., Gavine, F.M., and Kelly, L.A. 1993. Control and regulation of the environmental impact of fish farming in Scotland. *In: Proceedings of a workshop on fish farm effluents and their control in EC countries*, Hamburg, 23-25 November 1992. no page numbers
- Burbidge, P., Hendrick, V., Roth, E., and Rosenthal, H. 2001. Social and economic policy issues relevant to marine aquaculture. *Journal of Applied Ichthyology*, 17, 194-206.
- Buschmann, A.H., Lopez, D.A., and Medina, A., 1996. A review of the environmental effects and alternative production strategies of marine aquaculture in Chile. *Aquacultural Engineering*, 15(6), 397-421.
- Butler, E.I., Corner, E.D., and Marshall, S.M. 1969. On the nutrition and metabolism of zooplankton VI. Feeding efficiency of *Calanus* in terms of nitrogen and phosphorus. *Journal of the Marine Biological Association of the United Kingdom*, 49, 977-1001.
- Bye, J.A.T., and Heath, R.A. 1975. The New Zealand semi-diurnal tide. *Journal of Marine Research*, 33, 423-442.
- Campbell, D.E. and Newell, C.R. 1998. MUSMOD, a production model for bottom culture of the blue mussel, *Mytilus edulis* L. *Journal of Experimental Marine Biology and Ecology*, 219, 171-203.
- Caputi, N., Chubb, C., and Pearce, A. 2001. Environmental effects on recruitment of the western rock lobster, *Panulirus Cygnus*. *Marine and Freshwater Research*, 52, 559-582.
- Carrick, N.A., and Ostendorf, B. 2007. Development of a spatial decision support system (DSS) for the Spencer Gulf penaid prawn fishery, South Australia. *Environmental Modelling and Software*, 22, 137-148.
- Carver, C.E.A. and Mallet, A.L. 1990. Estimating the carrying capacity of a coastal inlet for mussel culture. *Aquaculture*, 88, 39-53.
- Chamberlain, J., Fernandes, T.F., Read, P., Nickell, T.D., and Davies, I.M., 2001. Impacts of biodeposits from suspended mussel (*Mytilus edulis* L.) culture on the surrounding surficial sediments. *ICES Journal of Marine Science*, 58, 411-416.
- Chang, F.H., Vincent, W.F., and Woods, P.H. 1992. Nitrogen utilization by size-fractionated phytoplankton assemblages associated with an upwelling event. *New Zealand Journal of Marine and Freshwater Research*, 26, 287-301.
- Chang, F.H., MacKensie, L., Till, D., Hannah, D., and Rhodes, L. 1995. The first toxic shellfish outbreaks and the associated phytoplankton blooms in early 1993 in New Zealand. *In: Lassus, P., Arul, G., Erad, E., Gentein, P., and Marcallou, C. (Eds), Harmful marine algal blooms, Proceedings of the 6<sup>th</sup> International Conference on Toxic Marine Phytoplankton*, Lavoisier, Intercept Ltd, pp.145-150.
- Chang, F.H., Sharples, J., and Grieve, J.M. 1996. Temporal and spatial distribution of toxic dinoflagellates in Bay of Plenty, New Zealand, during the early 1993 toxic shellfish outbreaks. *In: Yasumoto, T., Oshima, Y., and Fukuyo, Y. (eds), Harmful and Toxic Algal Blooms*. Intergovernmental Oceanographic Commission of UNESCO 1996, pp. 235-238.

- Chang, F.H., Zeldis, J., Gall, M., and Hall, J. 2003. Seasonal and spatial variation of phytoplankton assemblages, biomass and cell size from spring to summer across the north-eastern New Zealand continental shelf. *Journal of Plankton Research*, 25(7), 737-758.
- Chapelle, A., Menesguen, A., Deslous-Paoli, J.-M., Souchu, P., Mazouni, N., Vaquer, A., and Millet, B., 2000. Modelling nitrogen, primary production and oxygen in a Mediterranean lagoon. Impact of oysters farming and inputs from the watershed. *Ecological Modelling*, 127, 161-181.
- Chiswell, S.M. 2000. Tidal energetics over the Chatam Rise, New Zealand. *Journal of Physical Oceanography*, 30, 2452-2460.
- Chiswell, S.M. 2001. Determining the internal structure of the ocean off the north-east coast of New Zealand from surface measurements. *New Zealand Journal of Marine and Freshwater Research*, 35, 289-306.
- Chiswell, S.M., and Moore, M.I. 1999. Internal tides in the south-west Pacific Ocean near the Kermadec Ridge. *Journal of Physical Oceanography*, 29, 1019-1035.
- Chiswell, S.M., and Schiel, D.R. 2001. Influence of along-shore advection and upwelling on coastal temperature at Kaikoura Peninsula, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 35, 307-317.
- Chivilev, S., and Ivanov, M., 1997. Response of the arctic benthic community to excessive amounts of non-toxic organic matter. *Marine Pollution Bulletin*, 35(7-12), 280-286.
- Cole, R. 2002. Impacts of marine farming on wild fish populations, NIWA report, June 2002, 49p.
- Cole, R., N. Alcock, G. Carbins and R. Stewart, 2003. Cape Runaway: Biological Information Relevant to a Potential Marine Reserve Proposal. Report prepared for Department of Conservation, i no. NEL2003-009.
- Collos, Y., Vaquer, A., Troussellier, M., Lautier, J., Chretiennot-Dinet, M.J., Neveux, J., and Machado, C. 1997. Variability in nitrate uptake rate of phytoplankton communities in a Mediterranean coastal lagoon. *Estuarine, Coastal, and Shelf Science*, 44, 369-375.
- Condie, S.A., Waring, J., Mansbridge, J.V. and Cahill, M.L. 2005. Marine connectivity patterns around the Australian continent, *Environmental Modelling and Software*, 20, 1149-1157.
- Conover, R.J. 1966. Assimilation of organic matter by zooplankton. *Limnology and Oceanography*, 11, 338-354.
- Costa, K.G.d, and Nalesso, R.C., 2006. Effects of mussel farming on macrobenthic community structure in Southeastern Brazil. *Aquaculture*, 258, 655-663.
- Cowen, R.K., Lwiza, K.M.M., Sponaugle, S., Paris, C.B., and Olson, D.B. 2000. Connectivity of marine populations: open or closed?. *Science*, 287, 857-859.
- Cowen, R.K., Paris, C.B. and Srinivasan, A. 2006. Scaling of connectivity in marine populations. *Science*, 311, 522-527.
- Cox, M.D. 1970. A mathematical model of the Indian Ocean. *Deep-Sea Research*, 17, 45-75.
- Cox, M.D. 1975. A baroclinic numerical model of the world ocean: Preliminary Results, In: *Numerical models of Ocean Circulation*, National Academy of Sciences, Washington D.C. pp.107-120.
- Cranford, P., Dowd, M., Grant, J., Hargrave, B., and McGladdery, S., 2003. Ecosystem level effects of marine bivalve aquaculture. In: *A scientific review of the potential environmental effects of aquaculture in aquatic ecosystems* (Vol. 1), Canadian Technical Report of Fisheries and Aquatic Sciences, 2450, pp. 51-84.
- Crawford, C.M., 2003. Environmental management of marine aquaculture in Tasmania, Australia. *Aquaculture*, 226, 129-138.
- Crawford, C.M., Macleod, C.K.A., Mitchell, I.M., 2003. Effects of shellfish farming on the benthic environment. *Aquaculture*, 224, 117-140.
- Crèpon, M., Richez, C., and Chartier, M. 1984. Effects of coastline geometry on upwellings. *Journal of Physical Oceanography*, 14, 1365-1382.
- Cross, S.F., and Kingzett, B.C., 1992. Biophysical criteria for shellfish culture in British Columbia: a site capability evaluation system. Aquametrix Research report for the British Columbia Ministry of Agriculture, Fisheries and Food, 40p.
- Csanady, G.T. 1981. Circulation in the coastal ocean. In: Saltzman, B. (ed.), *Advances in Geophysics*, Academic Press, New York, vol. 23, pp.101-183.
- Csanady, G.T. 1982. On the structure of transient upwelling events. *Journal of Physical Oceanography*, 12, 84-96.
- Csanady, G.T. 1997. On the theories that underlie our understanding of continental shelf circulation. *Journal of Physical Oceanography*, 53, 207-229.
- Cugier, P., Menesguen, A., and Guillaud, J.F. 2005. Three-dimensional (3D) ecological modelling of the Bay of Seine (English Channel, France). *Journal of Sea Research*, 54, 104-124.
- Cushman-Roisin, B., O'Brien, J.J., and Smith, R.L. 1983. On wind and ocean-velocity correlations in a coastal-upwelling system. *Journal of Physical Oceanography*, 13(3), 547-550.
- Dahlback, B., and Gunnarsson, L.A.H., 1981. Sedimentation and sulfate reduction under a mussel culture. *Marine Biology*, 63, 269-275.

- Dame, R.F., Dankers, N., Prins, T., Jongsma, H., and Smaal, A. 1991. The influence of mussel beds on nutrients in the West Wadden Sea and Eastern Scheldt estuaries. *Estuaries*, 14, 130-138.
- Dame, R.F. and Prins, T.C. 1998. Bivalve carrying capacity in coastal ecosystems, *Aquatic Ecology*, 31, 409-421.
- Dane, J.H., and Topp, G.C., eds., 2002, *Methods of Soil Analysis, Part 4*, Physical Methods. Madison, WI, Soil Science Society of America, Soil Science Society of America Book Series Number 5, ISBN 0-89118-810-X, 1692p
- Davenport, J., Smith, R.J.J.W., Packer, M. 2000 Mussels *Mytilus edulis*: significant consumers and destroyers of mesozooplankton. *Marine Ecology Progress Series*, 198, 131–137.
- Davies, A.M., and Hall, P. 2002. Numerical problems associated with coupling hydrodynamic models in shelf edge regions: the surge event of February 1994. *Applied Mathematical Modelling*, 26, 807-831.
- Davies, A.M., and Xing, J. 2003. On the interaction between internal tides and wind-induced near-inertial currents at the shelf edge. *Journal of Geophysical Research*, 108 (C3), 3099, doi:10.1029/2002JC001375.
- Day, P.R. 1965. Particle fraction and particle-size analysis, pp. 545-566. In C.A. Black et al. (eds.), *Methods of soil analysis*, Part 1. Agronomy No. 9. Madison, WI: American Society of Agronomy.
- de Jong, R.J. 1994. The effects of mussel farming on the benthic environment. MSc thesis, University of Auckland, Auckland, New Zealand, 150p.
- de Lange, W.P. 1991. Wave climate for No. 1 reach, Port of Tauranga, Tauranga Harbour. Report for the Port of Tauranga Ltd, September, 1991. Marine Geosciences Group, University of Waikato, New Zealand, 18p.
- de Lange, W.P., Bell, R., Gorman, R., and Reid, S. 2003. Physical oceanography of New Zealand waters. In: Goff, J., Nichol, S.L., and Rouse, H.L. (eds), *The New Zealand Coast Te Tai o Aotearoa*, Dunmore Press, Palmerston North, New Zealand, pp.59 – 77.
- Dean, W.E., 1974. Determination of Carbonate and Organic Matter in Calcareous Sediments and Sedimentary Rocks by Loss on Ignition: Comparison with Other Methods. *Journal of Sedimentary Petrology*, 44(1), 242-248.
- Demarcq, H., and Faure, V. (2000), Coastal upwelling and associated retention indices derived from satellite SST. Application to *Octopus vulgaris* recruitment. *Oceanologica Acta*, 23( 4), 391-408.
- Derber, J., and Rosati, A. 1989. A global oceanic data assimilation system. *Journal of Physical Oceanography*, 19(9), 1333-1347.
- Dever, E.P. 1997. Wind-forced cross-shelf circulation on the northern California shelf. *Journal of Physical Oceanography*, 14, 1566-1580.
- DHI, 2004. DHI eutrophication model 2 – ECOLab template: a scientific description, DHI water and environment, Horsholm, Denmark, 38p.
- Doonan, D. 2001. Aquaculture in the ages of integrated coastal management (ICM). In: Black, K.D. (ed), *Environmental impacts of aquaculture*. Sheffield Academic Press, Sheffield, U.K., 182-198.
- Dowd, M. 1997. On predicting the growth of cultured bivalves. *Ecological modelling*, 104, 113-131.
- Dowd, M. 2005. A bio-physical coastal ecosystem model for assessing environmental effects of marine bivalve aquaculture, *Ecological Modelling*, 183, 323-346.
- Doyle, A.C., Carter, L., Glasby, G.P., and Lewis, K.B. 1979. Bay of Plenty Sediments. *N.Z. Oceanographic Institute Chart*, Coastal Series, 1:200,000.
- Duarte, P., Meneses, R., Hawkins, A.J.S., Zhu, M., Fang, J., Grant, J., 2003. Mathematical modelling to assess the carrying capacity for multi-species culture within coastal waters. *Ecological Modelling*, 168, 109– 143.
- Dubois, J.M. and Habbane, M. 2002. Methodology of Regional aquaculture zoning using the aid of teledetection in IRS. *The Canadian Geographer* 46(2), 172-185.
- Duffy, C., 1998. Classification of New Zealand Coastal Units: Comment on East Coast Hawke's Bay Conservancy (Eastern Bay of Plenty: Opape to Matakaoa Point). Department of Conservation Internal Correspondence from Clinton Duffy, 26 October 1998.
- Dugdale, R.C. 1967. Nutrient limitation in the sea: dynamics, identification, and significance. *Limnology and Oceanography*, 12(4), 685-695.
- Dunbar, G.B., and Barret, P.J., 2005. Estimating paleobathymetry of wave-graded continental shelves from sediment texture. *Sedimentology*, 52, 253-269.
- Dyke, P.P.G. 2001. *Coastal and shelf sea modelling*, Springer Verlag, 272p.
- EBOP, 2006, Coastal use and value maps and discussion document: Aquaculture Management Areas, Environment Bay of Plenty (EBOP), Whakatane, New Zealand, 31p + 12 maps.
- Edwards C.A., Powell T.A., and Batchelder H.P. 2000. The stability of an NPZ model subject to realistic levels of vertical mixing. *Journal of Marine Research*, 58(1), 37-60.
- Egbert, G.D., Bennet, A.F., and Foreman, M.G.G. 1994. TOPEX/Poseidon tides estimated using a global inverse model. *Journal of Geophysical Research*, 99(C12), 24,821-24,852.

- Egbert, G.D., and Erofeeva, S.Y. 2002. Efficient inverse modeling of barotropic ocean tides. *Journal of Oceanic and Atmospheric Technology*, 19(N2), 183-204.
- Emery, W.J. and Thomson, R.E. 2004. *Data analysis methods in physical oceanography* (2<sup>nd</sup> rev. ed.), Elsevier B.V., Amsterdam, The Netherlands, 638p.
- Emmerson, C.W. 1990. Influence of sediment disturbance and water flow on the growth of the soft-shell clam *Mya arenaria* L. *Canadian Journal of Fisheries and Aquatic Science*, 47, 1655-1663.
- Eppley, R.W., 1972. Temperature and phytoplankton growth in the sea. *Fisheries Bulletin*, 70, 1063-1085.
- Eppley, R. W., Rogers, J. N., and McCarthy, J. 1969. Half-saturation constants for uptake of NO<sub>3</sub> and NH<sub>4</sub> by marine phytoplankton. *Limnology and Oceanography*, 14, 912-920.
- Eppley, R.W., Rogers, J.N., McCarthy, J.J., and Sournia, A. 1971. Light/dark periodicity in nitrogen assimilation of the marine phytoplankters *Skeletonema costatum* and *Coccolithus Huxley* in N-limitant chemostat culture. *Journal of Phycology*, 7, 150-154.
- Espinosa-Carreón, T.L., Strub, P.T., Beier, E., Ocampo-Torres, F., and Gaxiola-Castro, G. 2004. Seasonal and interannual variability of satellite-derived chlorophyll pigment, surface height, and temperature off Baja California. *Journal of Geophysical Research*, 109 (C3), C03039 doi:10.1029/2003JC002105.
- Evans, S.R. and Middleton, J.F. 1998. A regional model of shelf circulation near Bass Strait: A new upwelling mechanism. *Journal of Physical Oceanography*, 28, 1439-1457.
- FAO. 2002. The state of world fisheries and aquaculture, (SOFIA), Food and Agriculture Organisation of the United Nations, Rome, Italy, 150p.
- Ferreira, J.G., Duarte, P., and Ball, B. 1998. Trophic capacity of Carlingford Lough for oyster culture – analysis by ecological modelling. *Aquatic Ecology*, 31, 361–378.
- Feuillet-Girard, M., Héral, M., Abrioux, M., Fontugne, M., 1994. Carbone organique dissous et particulaire de la colonne d'eau et de l'interface eau-sédiment du bassin de Marennes-Oléron: influence des huîtres. *Oceanologica Acta*, 17, 271–284 (in French).
- Figueiras, F.G., Labarta, U., and Fernandez Reriz, M.J. 2002. Coastal upwelling, primary production and mussel growth in the Rias Baixas of Galicia. *Hydrobiologia*, 484, 121-131.
- Flaws, D.E. 1975. Aspects of the biology of mussels in the Cook Strait area. Unpublished PhD thesis, Victoria University, Wellington, New Zealand, 219p.
- Flemming, C.A. 1959. Notes on the New Zealand recent and tertiary mussels (Mytilidae). *Transactions of the Royal Society of New Zealand*, 87, 165-178.
- Frankic, A. and Hershner, C. 2003. Sustainable aquaculture: developing the promise of aquaculture. *Aquaculture International*, 11, 517-530.
- Franks, P.J.S., Wroblewski, J.S., and Flierl, G.R. 1986. Behaviour of a simple phytoplankton model with food-level acclimation by herbivores. *Marine Biology*, 91(1), 121-129
- Frechette, M., Bourget, E., 1985. Energy flow between the pelagic and benthic zones: factors controlling particulate organic matter available to inter-tidal mussel bed. *Canadian Journal of Fisheries and Aquatic Science*, 42, 1158-1165.
- Freeland, H.J., Boland, F.M., Church, J.A., Clarke, A.J., Forbes, A.M.G., Huyer, A., Smith, R.L., Thompson, R.O.R.Y., and White, N.J. 1986. The Australian coastal experiment: A search for coastal-trapped waves. *Journal of Physical Oceanography*, 16, 1230-1249.
- Fridley, R.B. 1995. The opportunities for engineering and technology in addressing the environmental, institutional, and economic constraints of marine aquaculture in the United States. *Estuaries* 18(1A), 18-24.
- Fulton, E.A., Smith, A.D.M. and Johnson, C.R. 2003. Effect of complexity on marine ecosystem models. *Marine Ecology Progress Series*, 253, 1-16.
- Gardner, J.P.A. 2002. Effects of seston variability on the clearance rate and absorption efficiency of the mussels *Aulacomya maoriana*, *Mytilus galloprovincialis* and *Perna canaliculus* from New Zealand. *Journal of Experimental Marine Biology and Ecology*, 268, 83-101.
- Garner, D.M. 1959. The subtropical convergence in New Zealand surface waters. *New Zealand Journal of Geology and Geophysics*, 2, 315-337.
- Garner, D.M. 1961. Hydrology of New Zealand coastal waters, 1955. *New Zealand Oceanographic Institute Memoir* no. 18, 85pp.
- Garvine, R.W. 1971. A simple model of coastal upwelling dynamics. *Journal of Physical Oceanography*, 1, 169-179.
- GESAMP (IMO/FAO/UNESCO-IOC/WMO/WHO/IAEA/UN/UNEP Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection). 1996. The contributions of science to coastal zone management. GESAMP Reports and Studies no.61, 66 p.
- Ghil, M., and Malanotte-Rizzoli, P. 1991. Data assimilation in meteorology and oceanography. *Advances in Geophysics*, 33, 141-266.
- Gibbs, M.M., James, M.R., Pickmere, S.E., Woods, P.H., Shakespeare, B.S., Hickman, R.W., and Illingworth, J. 1991. Hydrodynamic and water column properties at six stations associated with

- mussel farming in Pelorus Sound, 1984-85. *New Zealand Journal of Marine and Freshwater Research*, 25, 239-254.
- Gibbs, M.M., Pickmere, S.E., Woods, P.H., Payne, G.W., James, M.R., Hickman, R.W., and Illingworth, J. 1992. Nutrient and chlorophyll a variability at six stations associated with mussel farming in Pelorus Sound. *New Zealand Journal of Marine and Freshwater Research*, 26, 197-211.
- Gibbs, M.T., 2004. Interactions between bivalve shellfish farms and fishery resources. *Aquaculture*, 240, 367-396.
- Gibbs, M.T. 2007. Sustainability performance indicators for suspended bivalve aquaculture activities. *Environmental Indicators*, 7, 94-107.
- Goldman, J.C. and Caron, D.A. 1985. Experimental studies on an omnivorous microflagellate. Implications for grazing and nutrient regeneration in the marine microbial food chain. *Deep Sea Research*, 32, 889-915.
- Goring, D.G. 1995. Short-term variations in sea level (2-15 days) in the New Zealand region. *New Zealand Journal of Marine and Freshwater Research*, 29, 69-82.
- Goring, D.G., and Walters, R.A. 2002. Ocean-tide loading and Earth tides around New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 36, 299-309.
- Grange, K., and Cole, R., 1997. Mussel farming impacts. *Aquaculture Update*, (NIWA, NZ), 19(4), 1-3.
- Grant, J., Hatcher, A., Scott, D.B., Pocklington, P., Schafer, C.T., and Winter, G. 1995. A multidisciplinary approach to evaluating benthic impacts of shellfish aquaculture. *Estuaries*, 18, 124-144.
- Grant, J., Stenton-Dozey, J., Monteiro, P., Pitcher, G., and Heasman, K. 1998. Shellfish culture in the benguela upwelling system: a carbon budget of Saldanha Bay for raft culture of *Mytilus galloprovincialis*. *Journal of Shellfish Research*, 17(1), 41-49.
- Grant, J. and Bacher, C. 2001. A numerical model of flow modification induced by suspended aquaculture in a Chinese bay. *Canadian Journal of Fisheries and Aquatic Science*, 58, 1003-1011.
- Grant, J. Curran, K.J., Guyondet., T.L., Tita, G., Bacher, C., Koutitonsky, V., and Dowd, M. 2007. A box model of carrying capacity for suspended mussel aquaculture in Lagune de al Grade-Entrée, Iles-de-la-Madeleine, Quebec. *Ecological Modelling*, 200, 193-206.
- Grieg, M.J., Ridgeway, N.M., and Shakespeare, B.S. 1988. Sea surface temperature variations at coastal sites around New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 22, 391-400.
- Hall, J.A., Safi, K., James, M.R., Zeldis, J., and Weatherhead, M. 2006. Microbial assemblage during the spring-summer transition on the northeast continental shelf of New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 40, 195-210.
- Hampson, G., Hoagland, P., Kite-Powell, H., and Paul, W., 1999. Submerged coastal offshore mussel aquaculture system (SCOMAS), Poster displayed at Rinehart coastal research center, Woods Hole Oceanographic Institute, June 11, 1999.
- Handley, S. and Jeffs, A. 2003. Assessment of future expansion of Pacific oyster (*Crassostrea gigas*) farming in Northland. NIWA Client Report AKL2003-027, 32p.
- Hardman-Mountford, N.J., and McGlade, J.M. 2003. Seasonal and interannual variability of oceanographic processes in the Gulf of Guinea: an investigation using AVHRR sea surface temperature data. *International Journal of Remote Sensing*, 24(16), 3247-3268.
- Harrison, S., Lee, R., and Black, K. 2007. An integrated catchment and receiving model for Port Phillip Bay and Western Port to address pressing environmental concerns. *Coasts and Ports 2007 conference*, Adelaide, Australia, paper no. 77.
- Hartstein, N.D., and Rowden, A.A., 2004. Effects of biodeposits from mussel culture on macroinvertebrate assemblages at sites of different hydrodynamic regime. *Marine Environmental Research*, 54, 339-357.
- Hasumi, H., and Sugimoto, N. 1999. Sensitivity of a global ocean circulation model to tracer advection schemes. *Journal of Physical Oceanography*, 29, 2730-2740.
- Hatcher, A., Grant, J., and Schofield, B., 1994. Effects of suspended mussel culture (*Mytilus* spp.) on sedimentation, benthic respiration and sediment nutrient dynamics in a coastal bay. *Marine Ecology Progress Series*, 115, 219-235.
- Hatton, S., Hayden, B.J., and James, M.R. 2005. The effects of food concentration and quality on the feeding rates of three size classes of the greenshell mussel, *Perna canaliculus*. *Hydrobiologia*, 548, 23-32.
- Hawkins, A.J.S., James, M.R., Hickman, R.W., Hatton, S., and Weatherhead, M. 1999. Modelling of suspension-feeding and growth in the green-lipped mussel *Perna canaliculus* exposed to natural and experimental variations of seston ability in the Marlborough Sounds, New Zealand. *Marine Ecology Progress Series*, 191, 217-232.
- Hay, D.N., de Lange, W.P., and Healy, T.R. 1991. Storm and oceanographic databases for the western Bay of Plenty, In: Bell, R.G., Hume, T.M., and Healy, T.R., (eds), Coastal Engineering – climate

- for change, *Proceedings of the 10<sup>th</sup> Australasian conference on coastal and ocean engineering*, Auckland. Water Quality Centre publication 31, pp.174-151.
- Hayden, B.J., 1995. Factors affecting recruitment of farmed greenshell mussels, *Perna canaliculus* (Gmelin) 1791, in Marlborough Sounds, unpublished PhD thesis, University of Otago, Dunedin, New Zealand. 168p.
- Healy, T.R., Harms, C., and de Lange, W. 1991. Dredge spoil and inner shelf investigations off Tauranga Harbour, Bay of Plenty, New Zealand. *Coastal Sediments '91*, American Society of Civil Engineers, pp. 2037-2051.
- Healy, T.R., Matthew, J., de Lange, W., and Black K. 1997. Adjustments towards equilibrium of a large flood-tidal delta after a major dredging program, Tauranga Harbour, New Zealand. *Proceedings of the 25<sup>th</sup> International Conference on coastal Engineering*, American Society of Civil Engineers, pp 3284-3294.
- Heasman, K.G., Pitcher, G., McQuaid, C.D. and Hecht, T. 1998. Shellfish mariculture in the Benguela system: food delivery to mussel rafts in a high production environment and the effect of rope spacing on food extraction, growth rate, yield and condition of mussels. *Journal of Shellfish Research*, 17, 33-39.
- Heath, R.A. 1984a. Tidal observations on the west coast, South Island. *New Zealand Journal of Marine and Freshwater Research*, 18, 251-261.
- Heath, R.A. 1984b. The depth of the mixed layer as an indicator of oceanic circulation around New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 18, 83-92.
- Heath, R.A. 1985. A review of the coastal physical oceanography of the seas around New Zealand – 1982, *New Zealand Journal of Marine and Freshwater Research*, 19, 79-124.
- Heiri, O., Lotter, A.F., and Lemcke, G., 2001. Loss on Ignition as a Method for Estimating Organic and Carbonate Content in Sediments: Reproducibility and Comparability of Results. *Journal of Paleolimnology*, 25, 101-110.
- Henderson, A., Gamito, S., Karakassis, I., Pederson, P. and Smaal, A. 2001. Use of hydrodynamic and benthic models for managing environmental impacts of marine aquaculture. *Journal of Applied Ichthyology*, 17, 163-172.
- Hickman, R.W. 1991. *Perna canaliculus* (Gmelin) in New Zealand. In: Menzel, W. (ed), *Estuarine and marine bivalve mollusc culture*. CRC press, Boca Raton, Florida, 325-334.
- Hickman, R.W., and Illingworth, J. 1980. Condition cycle of the green-lipped mussel. *Perna canaliculus* in New Zealand, *Marine Biology*, 60, 27-38.
- Hogan, K., 1992. Maraetai Bay Marine Reserve Proposal – A Site Survey and Discussion Paper. Department of Conservation, ISBN 0-478-01381-7.
- Holloway, G. 1986. Estimation of oceanic eddy transports from satellite altimetry, *Nature*, 323(6085), 243-244.
- Hopkins, G., and B. Robertson, 2001. Site Assessment for Proposed Marine Farm at Opotiki, Bay of Plenty. Part 1: Benthic Assessment. *Cawthron Report Number 672*. Prepared for Eastern Seafarms Limited, August 2001. 34p.
- Hsueh, Y., and O'Brien, J.J. 1971. Steady coastal upwelling induced by alongshore current. *Journal of Physical Oceanography*, 1, 180-186.
- Hume, T.M., Bell, R.G., de Lange, W.P., Healy, T.R., Hicks, D.M., and Kirk, R.M. 1992. Coastal oceanography and sedimentology in New Zealand, 1967-91. *New Zealand Journal of Marine and Freshwater Research*, 26, 1-36.
- Hume, T.M., Oldman, J.W., Black, K.P. 2000. Sediment facies and pathways of sand transport about a large deep water headland, Cape Rodney, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 34, 695-717.
- Huntsman, S.A. and Barber, R.T. 1977. Primary production off north-west Africa: the relationship to wind and nutrient conditions. *Deep Sea Research*, 24, 25-33.
- Huthnance, J.M. 1995. Circulation, exchange and water masses at the ocean margin: the role of physical processes at the shelf edge. *Progress in Oceanography*, 35, 353-431.
- Huyer, A. 1984. Hydrographic observations along the CODE central line of Northern California, 1981. *Journal of Physical Oceanography*, 14, 1647-1658.
- Huyer, A. 1990. Shelf Circulation, In: Le Mehaute, B., and Hanes, D.M. (eds), *The sea: ideas and observations on progress in the study of the seas*, *Ocean Engineering Science* 9 (Part A), New York, John Wiley and Sons, pp.423-466.
- Incze, L.S., Lutz, R.A., and True, E. 1981. Modeling carrying capacities for bivalve molluscs in open, suspended-culture systems. *Journal of the World Mariculture Society*, 12(1), 143-155.
- Inglis, G.J., Hayden, B.J., and Ross, A.H., 2000. An overview of factors affecting the carrying capacity of coastal embayments for mussel culture, Report for Ministry for the Environment, NIWA client report CHC00/69, 31p.
- Ivlev, V.S. 1961. *Experimental ecology of the feeding fishes*, Yale Univeristy Press, New Haven, Conneticut, 302p.



- Jahnke, R., Richards, M., Nelson, J., Robertson, C., Rao, A., and Jahnke, D. 2005. Organic matter remineralization and porewater exchange rates in permeable South Atlantic Bight continental shelf sediments. *Continental Shelf Research*, 25, 1433-1452.
- James, M.K., Armsworth, P.R., Mason, L.B., and Bode, L. 2002. The structure of reef fish metapopulations: modeling larval dispersal and retention patterns. *Proceedings of the Royal Society B: Biological Sciences*, 269, 2079-2086.
- James, M.R., and Ross, A. 1996. How many mussels can we farm? *Seafood New Zealand*, 4, 50-53.
- James, M.R., Weatherhead, M.A., and Ross, A.H. 2001. Size-specific clearance, excretion, and respiration rates, and phytoplankton selectivity for the mussel *Perna canaliculus* at low levels of natural food. *New Zealand Journal of Marine and Freshwater Research*, 35, 73-86.
- Jassby, A.D. and Goldman, C.R. 1974. Loss rates from a lake phytoplankton community. *Limnology and Oceanography*, 21, 540-547.
- Jeffs, A.G., Holland, R.C., Hooker, S.C., and Hayden, B.J. 1999. Overview and bibliography of research on the greenshell mussel, *Perna canaliculus*, from New Zealand waters. *Journal of Shellfish Research*, 18(2), 347-360.
- Jiang, W. and Gibbs, M.T. 2005. Predicting the carrying capacity of bivalve shellfish culture using a steady, linear food web model. *Aquaculture*, 244, 171-185.
- Jordan, A., Lawler, M., and Halley, V., 2003. The application of seabed habitat mapping for multiple marine resource management in Tasmanian coastal waters. In: Woodroffe, C.D., and Furness, R.A. (eds.), *Coastal GIS 2003: an integrated approach to Australian coastal issues*. Wollongong, Australia, pp. 158-174.
- Jorgensen, S.E., Nielsen, S., and Jorgensen, L. 1991. *Handbook of ecological parameters and ecotoxicology*, Elsevier, Amsterdam, 1263p.
- Kaiser, M.J. 2001. Ecological effects of shellfish cultivation. In: Black, K.D. (ed), *Environmental impacts of aquaculture*. Sheffield Academic / CRC Press, Sheffield, U.K. pp.51-75.
- Kaiser, M.J., Laing, I., Utting, S.D., and Burnell, G.M., 1998. Environmental impacts of bivalve mariculture. *Journal of Shellfish Research*, 17(1), 59-66.
- Kampf, J., Doubell, M., Griffin, D., Matthews, R.L., and Ward, T.M., 2004. Evidence of a large seasonal coastal upwelling system along the southern shelf of Australia, *Geophysical Research Letters*, 31(9), pL09310.
- Kapetsky, J.M. 1994. A Strategic Assessment of Warm Water Fish Farming Potential in Africa. CIFA Technical Paper No.27, Food and Agriculture Organisation of the United Nations, 67pp
- Kapetsky, J.M., McGregor, L., and Nanne, E.H., 1987. A geographical information system and satellite remote sensing plan for aquaculture development: A FAO-UNEP/GRID cooperative study in Costa Rica. *FOA Fisheries Technical Paper No.287*. FAO Rome, 51pp.
- Kapetsky, J.M., Hill, J.M., Worthy, L.D., and Evans, D.L. 1990. Assessing potential for aquaculture development with a geographic information system. *Journal of the World Aquaculture Society* 21(4), 241-249.
- Kapetsky, J.M. and Nath, S.S. 1997. A strategic assessment of the potential for freshwater fish farming in Latin America. *COPESCAL Technical Paper No.10*, FAO, Rome, 128pp.
- Kapetsky, J.M., Hill, J.M., and Worthy, L.D., 1988. A geographical information system for catfish farming development. *Aquaculture* 68, p311-320.
- Karr, J.R. 1981. Assessment of biotic integrity using fish communities, *Fisheries*, vol. 6, 21-27.
- Kaspar, H.F., Gillespie, P.A., Boyer, I.C., MacKenzie, A.L. 1985. Effects of mussel aquaculture on the nitrogen cycle and benthic communities in Kenepuru sound, Marlborough Sounds, New Zealand. *Marine Biology*, 85, 127-136.
- Kennedy, S.K., Meloy, T.P., and Durney, T.E., 1985. Sieve data-size and shape information. *Journal of Sedimentary Petrology*, 25, 101-110.
- Kimberly, M.M., 1990. A hand-operated, 180o rotating-scoop sediment sampler. *Continental Shelf Research*, 10(12) 1157 – 1169.
- Kinlan, B.P. and Gaines, S.D. 2003. Propagule dispersal in marine and terrestrial environments: a community perspective, *Ecology*, 84(8), 2007-2020.
- Kleppel, G.S. 1993. On the diets of calanoid copepods. *Marine Ecology Progress Series*, 99, 183-195.
- Kongsberg Staff, 2006. *EM3000 Multibeam Echosounder: Product Information*. Kongsberg Maritime A.S., Norway, 855-164771 Revision E. 20/6/2006., 4p.
- Konrad, J.G., Chesters, G., and Keeny, D.R., 1970. Determination of Organic and Carbonate Carbon in Freshwater Lake Sediments by a Micro-Combustion Procedure., *Journal of Thermal Analysis*, 2, 199-208.
- Kostylev, V.E., Todd, B.J., Fader, G.B.J., Courteney, R.C., Cameron, G.D.M., and Pickrill, R.A., 2001. Benthic habitat monitoring on the Scotian Shelf based on multibeam bathymetry, surficial geology and sea floor photographs. *Marine Ecology Progress Series*, 219, 121-137.
- Kunze E., and Toole, J.M. 1997. Tidally driven vorticity, diurnal shear, and turbulence atop Fieberling Seamount. *Journal of Physical Oceanography*, 27(2), 2663–2693.

- Laing, A.K., Stanton, B.R., Challenor, P.G. 1996. Variations in the East Auckland Current from satellite radar altimeter data. *New Zealand Journal of Marine and Freshwater Research*, 30, 175-199.
- Lamb, K. G., 1994. Numerical simulations of stratified inviscid flow over a smooth obstacle. *Journal of Fluid Mechanics*, 260, 1–22.
- Large, W.G., and Pond, S. 1981. Open ocean momentum flux measurements in moderate to strong winds. *Journal of Physical Oceanography*, 5, 642-669.
- Largier, J.L. 1994. The internal tide over the shelf of Cape Point Valley, South Africa. *Journal of Geophysical Research*, 99, 10023-10034.
- Lavender, S.J., Pinkerton, M.H., Froidefond, J-M., Morales, J., Aiken, J., Moore, G.F. 2004. SeaWiFs validation in European coastal waters using optical properties and bio-geochemical measurements. *International Journal of Remote Sensing*, 25(7-8), 1481-1488.
- Ledwell J. R., Watson, A.J., and Law, C.S. 1998. Mixing of a tracer in the pycnoclin., *Journal of Geophysical Research*, 103(C10), 21499–21529.
- Lee, R.S., Ajani, P.A., Wallace, S., Pritchard, T.R., and Black, K.P., 2001. Anomalous upwelling along Australia's East Coast. *Journal of Coastal Research Special Issue 34*, (ICS 2000, New Zealand), 87-95.
- Leendertse, J. J., Liu, S-K. 1975. Modelling of three dimensional flows in estuaries, *2nd Annual Symposium on Modelling Techniques*, Waterways Harbours and Coastal Engineering, American Society of Civil Engineers, pp. 625–642.
- Lehane, C. and Davenport, J. 2006. A 15-month study of zooplankton ingestion by farmed mussels (*Mytilus edulis*) in Bantry Bay, Southwest Ireland. *Estuarine, Coastal and Shelf Science*, 67, 645-652
- Leth, O. and Middleton, J.F. 2004. A mechanism for enhanced upwelling off central Chile: Eddy advection *Journal of Physical Geophysical Research*. 109, C12020, doi:10.1029/2003JC002129.
- Levin, L.A. and Gage, J.D., 1998. Relationships between oxygen, organic matter and the diversity of bathyal macrofauna. *Deep Sea Research II*, 45, 129-163.
- Levings, C.D., Ervik, A., Johannessen, P. and Aure, J. 1995. Ecological criteria used to help site fish farms in fjords. *Estuaries*, 18(1A) 81-90.
- Li, G., Shao, Y. (1998), Remote sensing of oceanic primary productivity and its GIS estimation model. *Acta Geographica Sinica*, 53(6), 546-553.
- Linz, 1997, Motiti Island to Pehitari Point, NZ542: *Land Information New Zealand* (Linz), Hydrographic Office of the Royal New Zealand Navy, scale 1:100,000, 1 sheet;.
- Linz, 2002. Hydrographical Chart, Cuvier Island to East Cape, NZ54: *Land Information New Zealand* (Linz), Hydrographic Office of the Royal New Zealand Navy, scale 1:300,000, 1 sheet.
- Lohse, L., Helder, W., Epping, E.H.G., and Balzer, W. 1998. Recycling of organic matter along a shelf-slope transect across the N.W. European continental margin (Gopan Spur). *Progress in Oceanography*, 42, 77-110.
- Longdill, P.C., Black, K.P., Park, S. and Healy, T.R., 2005. Bay of Plenty Shelf Water Properties Data Report 2003 – 2004: Aquaculture Management Areas. Report for Environment Bay of Plenty, ASR Ltd, P.O. Box 67, Raglan, NZ, 35p.
- Longdill, P.C., Healy, T.R., Black, K.P., and Mead, S.T., 2007a. Integrated sediment habitat mapping for aquaculture zoning. *Journal of Coastal Research*, SI50 (ICS2007), 173-179.
- Longdill, P.C., Healy, T.R., Black, K.P. *in press* (2007). Transient coastal upwelling on a lee coast shelf with variable width and orientation. *New Zealand Journal of Marine and Freshwater Research*.
- Loyd, B. D. 2003. Potential effects of mussel farming on New Zealand's marine mammals and seabirds: a discussion paper. Wellington, Department of Conservation. 34p.
- Lungu, T. (ed), 2006. QuikSCAT science data product user's manual: Overview and geophysical data products, JPL D18053-Rev.A, version 3.0, Jet Propulsion Laboratory, California, USA, 85p.
- Mallet, A.L., Carver, C.E., and Landry, T., 2006. Impact of suspended and off-bottom Easter oyster culture on the benthic environment in eastern Canada. *Aquaculture*, 255, 362-373.
- Malvern, 1985. *Malvern MasterSizer Reference Manual*: Software version 2.1 and later. Malvern Instruments Ltd, Worcestershire, England.
- Manizza, M., Le Quééré, C., Watson, A.J., and Buitenhuis, E.T. 2005. Bio-optical feedbacks among phytoplankton, upper ocean physics and sea-ice in a global model. *Geophysical Research Letters*, 32, L05603, doi:10.1029/2004GL020778.
- Mann, K.H., and Lazier, J.R.N. 1996. *Dynamics of marine ecosystems: biological-physical interactions in the oceans*, Blackwell Publishing, 496p.
- Maritime New Zealand, 2005. Guidelines for aquaculture management areas and marine farms, Maritime New Zealand, December, 2005, Wellington, New Zealand, 19p.
- Maritime New Zealand, 2006. Shipping routes around the New Zealand coast, Maritime Safety Authority, Wellington, New Zealand, 14p.

- Marsden, I.D., and Weatherhead, M.A. 1999. Shore-level induced variations in condition and feeding of the mussel *Perna canaliculus* from the east coas of the South Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 33, 611-622.
- Mattson, J., and Linden, O., 1983. Benthic macrofauna succession under mussels, *Mytilus edulis* L. (Bivalvia), cultured on a hanging long-line. *Sarsia*, 68, 97-102.
- McCarthy, L.H., Thomas, R.L., and Mayfield C.I., 2004. Assessing the toxicity of chemically fractionated Hamilton Harbour (Lake Ontario) sediment using selected aquatic organisms. *Lakes & Reservoirs: Research and Management*, 9(1) 89.
- McCave, I.N., and Langhorne, D.N., 1982. Sand waves and sediment transport at the end of a tidal sand bank. *Sedimentology*, 29(1), 1365.
- McClellan-Padman, J., and Padman, L. 1991. Summer upwelling on the Sydney inner continental shelf: the relative roles of local wind forcing and mesoscale eddy encroachment. *Continental Shelf Research*, 11(4), 321-345.
- McKindsey, C.W., Thetmeyer, H., Landry, T., and Silvert, W. 2006. Review of recent carrying capacity models for bivalve culture and recommendations for research and management. *Aquaculture*, 261, 451-462.
- McKnight, D. G., 1969. Infaunal Benthic Communities of the New Zealand Continental Shelf. *New Zealand Journal of Marine and Freshwater Research*, 3, 409-444.
- Mead, S.T., Scarfe, B., Blenkinsop, C., and Frazerhurst, J., 2003. Cape Runaway Marine Survey. ASR Ltd Report May 2003, prepared for Department of conservation, Hawkes Bay Conservancy.
- Mead, S.T., Longdill, P.C., Moores, A., Beamsley, B., and Black, K.P., 2005. Bay of Plenty Biological Survey: Aquaculture Management Areas. Report for Environment Bay of Plenty, ASR Ltd, P.O. Box 67, Raglan, NZ, 35p.
- Meaden, G. 1987. Where should trout farms be in Britian? *Fish Farmer*, 10(2):33-35.
- Meaden, G. and Kapetsky, J.M. 1991. Geographic Information Systems and Remote Sensing in Inland Fisheries and Aquaculture. FAO Fisheries Technical Paper. No.318. FAO, Rome, p186-233.
- Menesguen, A., Cuiger, P., Loyer, S., Vanhoutte-Brunier, A., Hoch, T., Guillard, J.F., and Gohin, F. 2007. Two- or three- layered box models versus fine 3D models for coastal ecological modeling? A comparative study in the English Channel (Western Europe). *Journal of Marine Systems*, 64, 47-65.
- Middleton, J.F. 2006. The coastal-trapped wave paddle and open boundary conditions. *Ocean Modelling*, 12, 224-236.
- Middleton, J. F., Black, K. P. 1994. The low frequency circulation in and around Bass Strait: a numerical study. *Continental Shelf Research*, 14 (13/14), 1495–1521.
- Middleton, J.F. and Platov, G. 2003. The mean summertime circulation along Australia's Southern shelves: A numerical study. *Journal of Physical Oceanography*, 33, 2270-2287.
- Milliman, J.D, 1994. Organic matter content on the U.S. Atlantic continental slope sediments: Decoupling the grain size factor. *Deep Sea Research II*, 41, 797-808.
- Ministry for the Environment, 2003. Aquaculture reforms – resource management (aquaculture moratorium extension) bill, Ministry for the Environment, Info 46, Wellington, New Zealand, December 2003, 2p.
- Ministry for the Environment, 2005. Aquaculture reform 2004: An overview, Ministry for the Environment, Info 83, Wellington, New Zealand, January 2005, 6p.
- Ministry of Economic Development, 2007. Our blue horizon: The government's commitment to aquaculture, Ministry of Economic Development, Wellington, New Zealand, 47p.
- Mitchell, I.M., 2006. In situ biodeposition rates of Pacific Oysters (*Crassostrea gigas*) on a marine farm in Southern Tasmania (Australia). *Aquaculture*, 257, 194-203.
- Mitchum, G.T. and Clarke A.J., 1986. Evaluation of frictional, wind-forced long-wave theory on the west Florida shelf. *Journal of Physical Oceanography*, 6, 1029-1037.
- Morel, F. 1983. *Principles of aquatic chemistry*, John Wiley and Sons, New York, 150p.
- Morris, M., Roemmich, D., and Cornuelle, B. 1996. Observations of variability in the South Pacific Subtropical Gyre. *Journal of Physical Oceanography*, 26, 2359-2380.
- Morrisey, D.J., and Swales, A. 1997. *Towards the development of guidelines for sustainable management of aquaculture in the Auckland region*. Auckland Regional Council Technical Publication No.80, December 1997, 177p.
- Morrisey, D.; Turner, S. and MacDiarmid, A.B., 1998. Subtidal Assemblages of Soft Substrata. In: Kingsford, M, and Battershill, C. (eds), *Studying Temperate Marine Environments*, Canterbury University Press, Christchurch, NZ, 335p.
- Morton, J.E., and Miller, M.M. 1973. *The New Zealand sea shore* 2<sup>nd</sup> ed., Collins, Auckland, NZ, 638p.
- Mulhearn, P.J. 1987. The Tasman Front: a study using satellite infrared imagery. *Journal of Physical Oceanography*, 17, 1148-1155.

- Murphy, R.J., Pinkerton, M.H., Richardson, K.R.M., Bradford-Grieve, J.M., and Boyd, P.W. 2001. Phytoplankton distributions around New Zealand derived from SeaWiFS remotely-sensed ocean colour data. *New Zealand Journal of Marine and Freshwater Research*, 35, 343-362.
- Mussel Industry Council, 2005. The greenshell story, Mussel Industry Council of New Zealand, 15p.
- Nakanashi, H., Ukita, M., and Kawai, Y. 1986. Study on the modelling of the behaviour of phosphorus released from sediments. *Ecological Modelling*, 31(1-4), 105-123.
- Nath, S.S., Bolte, J.P., Ross, L.G., Aguilar-Manjarrez, J. 2000. Applications of geographic information systems (GIS) for spatial decision support in aquaculture. *Aquacultural Engineering*, 23, 233-278.
- Navarro, E., Iglesias, J.I.P., Perez-Camacho, A., Labarta, U., and Berias, R. 1991. The physiological energetics of mussels (*Mytilus galloprovincialis* Lmk) from different cultivation rafts in the Ria de Arosa (Galicia, N.W. Spain). *Aquaculture*, 94, 197-212.
- Naylor, R.L., Goldberg, R.J., Primavera, J.H., Kautsky, N., Beveridge, M.C.M., Clay, J., Folke, C., Lubchenco, J., Mooney, H. and Troell, M. 2000. Effect of aquaculture on world fish supplies. *Nature*, 405(29), 1017-1024.
- New Zealand Government, 2004. Aquaculture reform bill 2004: Explanatory note, Bill 181-1, New Zealand Government, December 2004. 162p.
- NWASCO, 1982. Physical and chemical methods for water quality analysis, *Water and Soil Miscellaneous Publication No. 38*. (National Water and Soil conservation Authority: Wellington, New Zealand).
- O'Neil, R.V., DeAngelis, D.L., Pastor, J.J., Jackson, B.J., and Post, W.M. 1989. Multiple nutrient limitations in ecological models. *Ecological Modelling*, 46, 147-163.
- Odum, E.P. 1983. *Basic ecology*, WB Saunders, Philadelphia, 320p.
- Officer, C.B., Smayda, T.J., and Mann, R. 1982. Benthic filter feeding: a natural eutrophication control. *Marine Ecology Progress Series*, 9, 203-210.
- Ogilvie, S.C., Ross, A.H., and Schiel, D.R., 2000. Phytoplankton biomass associated with mussel farms in Beatrix Bay, New Zealand. *Aquaculture*, 181, 71-80
- Ogilvie, S.C., Ross, A.H., James, M.R., and Schiel, D.R. 2003. In situ enclosure experiments on the influence of cultured mussels (*Perna canaliculus*) on phytoplankton at times of high and low ambient nitrogen. *Journal of Experimental Marine Biology and Ecology*, 25, 23-39.
- Oke, P.R., Middleton, J.H. 2000. Topographically induced upwelling off eastern Australia. *Journal of Physical Oceanography*, 30, 512-531.
- Oliver, J.K. 1995. Is the 'limits of acceptable change' concept useful for environmental managers? A case study from the Great Barrier Reef Marine Park. In: Grigg, G.C., Hale, P.T., and Lunney, D. (eds), *Conservation through sustainable use of wildlife*, pp. 11-19.
- Operational Oceanography Group: Global Temperature-Salinity Profile Program. June 2006. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Oceanographic Data Center, Silver Spring, Maryland, 20910.
- Paffenhöfer, G.A., Orcutt, J.D. 1986. Feeding, growth and food conversion of the marine cladoceran *Penilia avirostris*. *Journal of Plankton Research*, 8(4), 741-754.
- Paffenhofer, G.A., Atkinson, L.P., Blanton, J.O., Lee, T.N., Pomeroy, L.R., and Yoder, J.A. 1987. Summer upwelling on the southeastern continental shelf of the U.S.A. during 1981. *Progress in Oceanography*, 19, 221-230.
- Park, S. 1991. Bay of Plenty Regional Council Coastal Overview Report – 1991. Bay of Plenty Regional Council Technical Publication Number 3, August 1991.
- Park, S. 1998. Bay of Plenty Coastal Water Quality: 1996-1997. A report prepared of Environment Bay of Plenty, no. 98/5.
- Parker, M.R., Beale, B.F., Congelton, W.R.Jr, Pearce, B.R., and Morin, L. 1998. Utilization of GIS and GPS for shellfish growout site selection. *Journal of Shellfish Research*, 17, 1491-1495.
- Paw, J.N., Diamante, D.A.D., Robles, N.A., Chua, T.E., Quitos, L.N., and Cargamento, A.G.A. 1992. Site selection for brackish water aquaculture development and mangrove reforestation in Lingayen Gulf, Phillipines using geographic information systems. Proceedings of Canadian Conference on GIS, Ottawa, Canada, March 1992. p821-823.
- Pawlowicz, R., Beardsley, B., and Lentz, S. 2002. Classical tidal harmonic analysis including error estimates in MATLAB using T\_TIDE. *Computers and Geosciences*, 28, 929-937.
- Pearson, T.H., and Rosenberg, R., 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. In: Barnes, H. (ed.), *Oceanography and Marine biology: An Annual Review*, 16, 229-311.
- Perez, O.M., Telfer, T.C., and Ross, L.G. 2003. Use of GIS-based models for integrating and developing marine fish cages within the tourism industry in Tenerife (Canary Islands). *Coastal Management*, 31, 355-366.
- Perez, O.M., Telfer, T.C., and Ross, L.G., 2005. Geographical information systems-based models for offshore floating marine fish cage aquaculture site selection in Tenerife, Canary Islands. *Aquaculture Research*, 36, 946-961.

- Pérez-Camacho, A., Labarta, U., and Beiras, R. 1995. Growth of mussel (*Mytilus edulis galloprovincialis*) in cultivation raft: Influence of seed source, cultivation site and food availability. *Aquaculture*, 138, 349–362.
- Perrels, P.A.J., and Karelse, M. 1982. A two-dimensional laterally averaged model for salt intrusion in estuaries. Delft Hydraulics Laboratory Report No. 262.
- Peters, R.H., Downing, J.A. 1984. Empirical analysis of zooplankton filtering and feeding rates. *Limnology & Oceanography* 29(4), 763-784.
- Pickard, G.L., and Emery, W.J. 1990. *Descriptive physical oceanography: An Introduction* (5<sup>th</sup> ed), Pergamon Press, Oxford, England, 320p.
- Pickrill, R.A., and Mitchell, J.S. 1979. Ocean wave characteristics around New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 13, 319-324.
- Pillay, T.V.R., 2004. *Aquaculture and the Environment* (2nd ed). Blackwell Scientific Publishing, Oxford, U.K. 195p.
- Pinkerton, M.H., Gall, M.P., and Hill, P. 2005. Bio-Optics Research Voyage EBoP0604 supporting Ocean Colour Remote Sensing of the Bay of Plenty. National Institute of Water and Atmospheric Research Ltd. Client Report WLG2002-00, NIWA project EBOP04302, 27p.
- Pinkerton, M.H., Moore, G.F., Lavender, S.J., Gall, M.P., Oubelkheir, K., Richardson, K.M., Boyd, P.W., Aiken, J. 2006. A method for estimating inherent optical properties of New Zealand continental shelf waters from satellite ocean colour measurements. *New Zealand Journal of Marine and Freshwater Research*, 40, 227-247.
- Pitcher, G.C. and Calder, D. 1998. Shellfish mariculture in the Benguela system: phytoplankton and the availability of food for commercial mussels farms in Saldanha Bay, South Africa. *Journal of Shellfish Research*, 17, 15-25.
- Pitts, J. 1990. Washington State's environmental impact statement (EIS) on fish culture in floating net-pens. *In: Proceedings of Aquaculture International*, 4-7 September 1990, Vancouver, Canada: 285-289.
- Plackett, R.L. and Burman, J.P. 1946. The design of optimum multifactorial experiments. *Biometrika*, 33, 305-325.
- Plew, D.R. 2005. The hydrodynamic effects of long-line mussel farms, PhD thesis, University of Canterbury, Christchurch, New Zealand, 330p.
- Plew, D.W., Stevens, C.L., Spigel, R.H., Hartstein, N.H. 2005. Hydrodynamic implications of large offshore mussel farms. *IEEE Journal of Oceanic Engineering*, 30(1), 95-108.
- Pond, S., and Pickard, G.L. 1983. *Introductory Dynamical Oceanography* (2<sup>nd</sup> ed.), Pergamon Press, Oxford, U.K., 329p.
- Poppe, L.J., Eliason, A.H., Fredericks, J.J., Rendigs, R.R., Blackwood D. and Polloni, C.F., 2000, Grain-size Analysis of Marine Sediments: Methodology and Data Processing. U.S. Geological Survey Open-file Report 00-358. U.S. Geological Survey, Woods Hole, MA 02543.
- Preston, N., Macleaoir, I., Rothlisberg, P., and Lang, B. 1997. Environmentally sustainable aquaculture production – an Australian perspective. *In: Hancock, D.A., Smith, D.C., Grant, A., and Beumer, J.P. (eds), Developing and sustaining world fisheries resources: The state of science and management*, 2<sup>nd</sup> World fisheries congress, CSIRO, Australia, 1997, 471-477.
- Price, J.M., Johnson, W.R., Ji, Z.G., Marshall, C.F., and Rainey, G.B. 2004. Sensitivity testing for improved efficiency of a statistical oil-spill risk analysis model. *Environmental Modelling and Software*, 19, 671-679.
- Prins, T.C., Smaal, A.C., and Dame, R.F. 1998. A review of the feedbacks between bivalve grazing and ecosystem processes. *Aquatic Ecology*, 31, 349-359.
- Prior, D.B., and Bornhold, B.D., 1989. Submarine sedimentation on a developing Holocene fan delta. *Sedimentology*, 36(6) 1053.
- Procter, R., and Greig, M.J.N. 1989. A numerical model investigation of the residual circulation in Hauraki Gulf, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 23, 421-442.
- Raillard, O., and Menesguen, A. 1994. An ecosystem box model for estimating the carrying capacity of a macrotidal shellfish system. *Marine Ecology Progress Series*, 115, 117-130.
- Rajitha, K., Mukherjee, C.K., Vinu-Chandran, R. 2007. Applications of remote sensing and GIS for sustainable management of shrimp culture in India. *Aquacultural Engineering*, 36, 1-17.
- Rawle, A., 1995. *The basic principles of particle size analysis*. Technical Paper, Malvern Instruments Ltd, England, 8p.
- Redfield A.C. 1934. On the proportions of organic derivations in sea water and their relation to the composition of plankton. *In: Daniel, R.J. (ed) James Johnson Memorial Volume*. University Press of Liverpool, pp. 177-192.
- Redfield, A.C. 1958. The biological control of chemical factors in the environment. *American Scientist*, 46, 205-221.

- Ren, J.S. and Ross, A.H., 2005. Environmental influence on mussel growth: A dynamic energy budget model and its application to the greenshell mussel *Perna canaliculus*. *Ecological Modelling*, 189, 347-362.
- Rew, R., Davis, G., Emmerson, S., Davies, H., and Hartnet, E. 2007. The netCDF users guide: Data model, programming interfaces, and format for self-describing portable data. Unidata Program Center, 15 February 2007, 60p.
- Reynaud, T., Grant-Ingram, R., Freeland, H.J., Weaver, A.J. 1992. Propagation of coastal-trapped waves under an ice cover in Hudson Bay. *Atmosphere-Ocean*, 30(4), 593-620.
- Richardson, K.M., Pinkerton, M.H., Uddstrom, M.J., Gall, M.P., and Hill, P. 2005. Remote Sensing Survey of the Bay of Plenty: Report on Sea Surface Temperature and Ocean Colour Product Generation for Environment Bay of Plenty, National Institute of Water and Atmospheric Research Ltd, NIWA Report EBOP04302, 27p.
- Ridgeway, N.M., and Greig, M.J.N. 1986. Water movements in Bay of Plenty, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 20, 447-453.
- Robinson, C.L.K., Morrison, J. and Foreman, M.G.G. 2005. Oceanographic connectivities among marine protected areas on the north coast of British Columbia, Canada. *Canadian Journal of Fisheries and Aquatic Sciences*, 62(6), 1350-1362.
- Rochford, D.J., 1975. Nutrient enrichment of east Australian coastal waters II, Laurieton upwelling. *Australian Journal of Marine and Freshwater Research*, 26, 233-243.
- Rodhouse, P.G., Roden, C.M. 1987. Carbon budget for a coastal inlet in relation to intensive cultivation of suspension-feeding bivalve molluscs. *Marine Ecology Progress Series*, 36, 225-236.
- Rodhouse, P.G., Roden, C.M., Hensey, M.P., and Ryan, T.H. 1985. Production of mussels, *Mytilus edulis*, in suspended culture and estimates of carbon and nitrogen flow: Killaroy Harbour, Ireland. *Journal of the Marine Biology Association U.K.* 65, 55-68.
- Roemmich, D., and Cornuelle, B. 1990. Observing the fluctuations of gyre scale circulation: a study of the subtropical South Pacific. *Journal of Physical Oceanography*, 20, 1919-1934.
- Roemmich, D., and Sutton, P. 1998. The mean and variability of ocean circulation past northern New Zealand: determining the representativeness of hydrographic climatologies. *Journal of Geophysical Research*, 103, 13041-13054.
- Ross, A.H., James, M.R., Hadfield, M., and Gibbs, M.M. 1999. Estimating the sustainable production of mussel aquaculture using numerical simulation models. International Conference on Sustainable Management of Coastal Ecosystems, Fernando Pessoa University, Porto, Portugal, November 1999.
- Ross, L.G., Mendoza Q.M., E.A., and Beveridge, M.C.M. 1993. The application of geographical information systems to site selection for coastal aquaculture: an example based on salmonid cage culture. *Aquaculture*, 112, 165-178.
- Roughan, M., and Middleton, J.H. 2002. A comparison of observed upwelling mechanisms off the east coast of Australia. *Continental Shelf Research*, 22, 2551-2572.
- Rowe, G.T., Hovey-Clifford, C., Smith, K.L. Jr., Lawrence-Hamilton, P. 1975. Benthic nutrient regeneration and its coupling to primary productivity in coastal waters. *Nature*, 255, 215-217.
- Rubec, P.J., Christensen, J.D., Arnold, W.S., Norris, H., Steele, P., and Monaco, M.E. 1998. GIS and modeling: coupling habitats to Florida fisheries. *Journal of Shellfish Research*, 17(5), 1451-1457
- Sahu, B.K., 1965. Theory of sieving. *Journal of Sedimentary Petrology*, 35, 750-753.
- Saunders, P.M., Coward, A.C., and de Cuevas, B.A., 1999. Circulation of the Pacific Ocean seen in a global ocean model: Ocean circulation and climate advanced modelling project (OCCAM). *Journal of Geophysical Research*, 104(C8), 18,281-18,300.
- Shackley, S.E., and Collins, M., 1984. Variations in sublittoral sediments and their associated macro-infauna in response to inner shelf processes; Swansea Bay, U.K. *Sedimentology*, 31(6), 793.
- Sharples, J. 1997. Cross-shelf intrusion of subtropical water into the coastal zone of northeast New Zealand. *Continental Shelf Research*, 17(7), 835-857.
- Sharples, J., and Greig, M.J.N. 1998. Tidal currents, mean flows, and upwelling on the north-east shelf of New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 32, 215-231.
- Shears, N.T., and Babcock, R.C., 2003. Quantitative classification of New Zealand rocky coastal community types. Leigh Marine Laboratory, University of Auckland, 53p.
- Shevyrnogov, A., Vysotskaya, G., and Shevyrnogov, E. 2004. Spatial and temporal anomalies of sea surface temperature in global scale (by space-based data). *Advances in Space Research*, 33, 1179-1183.
- Siddal, S.E., 1980. A clarification of the genus *Perna* (*Mytilidae*). *Bulletin of Marine Science*, 30, 858-870.
- Silker, W.B. 1972. Horizontal and vertical distributions of radionuclides in the north Pacific Ocean. *Journal of Geophysical Research*, 77, 1061-1070.
- Silvert, W. and Cromey, C.J. 2001. Modelling impacts. In: Black, K.D. (ed), *Environmental impacts of aquaculture*. Sheffield Academic Press, Sheffield, U.K., 154-181.

- Simms, A. 2002. GIS and aquaculture: Assessment of soft-shell clam sites. *Journal of Coastal Conservation*, 8, 35-47.
- Smaal, A.C., and Prins, T.C. 1993. The uptake of organic matter and the release of inorganic nutrients by bivalve suspension feeder beds. In: Dame, R.F. (ed). *Bivalve filter feeders in estuarine and coastal ecosystems processes*. NATO ASI series vol. G33, Springer Verlag, Berlin, 271-298.
- Smaal, A.C. and Heral, M. 1998. Preface: modelling bivalve carrying capacity. *Aquatic Ecology*, 31, 347.
- Smaal, A.C., Prins, T.C., Dankers, N., and Ball, B. 1998. Minimum requirements for modelling bivalve carrying capacity. *Aquatic Ecology*, 31, 423-428.
- Smaal, A.C., van Stralen, M., and Schuiling, E. 2001. The interaction between shellfish culture and ecosystem processes. *Canadian Journal of Fisheries and Aquatic Science*, 58, 991-1002.
- Smith, K.A. 2000. Active and passive dispersal of *Centroberyx affinis* (Berycidae) and *Gonorynchus greyi* (Gonorynchidae) larvae on the Sydney shelf. *Marine and Freshwater Research*, 51, 229-234.
- Smith, R.L. 1981. A comparison of the structure and variability of the flow field in three coastal upwelling regions: Oregon, northwest Africa, and Peru. In: Richards, F.A. (ed), *Coastal Upwelling*, American Geophysical Union, Washington, pp.107-118.
- Smyth, T.J., Moore, G.F., Hirata, T., Aiken, J. 2006. Semianalytical model for the derivation of ocean color inherent optical properties: description, implementation and performance assessment. *Applied Optics*, 45(31), 8116-8131.
- Soniat, T.M., and Ray, S.M. 1985. Relationship between possible available food and the composition, condition and reproductive state of oysters from Galveston Bay, Texas. *Contributions in Marine Science*, 28, 109-121.
- Soniat, T.M., and Brody, M.S. 1988. Field validation of a habitat suitability index model for the American oyster. *Estuaries*, 11(2), 87-95.
- Spencer, B.E., 2002. *Molluscan shellfish farming*, Blackwell science, Oxford, U.K. 296p.
- Stanton, B.R. 1978. Preliminary observations of the internal tide over the shelf west of New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 11, 703-711.
- Stanton, B.R. 1981. An oceanographic survey of the Tasman Front. *New Zealand Journal of Marine and Freshwater Research*, 15, 289-298.
- Stanton, B.R. 1995. Sea level variability on the west coast of New Zealand. *Journal of Physical Oceanography*, 25, 1265-1272.
- Stanton, B.R. 2001. Estimating the East Auckland Current transport from model winds and the island rule. *New Zealand Journal of Marine and Freshwater Research*, 35, 531-540.
- Stanton, B.R., Sutton, P.J., Chiswell, S.M. 1997. The East Auckland Current, 1994-95. *New Zealand Journal of Marine and Freshwater Research*, 31, 537-549.
- Stanton, B., Goring, D.G., and Bell, R.G. 2001. Observed and modeled tidal currents in the New Zealand region. *New Zealand Journal of Marine and Freshwater Research*, 35, 397-415.
- Stanton, B.R., and Sutton, P.J. 2003. Velocity measurements in the East Auckland Current north-east of North Cape, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 37, 195-204.
- Stead, D.H., 1971. A preliminary survey of mussel stocks in Pelorus Sound, New Zealand Fisheries Technical Report 61. 16p.
- Steele, J.H., 1965. Notes on some theoretical problems in production ecology. In: C.R. Goldman (ed.), *Primary production in aquatic environments*. University of California Press, Berkeley, California, pp393-398.
- Steiger, J.H. 1980. Tests for comparing elements in a correlation matrix. *Psychological Bulletin*, 87, 245-251.
- Stenton-Dozey, J.M.E., Jackson, L.F., and Busby, A.J. 1999. Impact of mussel culture on macrobenthic community structure in Saldanha Bay, South Africa. *Marine Pollution Bulletin*, 39, 357-366.
- Stephens, S.A., Bell, R.G. and Black, K.P. 2001. Complex circulation in a coastal embayment: shelf-current wind and density-driven circulation in Poverty Bay, New Zealand. *Journal of Coastal Research*, SI 34 (ICS2000), 45-59.
- Strohmeier T., Aure J., Duinker A., Castberg T., Svoldal A. and Strand Ø. 2005. Flow reduction, seston depletion, meat content and distribution of diarrhetic shellfish toxins in a long-line blue mussel (*Mytilus edulis*) farm. *Journal of Shellfish Research*, 24, 15-23.
- Sutton, P.J.H., and Roemmich, D., 2001. Ocean temperature climate of north-east New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 35, 553-565.
- Sutton, P.J.H., and Chereskin, T.K. 2002. Absolute geostrophic currents in the East Auckland Current region. *New Zealand Journal of Marine and Freshwater Research*, 36, 751-762.
- Takahashi, W. and Kawamura, H. 2005. Detection method of the Kuroshio front using the satellite-derived chlorophyll-a images. *Remote Sensing of Environment*, 97, 83-91.
- Taylor, J.R., and Park, S.G. 2001. Natural environment regional monitoring network: Bay of Plenty rivers' water quality (1989-2000). Environment BOP environmental report 2001/12, 118p.

- Telfor, T., Robinson, K. 2003. Environmental quality and carrying capacity for aquaculture in Mulroy Bay Co. Donegal. Marine Environment and Health Series No. 9
- Tenore, K.R., Boyer, L.F., Cal, R.M., Corral, J., Garcia-Fernandez, C., Gonzalez, N., Gonzalez-Gurriaran, E., Hanson, R.B., Iglesias, J., Krom, M., Lopez-Jamar, E., McClain, J., Pamatmat, M.M., Perez, A., Rhoads, D.C., de Santiago, G., Tietjen, J., Westrich, J., Windom, H.L., 1982. Coastal upwelling in the Rias Bajas, NW Spain: contrasting the benthic regimes of the Rias de Arosa and de Muros. *Journal of Marine Research*, 40, 701–772.
- Tetra Tech, 1980. Methodology for evaluation of multiple power plant cooling system effect, Volume V. Methodology Application to Prototype – Cayuga Lake. Tetra Tech, Inc., Lafayette, California. For Electric Power Research institute. Report EPRI EA-III.
- Thomann, R.V., and Kirkpatrick, J.J., 1982. Calibration and validation of a mathematical model of the eutrophication of the Potomac estuary, Prepared for the Department of Environmental Services, Government of the District of Colombia, HydroQual Inc, Mahwah, NJ.
- Thomson, N.W., 1996. Trends in Australasian open water aquaculture, *In: Polk, M (ed), Open ocean aquaculture, Proceedings of an international conference*, May 8-10, 1996, Portland, Maine, 223-234.
- Tilburg, C.E., Hurlburt, H.E., O'Brien, J.J., Shriver, J.F. 2001. The dynamics of the East Australian Current system: the Tasman Front, the East Auckland Current, and the East Cape Current. *Journal of Physical Oceanography*, 31, 2917-2943.
- Tilburg, C.E., and Garvine, R.W. 2003. Three-dimensional flow in a shallow coastal upwelling zone: Alongshore convergence and divergence on the New Jersey shelf. *Journal of Physical Oceanography*, 33, 2113-2125.
- Tomczak, M., and Godfrey, J.S. 1994. *Regional Oceanography: An Introduction*, Pergamon Press, Oxford, UK. 422p.
- Treml, E.A., Halpin, P.N., Urban, D.L., and Pratson, L.F. (in press). Modeling Population Connectivity by Ocean Currents, a Graph-theoretic Approach for Marine Conservation. *Landscape Ecology*.
- Turner, M., Walker, J. P., Oke, P. and Grayson R. B., 2005. Assimilation of sea-surface temperature into a hydrodynamic model of Port Phillip Bay, Australia, *Coasts and Ports Australasian Conference*, Adelaide Australia, 21-23 September.
- Uddstrom, M.J., Oien, N.A. 1999. On the use of high-resolution satellite data to describe the spatial and temporal variability of sea surface temperatures in the New Zealand region. *Journal of Geophysical Research – Oceans*, 104, 20,729-20,751.
- Urbanski, J.A., and Szymelfenig, M. 2003. GIS-based mapping of benthic habitats. *Estuarine, Coastal and Shelf Science*, 56, 99-109.
- Valavanis, V. 2002. *Geographic Information Systems in Oceanography and Fisheries*. Taylor and Francis, London, 209p.
- Valavanis, V.D., Drakopoulos, P., and Georgakarakos, S. 1999. A study of upwellings using GIS. In *Proceedings of Coast GIS 1999, International Conference of GIS and New Advances in Coastal Management*, September 1999, Brest, France, IFREMER.
- Valavanis, V.D., Kapantagakis, A., Katara, I., and Palialexis, A. 2004. Critical regions: A GIS-based model of marine productivity Hotspots. *Aquatic Sciences*, 66, p139-148.
- Valiky, J.M., 1989. The biology and culture of mussels of the genus *Perna*. ICLARM Studies and Reviews 17, International Center for Living Aquatic Resource Management, Manila, Philippines, 63p.
- Vennel, R., and Moore, M. 1993. Acoustic Doppler Current Profiler measurements of the semi-diurnal internal tide on the west coast of New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 27, 31-38.
- Verity, P.G. 1985. Grazing, respiration, excretion and growth rates of tintinnids. *Limnology and Oceanography*, 30(6), 1268-1282.
- Vincenzi, S., Caramori, G., Rossi, R., and De Leo, G.A. 2006. A GIS-based habitat suitability model for commercial yield estimation of *Tapes philippinarum* in a Mediterranean coastal lagoon (Sacca di Goro, Italy). *Ecological Modelling*, 193, 90-104.
- Waite, R.P. 1989. The nutritional biology of *Perna canaliculus* with special reference to intensive mariculture systems. Unpublished PhD thesis, University of Canterbury, New Zealand, 136p.
- Walters, R.A., Goring, D.G., and Bell, R.G. 2001. Ocean tides around New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 35, 567-579.
- Webb, D.J., de Cuevas, B.A., and Coward, A.C., 1998. The first main run of the OCCAM global ocean model, internal report of James Rennell Div., Southampton Oceanography Centre, 50 pp., Southampton, England, U.K.
- Weisburg, S.B., Ranasinghe, J.A., Dauer, D.A., Schaffner, L.C., Diaz, R.J., and Frithsen, J.B. 1997. An estuarine benthic index of biotic integrity (B-IBI) for the Chesapeake Bay. *Estuaries*, 20, 149–158.
- Widdows, J., Fieth, P., and Worrall, C.M. 1979. Relationships between seston, available food and feeding activity in the common mussel *Mytilus edulis*. *Marine Biology*, 50, 195-207.



- WILDCO (Wildlife Supply Company Ltd), 2006. Wildlife Supply Company Catalogue. *Wildlife Supply Company*, 95 Botsford Place, Buffalo, NY, USA.
- Winter, J.E. 1978. A review on the knowledge of suspension feeding in lamellibranchiate bivalves, with special reference to artificial aquaculture systems. *Aquaculture*, 13, 1-33.
- Wolff, M. 1994. A trophic model for Tongoy Bay – a system exposed to suspended scallop culture (Northern Chile). *Journal of Experimental Marine Biology and Ecology*, 182, 149-168.
- Wu, J. 1982. Wind stress coefficients over sea surface from sea breeze to hurricanes. *Journal of Geophysical Research*, 87, 9704-9706.
- Yankovsky, A.E. 2003. The cold-water pathway during an upwelling event on the New Jersey shelf. *Journal of Physical Oceanography*, 33, 1954-1966.
- Young, I.R., Black, K.P., Heron, M.L. 1994. Circulation in the ribbon reef region of the Great Barrier Reef. *Continental Shelf Research*, 14 (2/3), 117-142.
- Young, J.A., and Silker, W.B. 1974. The determination of air-sea exchanges and oceanic mixing rates using <sup>7</sup>Be during the bomex experiment. *Journal of Geophysical Research*, 79, 4481-4489.
- Zeldis, J.R., Walters, R.A., Greig, M.J.N., and Image, K. 2004a. Circulation over the northeastern New Zealand continental slope, shelf and adjacent Hauraki Gulf, during spring and summer. *Continental shelf Research*, 24, 543-561.
- Zeldis, J., Robinson, K., Ross, H., and Hayden, B. 2004b. First observations of predation by New Zealand greenshell mussels (*Perna canaliculus*) on zooplankton. *Journal of Experimental Marine Biology and Ecology*, 311, 287-299.
- Zeldis, J., Felsing, M., and Wilson, J. 2006. Limits of acceptable change: a framework for marine farming. *Water and Atmosphere*, 14(2), 16-17.
- Zeng, T.Q., Dorman, F., Ogburn, D., Derwent, L., and Williams, R. 2003. Aquaculture management with Geographical Information Systems (GIS) in NSW fisheries, Australia, in: Woodroffe, C.D. and Furness, R.A (eds), *Coastal GIS 2003: an integrated approach to Australian coastal issues*, Wollongong papers in maritime policy, 14, 454-466.

