

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Editors: **W. Clegg** and **D. G. Watson**

(6-Acetoxy-2-acetylphenyl- $\kappa^2 C^1, O^1$)tetracarbonylmanganese(I)

Victor D. Fester, Lyndsay Main and Brian K. Nicholson

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site or institutional repository provided that this cover page is retained. Republication of this article or its storage in electronic databases other than as specified above is not permitted without prior permission in writing from the IUCr.

For further information see <http://journals.iucr.org/services/authorrights.html>

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(6-Acetoxy-2-acetylphenyl- κ^2C^1,O^1)-tetracarbonylmanganese(I)

Victor D. Fester, Lyndsay Main and Brian K. Nicholson*

Chemistry Department, University of Waikato, Private Bag 3105, Hamilton 3240, New Zealand

Correspondence e-mail: b.nicholson@waikato.ac.nz


Received 18 September 2007; accepted 27 September 2007

Key indicators: single-crystal X-ray study; $T = 168$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.026; wR factor = 0.073; data-to-parameter ratio = 12.7.

The title compound, $[Mn(C_{10}H_9O_3)(CO)_4]$, is formed by orthomanganation of 3'-acetoxyacetophenone at the sterically crowded *ortho* site. The atoms of the benzene and the cyclometallated rings are coplanar to within <0.018 Å, and there are no significant intramolecular interactions between the $Mn(CO)_4$ group and the adjacent acetoxy group.

Related literature

The preparation and structures of related orthomanganated aryl ketones have been reviewed (Main & Nicholson, 1994). Preference in cyclometallation reactions for the more sterically crowded isomer for other 3'-substituted acetophenones has been observed (Cooney *et al.*, 1988, 2001; Liebeskind *et al.*, 1989).



Experimental

Crystal data

$[Mn(C_{10}H_9O_3)(CO)_4]$
 $M_r = 344.15$
 Triclinic, $P\bar{1}$
 $a = 9.306$ (2) Å
 $b = 9.635$ (2) Å
 $c = 9.954$ (2) Å
 $\alpha = 68.594$ (3)°
 $\beta = 67.663$ (3)°

$\gamma = 68.422$ (3)°
 $V = 740.9$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.92$ mm⁻¹
 $T = 168$ (2) K
 $0.72 \times 0.65 \times 0.15$ mm

Data collection

Bruker SMART CCD diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1998)
 $T_{min} = 0.711$, $T_{max} = 0.870$

9684 measured reflections
 2988 independent reflections
 2730 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.026$
 $wR(F^2) = 0.073$
 $S = 1.02$
 2988 reflections

235 parameters
 All H-atom parameters refined
 $\Delta\rho_{max} = 0.26$ e Å⁻³
 $\Delta\rho_{min} = -0.41$ e Å⁻³

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999).

The authors thank Dr Jan Wikaira, University of Canterbury, for collection of X-ray intensity data.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2041).

References

- Bruker (2001). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Cooney, J. M., Gommans, L. H. P., Main, L. & Nicholson, B. K. (1988). *J. Organomet. Chem.* **349**, 197–208.
 Cooney, J. M., Gommans, L. H. P., Main, L. & Nicholson, B. K. (2001). *J. Organomet. Chem.* **634**, 157–166.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Liebeskind, L. S., Gasdaska, J. R., McCallum, J. S. & Tremont, S. J. (1989). *J. Org. Chem.* **54**, 669–677.
 Main, L. & Nicholson, B. K. (1994). *Adv. Met.-Org. Chem.* **3**, 1–50.
 Sheldrick, G. M. (1997). SHELXS97 and SHELXL97. University of Göttingen, Germany.
 Sheldrick, G. M. (1998). SADABS. University of Göttingen, Germany.

supplementary materials

Acta Cryst. (2007). E63, m2655 [doi:10.1107/S1600536807047605]

(6-Acetoxy-2-acetylphenyl- κ^2C^1, O^1)tetracarbonylmanganese(I)

V. D. Fester, L. Main and B. K. Nicholson

Comment

(2-Acetyl- κO^1 -6-acetoxyphenyl- κC^1)tetracarbonylmanganese(I) is formed by orthomanganation of 3'-acetoxyacetophenone at the sterically crowded *ortho* site. The atoms of the phenyl and the cyclometallated rings are coplanar to within <0.018 Å, and there are no significant intramolecular interactions between the Mn(CO)₄ group and the adjacent acetoxy one. The Mn1—C12 distance *trans* to O1 (1.8046 (18) Å) is shorter than the Mn1—C11 distance *trans* to C1 (1.8367 (19) Å), while the two Mn—CO distances *trans* to each other are the longest (av. 1.857 (2) Å). The Mn1—C1 distance is significantly shorter than the Mn1—O1 distance (2.0445 (15) and 2.0548 (11) Å respectively) contrary to the trend expected purely on covalent radii. These conform to the pattern found for other orthomanganated arenes (Main and Nicholson, 1994).

Experimental

The title compound was prepared by reaction of PhCH₂Mn(CO)₅ with 3'-acetoxyacetophenone in refluxing heptane, under the usual conditions (Main and Nicholson, 1994). The structure was determined to confirm that cyclometallation had occurred at the sterically crowded 2'-position, rather than the alternative 6'-position, and to detect any residual interactions with the adjacent acetoxy group which might have directed this preference.

Refinement

All H-atoms were located as the highest peaks in a penultimate difference map and were refined with isotropic temperature factors.

Figures




Fig. 1. Structure of (2-Acetyl- κO^1 -6-acetoxyphenyl- κC^1)tetracarbonylmanganese(I), with ellipsoids drawn at the 30% probability level.

(6-Acetoxy-2-acetylphenyl- κ^2C^1, O^1)tetracarbonylmanganese(I)

Crystal data

[Mn(C₁₀H₉O₃)(CO)₄]

Z = 2

M_r = 344.15

*F*₀₀₀ = 348

Triclinic, *P* $\bar{1}$

D_x = 1.543 Mg m⁻³

supplementary materials

Hall symbol: -P 1

$a = 9.306$ (2) Å

$b = 9.635$ (2) Å

$c = 9.954$ (2) Å

$\alpha = 68.594$ (3)°

$\beta = 67.663$ (3)°

$\gamma = 68.422$ (3)°

$V = 740.9$ (3) Å³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 3920 reflections

$\theta = 2-26^\circ$

$\mu = 0.92$ mm⁻¹

$T = 168$ (2) K

Block, yellow

$0.72 \times 0.65 \times 0.15$ mm

Data collection

Bruker SMART CCD
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 168$ (2) K

ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1998)

$T_{\min} = 0.711$, $T_{\max} = 0.870$

9684 measured reflections

2988 independent reflections

2730 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.015$

$\theta_{\text{max}} = 26.4^\circ$

$\theta_{\text{min}} = 2.3^\circ$

$h = -11 \rightarrow 11$

$k = -12 \rightarrow 11$

$l = -12 \rightarrow 12$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.026$

$wR(F^2) = 0.073$

$S = 1.02$

2988 reflections

235 parameters

Primary atom site location: structure-invariant direct
methods

Secondary atom site location: difference Fourier map

Hydrogen site location: difference Fourier map

All H-atom parameters refined

$$w = 1/[\sigma^2(F_o^2) + (0.0452P)^2 + 0.1914P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.004$

$\Delta\rho_{\text{max}} = 0.26$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.41$ e Å⁻³

Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.15624 (3)	0.30950 (3)	0.24687 (2)	0.03251 (9)
C1	0.37221 (17)	0.14695 (17)	0.24738 (15)	0.0287 (3)
C2	0.46885 (18)	0.18432 (17)	0.30330 (16)	0.0311 (3)
C3	0.6217 (2)	0.09150 (19)	0.31439 (19)	0.0380 (3)
C4	0.6832 (2)	-0.0424 (2)	0.2682 (2)	0.0428 (4)
C5	0.5919 (2)	-0.08368 (19)	0.21356 (19)	0.0389 (3)
C6	0.44005 (18)	0.00979 (17)	0.20521 (16)	0.0314 (3)
C7	0.3934 (2)	0.32656 (18)	0.35101 (16)	0.0337 (3)
C8	0.4716 (3)	0.3870 (2)	0.4155 (2)	0.0425 (4)

C9	0.29739 (19)	-0.15989 (18)	0.22594 (19)	0.0378 (3)
C10	0.2255 (3)	-0.2011 (2)	0.1395 (3)	0.0512 (5)
C11	-0.0262 (2)	0.4716 (2)	0.2641 (2)	0.0509 (4)
C12	0.0719 (2)	0.2290 (2)	0.1657 (2)	0.0415 (4)
C13	0.08616 (19)	0.1809 (2)	0.43689 (19)	0.0395 (4)
C14	0.2643 (2)	0.4043 (2)	0.0558 (2)	0.0470 (4)
O1	0.25602 (14)	0.39925 (12)	0.33756 (13)	0.0383 (3)
O2	0.35525 (14)	-0.03417 (12)	0.14249 (12)	0.0359 (2)
O3	0.30441 (17)	-0.22465 (16)	0.35196 (16)	0.0559 (4)
O11	-0.1391 (2)	0.5708 (2)	0.2727 (2)	0.0809 (5)
O12	0.01425 (17)	0.18291 (17)	0.11287 (17)	0.0599 (4)
O13	0.05411 (16)	0.09520 (18)	0.55063 (15)	0.0572 (4)
O14	0.3357 (2)	0.4562 (2)	-0.06173 (17)	0.0813 (5)
H3	0.679 (2)	0.117 (2)	0.353 (2)	0.046 (5)*
H4	0.788 (2)	-0.112 (2)	0.276 (2)	0.048 (5)*
H5	0.629 (2)	-0.178 (2)	0.182 (2)	0.044 (5)*
H81	0.574 (3)	0.399 (3)	0.353 (3)	0.075 (8)*
H82	0.489 (3)	0.321 (3)	0.506 (3)	0.063 (6)*
H83	0.406 (3)	0.481 (3)	0.435 (2)	0.059 (6)*
H101	0.304 (3)	-0.215 (3)	0.049 (3)	0.081 (8)*
H102	0.143 (3)	-0.120 (3)	0.113 (3)	0.072 (7)*
H103	0.185 (3)	-0.288 (4)	0.196 (3)	0.093 (9)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Mn1	0.03310 (14)	0.02938 (14)	0.03785 (15)	-0.00427 (9)	-0.01615 (10)	-0.00960 (10)
C1	0.0313 (7)	0.0291 (7)	0.0266 (7)	-0.0100 (6)	-0.0102 (6)	-0.0042 (5)
C2	0.0357 (7)	0.0310 (7)	0.0297 (7)	-0.0133 (6)	-0.0121 (6)	-0.0039 (6)
C3	0.0358 (8)	0.0409 (9)	0.0417 (8)	-0.0121 (7)	-0.0175 (7)	-0.0069 (7)
C4	0.0341 (8)	0.0444 (9)	0.0480 (9)	-0.0038 (7)	-0.0168 (7)	-0.0115 (8)
C5	0.0402 (8)	0.0337 (8)	0.0406 (8)	-0.0032 (7)	-0.0123 (7)	-0.0129 (7)
C6	0.0374 (8)	0.0307 (7)	0.0289 (7)	-0.0109 (6)	-0.0115 (6)	-0.0068 (6)
C7	0.0448 (8)	0.0304 (7)	0.0302 (7)	-0.0155 (6)	-0.0148 (6)	-0.0030 (6)
C8	0.0582 (11)	0.0383 (9)	0.0444 (10)	-0.0193 (8)	-0.0241 (9)	-0.0089 (8)
C9	0.0356 (8)	0.0287 (7)	0.0500 (10)	-0.0049 (6)	-0.0148 (7)	-0.0124 (7)
C10	0.0623 (12)	0.0427 (10)	0.0631 (13)	-0.0209 (10)	-0.0232 (11)	-0.0170 (9)
C11	0.0477 (10)	0.0448 (10)	0.0624 (12)	0.0007 (8)	-0.0264 (9)	-0.0182 (9)
C12	0.0398 (8)	0.0392 (9)	0.0482 (9)	-0.0061 (7)	-0.0203 (7)	-0.0105 (7)
C13	0.0323 (8)	0.0479 (10)	0.0430 (9)	-0.0094 (7)	-0.0142 (7)	-0.0147 (8)
C14	0.0565 (10)	0.0435 (10)	0.0458 (10)	-0.0138 (8)	-0.0259 (9)	-0.0045 (8)
O1	0.0462 (6)	0.0299 (5)	0.0451 (6)	-0.0062 (5)	-0.0207 (5)	-0.0128 (5)
O2	0.0471 (6)	0.0323 (6)	0.0362 (6)	-0.0136 (5)	-0.0170 (5)	-0.0091 (4)
O3	0.0630 (8)	0.0504 (8)	0.0587 (8)	-0.0268 (7)	-0.0330 (7)	0.0092 (6)
O11	0.0606 (9)	0.0622 (10)	0.1086 (14)	0.0228 (8)	-0.0392 (9)	-0.0332 (10)
O12	0.0597 (8)	0.0649 (9)	0.0777 (10)	-0.0146 (7)	-0.0405 (8)	-0.0235 (8)
O13	0.0491 (7)	0.0742 (10)	0.0447 (7)	-0.0256 (7)	-0.0133 (6)	-0.0023 (7)
O14	0.0975 (13)	0.0914 (13)	0.0457 (9)	-0.0431 (11)	-0.0189 (8)	0.0090 (8)

supplementary materials

Geometric parameters (Å, °)

Mn1—C12	1.8046 (18)	C6—O2	1.4161 (18)
Mn1—C11	1.8367 (19)	C7—O1	1.243 (2)
Mn1—C14	1.849 (2)	C7—C8	1.490 (2)
Mn1—C13	1.8665 (18)	C8—H83	0.93 (2)
Mn1—C1	2.0443 (15)	C8—H82	0.93 (2)
Mn1—O1	2.0548 (11)	C8—H81	0.94 (3)
C1—C6	1.382 (2)	C9—O3	1.195 (2)
C1—C2	1.419 (2)	C9—O2	1.3611 (19)
C2—C3	1.395 (2)	C9—C10	1.493 (2)
C2—C7	1.453 (2)	C10—H101	0.94 (3)
C3—C4	1.376 (2)	C10—H102	0.91 (3)
C3—H3	0.90 (2)	C10—H103	0.94 (3)
C4—C5	1.391 (2)	C11—O11	1.131 (2)
C4—H4	0.97 (2)	C12—O12	1.150 (2)
C5—C6	1.383 (2)	C13—O13	1.138 (2)
C5—H5	0.97 (2)	C14—O14	1.133 (2)
C12—Mn1—C11	90.40 (8)	C4—C5—H5	122.5 (11)
C12—Mn1—C14	89.72 (8)	C1—C6—C5	123.10 (14)
C11—Mn1—C14	95.64 (9)	C1—C6—O2	119.06 (13)
C12—Mn1—C13	90.80 (8)	C5—C6—O2	117.74 (14)
C11—Mn1—C13	96.13 (8)	O1—C7—C2	117.28 (13)
C14—Mn1—C13	168.21 (8)	O1—C7—C8	119.36 (15)
C12—Mn1—C1	100.08 (7)	C2—C7—C8	123.35 (15)
C11—Mn1—C1	169.49 (7)	C7—C8—H83	110.0 (14)
C14—Mn1—C1	85.22 (7)	C7—C8—H82	112.0 (14)
C13—Mn1—C1	83.09 (6)	H83—C8—H82	106.7 (19)
C12—Mn1—O1	179.05 (6)	C7—C8—H81	112.6 (15)
C11—Mn1—O1	90.46 (7)	H83—C8—H81	110 (2)
C14—Mn1—O1	89.80 (7)	H82—C8—H81	105 (2)
C13—Mn1—O1	89.51 (6)	O3—C9—O2	122.64 (15)
C1—Mn1—O1	79.06 (5)	O3—C9—C10	126.36 (16)
C6—C1—C2	115.00 (13)	O2—C9—C10	111.00 (15)
C6—C1—Mn1	132.64 (11)	C9—C10—H102	110.0 (16)
C2—C1—Mn1	112.36 (11)	C9—C10—H103	111.9 (17)
C3—C2—C1	123.14 (14)	H102—C10—H103	108 (2)
C3—C2—C7	122.76 (14)	C9—C10—H101	108.0 (16)
C1—C2—C7	114.09 (13)	H102—C10—H101	106 (2)
C4—C3—C2	118.92 (15)	H103—C10—H101	113 (2)
C4—C3—H3	120.1 (13)	O11—C11—Mn1	178.71 (19)
C2—C3—H3	120.9 (13)	O12—C12—Mn1	177.51 (16)
C3—C4—C5	119.73 (15)	O13—C13—Mn1	175.05 (15)
C3—C4—H4	122.1 (12)	O14—C14—Mn1	176.86 (19)
C5—C4—H4	118.1 (12)	C7—O1—Mn1	117.18 (10)
C6—C5—C4	120.10 (15)	C9—O2—C6	117.20 (12)
C6—C5—H5	117.4 (11)		

Fig. 1

