

**Correspondence to:**

Professor W. Henderson,  
Department of Chemistry,  
University of Waikato,  
Private Bag 3105,  
Hamilton,  
New Zealand  
e-mail w.henderson@waikato.ac.nz  
FAX 0064-7-838-4219

**Spanning  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$  metalloligands with 1,4-dimercurated  
durene**

Nicola E. Cameron,<sup>a</sup> Rachael A. Linklater,<sup>a</sup> William Henderson,<sup>a,\*</sup> Brian K. Nicholson<sup>a</sup>  
and T. S. Andy Hor<sup>b</sup>

<sup>a</sup> *Department of Chemistry, University of Waikato, Private Bag 3105, Hamilton,  
New Zealand*

<sup>b</sup> *Department of Chemistry, National University of Singapore, 3 Science Drive 3,  
Singapore 117543.*

Received:

**Abstract** Reaction of the metalloligand  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$  with 0.5 mole equivalents of durene-1,4-bis(mercuric acetate)  $[\text{AcOHgC}_6\text{Me}_4\text{HgOAc}]$  in methanol gives the polynuclear complex  $[\{\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\}_2(\mu\text{-1,4-C}_6\text{Me}_4\text{Hg}_2)]^{2+}$ , isolated as its  $\text{PF}_6^-$  and

BPh<sub>4</sub><sup>-</sup> salts. Positive-ion ESI mass spectra indicate that [ $\{\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\}_2(\mu\text{-1,4-C}_6\text{Me}_4\text{Hg}_2)\text{]}^{2+}$  undergoes fragmentation by successive loss of PPh<sub>3</sub> ligands, while the ESI mass spectrum of the BPh<sub>4</sub><sup>-</sup> salt showed additional ions [ $\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4(\text{HgC}_6\text{Me}_4\text{HgPh})\text{]}^+$  and [ $\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\text{HgPh}\text{]}^+$  as a result of phenyl transfer from BPh<sub>4</sub><sup>-</sup> to Hg. A single-crystal X-ray structure determination on [ $\{\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\}_2(\mu\text{-1,4-C}_6\text{Me}_4\text{Hg}_2)\text{}(\text{BPh}_4)_2$ ] shows that the cation crystallises on a centre of symmetry, with structural features that are comparable to those of the previously characterised complex [ $\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\text{HgPh}\text{}]\text{BPh}_4$ .

**Keywords** Platinum complexes · Organomercury complexes · Sulfide complexes · Crystal structure

## Introduction

The ability of the platinum(II) sulfido complex [ $\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4$ ] **1** to act as a metalloligand is well established.<sup>1</sup> Closely related analogues with alternative phosphine ligands are known to behave in the same way.<sup>2,3,4,5</sup> Through such reactions, larger homo- and hetero-metallic sulfido-bridged aggregates of higher nuclearities can easily be assembled. We are interested in developing systematic methods for assembling larger assemblies containing  $\{\text{Pt}_2\text{S}_2\}$  cores as a basic building block, because such assemblies are relatively rare,<sup>2,6,7</sup> and reasoned that the use of organomercury complexes could offer a strategy to achieve this goal. A range of organomercury complexes has previously been reacted with [ $\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4$ ], using electrospray ionisation mass spectrometry (ESI MS) as a powerful tool to screen reactions for the successful formation of (cationic)

products.<sup>8</sup> Isolated complexes include the simple adducts  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\text{HgPh}]\text{BPh}_4$  **2**,<sup>8</sup>  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\text{HgEt}]\text{PF}_6$ ,<sup>8</sup>  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\text{HgFc}]\text{PF}_6$ ,<sup>9</sup> [Fc =  $\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\eta^5\text{-C}_5\text{H}_4)$ ], the dimercurated ferrocene derivative  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\text{Hg}_2\text{Fc}^*](\text{PF}_6)_2$ ,<sup>9</sup> [Fc\* =  $\text{Fe}(\eta^5\text{-C}_5\text{H}_4)_2$ ] and the organomercury adducts of the monobenzylated  $\{\text{Pt}_2\text{S}_2\}$  core,  $[\text{Pt}_2(\mu\text{-SCH}_2\text{Ph})(\mu\text{-SHgR})(\text{PPh}_3)_4](\text{PF}_6)_2$  (R = Ph, Fc).<sup>10</sup> In this paper we describe the reactivity of  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$  towards dimercurated durene (1,2,4,5-tetramethylbenzene),  $\text{AcOHgC}_6\text{Me}_4\text{HgOAc}$  **3**, which leads to the formation of hexametallic assemblies of two  $\{\text{Pt}_2\text{S}_2\}$  units spanned by the dimercurated arene moiety  $[\text{HgC}_6\text{Me}_4\text{Hg}]^{2+}$ . The availability of a wide range of poly-mercurated organic<sup>11,12</sup> and organometallic<sup>13,14</sup> compounds suggests that the assembly of even larger  $\{\text{Pt}_2\text{S}_2\}$ -Hg aggregates might ultimately be possible by this methodology.

## Results and discussion

The reaction of a methanol suspension of  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$  **1** with 0.5 mole equivalents of durene-1,4-bis(mercuric acetate) **3** leads to fairly rapid formation of a bright yellow solution which was shown by positive-ion ESI MS to contain almost exclusively the cation  $[\{\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\}_2(\mu\text{-1,4-C}_6\text{Me}_4\text{Hg}_2)]^{2+}$  [observed  $m/z$  1769.754, calculated  $m/z$  1769.755]. This complex contains two  $\{\text{Pt}_2\text{S}_2\}$  metalloligands bridged by the  $[\text{p-HgC}_6\text{Me}_4\text{Hg}]^{2+}$  fragment. The product was readily isolated from solution as its hexafluorophosphate salt **4a** by addition of excess  $\text{NH}_4\text{PF}_6$  to the filtered reaction solution. The positive-ion ESI mass spectrum of isolated **4a** in  $\text{CH}_2\text{Cl}_2\text{-CH}_3\text{OH}$  showed exclusively  $[\{\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\}_2(\mu\text{-1,4-C}_6\text{Me}_4\text{Hg}_2)]^{2+}$  at low cone voltages (up to 50 V), but as the cone voltage is increased, successive loss of  $\text{PPh}_3$  ligands occurs to give the

series of dications  $[M - nPPh_3]^{2+}$  (**n**,  $m/z$ : 0, 1769.5; **1**, 1638.5; **2**, 1507.5; **3**, 1376.5; **4**, 1245.5) at a cone voltage of 90 V. The complex shows a single resonance in its  $^{31}P$ - $\{^1H\}$  NMR spectrum, showing coupling to  $^{195}Pt$  of 3038 Hz, which is essentially the same as that observed for  $[Pt_2(\mu-S)_2(PPh_3)_4HgPh]PF_6$ .<sup>8</sup>

The  $[\{Pt_2(\mu-S)_2(PPh_3)_4\}_2(\mu-1,4-C_6Me_4Hg_2)]^{2+}$  cation can also be isolated as its tetraphenylborate salt **4b**, by addition of excess  $NaBPh_4$  to the reaction solution. However in this case, the isolated solid product was found to contain additional minor species when analysed by ESI MS. One species at  $m/z$  2113 (*ca.* 10% relative intensity) was not observed at all in the reaction solution. Exact mass measurement indicates that this species is the *phenyl*-mercury adduct  $[Pt_2(\mu-S)_2(PPh_3)_4(HgC_6Me_4HgPh)]^+$  **5** (observed  $m/z$  2113.326, calculated  $m/z$  2113.311). A number of other very minor species were observed in isolated **4b**, including the monocationic phenyl derivative  $[Pt_2(\mu-S)(\mu-SHgPh)(PPh_3)_4]^+$  **2** (observed  $m/z$  1781.247, calculated  $m/z$  1781.248). The source of the phenyl group is undoubtedly the  $BPh_4^-$  ion, since the phenylating ability of tetraphenylborate is well known,<sup>15,16,17,18</sup> including phenylation of mercury species, primarily for analytical purposes.<sup>19,20,21,22</sup> Consistent with this, the reaction of  $[Pt_2(\mu-S)_2(PPh_3)_4]$  with one equivalent of  $Hg(OAc)_2$  followed by precipitation with  $NaBPh_4$  gave a yellow product that showed **2** ( $m/z$  1781, 100% relative intensity) together with the known species<sup>23,24</sup>  $[\{Pt_2(\mu-S)_2(PPh_3)_4\}_2Hg]^{2+}$  ( $m/z$  1603.5, 85% relative intensity) as the only significant species. The isolated  $BPh_4^-$  salt **4b** also yielded an ion identified as the mono-mercurated species  $[Pt_2(\mu-S)_2(PPh_3)_4Hg(C_6Me_4H)]^+$  **6** (observed  $m/z$  1837.307, calculated  $m/z$  1837.310); this species was observed as a trace component of the reaction solution before precipitation, but as a more significant component of the

tetraphenylborate-precipitated solid. This species could arise from traces of mono-mercurated durene in the starting material, or from Hg-C bond cleavage in the product mediated by  $\text{BPh}_4^-$ . Recrystallisation of **4b** from dichloromethane-diethyl ether furnished a relatively pure sample of the compound.

In order to characterise the  $\{\text{Pt}_2\text{S}_2\}$ -Hg binding in this polymetallic system, and to compare it with the mono-mercury complex  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\text{HgPh}]^+$  **2**,<sup>8</sup> an X-ray structural determination of **4b** was carried out; the structure of the cation is shown in Figure 1, and selected bond lengths and angles are given in Table 1. The cation of complex **4b** crystallises on a centre of symmetry, so only half of the cation is unique. Overall, the structural features are similar to those of  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\text{HgPh}]^+$  **2**, with the mercury in each complex being strongly coordinated to one sulfur and weakly coordinated to the other, resulting in a ‘T-shaped’ coordination geometry. Thus, in **4b** the Hg-S(1) and Hg-S(2) bond distances are 2.3990(7) and 2.9473(7) Å, with an S(1)-Hg(1)-C(1) bond angle of 173.49(9)°, compared to the corresponding Hg-S bond distances of 2.4709(12) and 2.9286(12) Å, and an S-Hg-C bond angle of 174.59(16)° in **2**. The asymmetric mercury binding is, not unexpectedly, transmitted to the  $\{\text{Pt}_2\text{S}_2\}$  core of **4b**, with sulfur S(2) [which is more weakly coordinated to Hg] forming the shorter Pt-S bonds [Pt(1)-S(2) 2.3232(7) and Pt(2)-S(2) 2.3519(7) Å, compared with Pt(1)-S(1) 2.3675(7) and Pt(2)-S(1) 2.3797(7) Å]. Consequently, the Pt-P bonds *trans* to S(2) are slightly longer than those *trans* to S(1); for example Pt(1)-P(1) [*trans* to S(2)] is 2.2939(7) Å, while Pt(1)-P(2) [*trans* to S(1)] is 2.2661(7) Å. This indicates that sulfur S(2) displays a stronger *trans*-influence at the Pt centre than S(1). The same trend is observed in the phenyl derivative **2**.

The Hg $\cdots$ Pt non-bonded distances in **4b** [3.0707(1) and 3.5020(1) Å] can be compared with 3.078 and 3.515 Å in **2**, indicating that in both complexes the Hg-arene group is tilted towards one of the Pt(PPh<sub>3</sub>)<sub>2</sub> groups. The fold angle [angle between the two S-Pt-S planes] of the {Pt<sub>2</sub>S<sub>2</sub>} butterfly of **4b** [133.4°] is very similar to the corresponding angle of 132.9° in **2**. The durene ring is also twisted at an angle of 46.6° to the Hg-S(1)-S(2) plane, presumably to minimise steric interactions.

## Experimental

[Pt<sub>2</sub>( $\mu$ -S)<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>] **1** was prepared by the literature procedure from *cis*-[PtCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>] and Na<sub>2</sub>S $\cdot$ 9H<sub>2</sub>O in benzene suspension,<sup>25</sup> while the organomercury compound 1,4-AcOHgC<sub>6</sub>Me<sub>4</sub>HgOAc **3** was prepared by mercuration of 1,2,4,5-tetramethylbenzene (durene, Aldrich) with Hg(OAc)<sub>2</sub>, as described in the literature.<sup>26</sup> Sodium tetrphenylborate (BDH) and ammonium hexafluorophosphate (Aldrich) were used as supplied. Reactions were carried out in LR grade methanol, without exclusion of light, air or moisture.

Routine ESI mass spectra were recorded in positive-ion mode on a VG Platform II instrument using a methanol mobile phase. Isolated products were dissolved in a couple of drops of dichloromethane prior to dilution to *ca.* 0.1 mg mL<sup>-1</sup> with methanol. High resolution mass measurements were made on a Bruker Daltonics MicrOTOF instrument, calibrated using a sodium formate solution. <sup>31</sup>P-<sup>1</sup>H} NMR spectra were recorded on a Bruker AC300P instrument at 121.5 MHz in CDCl<sub>3</sub> solution, and spectra were referenced relative to external H<sub>3</sub>PO<sub>4</sub>. Elemental microanalyses were performed by

the Campbell Microanalytical Laboratory, Department of Chemistry, University of Otago, New Zealand.

#### **Synthesis of $[\{\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\}_2(\mu\text{-1,4-C}_6\text{Me}_4\text{Hg}_2)](\text{PF}_6)_2$ **4a****

A suspension of  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$  **1** (300 mg, 0.200 mmol) and **3** (65 mg, 0.100 mmol) in methanol (30 mL) was stirred at room temperature for 2 hours, giving a bright yellow solution. ESI MS showed a single peak due to the product cation  $[\{\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\}_2(\mu\text{-1,4-C}_6\text{Me}_4\text{Hg}_2)]^{2+}$ . After filtration to remove a trace of insoluble grey matter, solid  $\text{NH}_4\text{PF}_6$  (200 mg, 1.23 mmol) was added to the stirred filtrate, giving a yellow precipitate. Water (20 mL) was added to assist the precipitation, the product isolated by filtration, washed successively with water (2 x 20 mL) and diethyl ether (2 x 20 mL), and dried *in vacuo* to give **4a** (295 mg, 77%) as a bright yellow solid. Found: C, 47.7; H, 3.6.  $\text{C}_{154}\text{H}_{132}\text{F}_{12}\text{Hg}_2\text{P}_{10}\text{Pt}_4\text{S}_4$  requires C, 48.3; H, 3.5%.  $^{31}\text{P}\text{-}\{^1\text{H}\}$  NMR  $[(\text{CD}_3)_2\text{SO}]$ ,  $\delta$  20.1 [s,  $^1\text{J}(\text{PtP})$  3038 Hz].  $^1\text{H}$  NMR,  $\delta$  7.78-7.08 (m, Ph) and 2.57 (s,  $\text{CH}_3$ ).

#### **Synthesis of $[\{\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4\}_2(\mu\text{-1,4-C}_6\text{Me}_4\text{Hg}_2)](\text{BPh}_4)_2$ **4b****

Following the procedure for **4a**,  $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$  **1** (300 mg, 0.200 mmol) with **3** (65 mg, 0.100 mmol) were reacted. To the resulting yellow solution after filtration was added solid  $\text{NaBPh}_4$  (200 mg, 0.585 mmol), the product filtered, washed successively with methanol (20 mL), water (20 mL), methanol (20 mL) and diethyl ether (20 mL) and dried *in vacuo* to give **4b** (305 mg, 73%) as a yellow powder. Found: C, 57.2; H, 4.1.  $\text{C}_{202}\text{H}_{172}\text{B}_2\text{Hg}_2\text{P}_8\text{Pt}_4\text{S}_4$  requires C, 58.1; H, 4.2%.  $^{31}\text{P}\text{-}\{^1\text{H}\}$  NMR  $[(\text{CD}_3)_2\text{SO}]$ ,  $\delta$  21.3 [s,  $^1\text{J}(\text{PtP})$  3038 Hz].  $^1\text{H}$  NMR,  $\delta$  7.46-6.85 (m, Ph) and 2.54 (s,  $\text{CH}_3$ ).

### **Reaction of [Pt<sub>2</sub>(μ-S)<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>] with Hg(OAc)<sub>2</sub> and NaBPh<sub>4</sub>**

Mercury(II) acetate (21.6 mg, 0.068 mmol) was added to a stirred suspension of [Pt<sub>2</sub>(μ-S)<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>] (102 mg, 0.068 mmol) in methanol (20 mL), whereupon the reactants rapidly dissolved to give a clear yellow solution. After stirring for 1 h, NaBPh<sub>4</sub> (200 mg, 0.585 mmol) was added giving a yellow precipitate that was isolated by filtration, washed with water (10 mL) and methanol (10 mL) and dried *in vacuo* to give a yellow powder (105 mg). ESI MS analysis showed [Pt<sub>2</sub>(μ-S)<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>HgPh]<sup>+</sup> **2** (*m/z* 1781, 100%) and [{Pt<sub>2</sub>(μ-S)<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>]<sub>2</sub>Hg]<sup>2+</sup> (*m/z* 1603.5, 85%).

### **X-ray structure determination of [{Pt<sub>2</sub>(μ-S)<sub>2</sub>(PPh<sub>3</sub>)<sub>4</sub>]<sub>2</sub>(μ-1,4-C<sub>6</sub>Me<sub>4</sub>Hg<sub>2</sub>)](BPh<sub>4</sub>)<sub>2</sub>·4CH<sub>2</sub>Cl<sub>2</sub>**

Bright yellow crystals suitable for an X-ray structural study were obtained by vapour diffusion of diethyl ether into a dichloromethane solution of the complex at room temperature. Intensity data were obtained on a Bruker Apex II CCD diffractometer. Crystal, collection and refinement data are given in Table 2. The complex crystallises with four molecules of CH<sub>2</sub>Cl<sub>2</sub> per cation.

Crystallographic data have been deposited with the Cambridge Crystallographic Data Centre, CCDC No. 685996. Copies of this information can be obtained free of charge from The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (Fax: +44-1223-336033; e-mail deposit@ccdc.cam.ac.uk or www: <http://www.ccdc.cam.ac.uk>).



## **Acknowledgements**

We thank the University of Waikato (UW) and the National University of Singapore (NUS) for financial support of this work, including Summer Research Scholarships (NEC and RAL). We also thank Dr. Tania Groutso (University of Auckland) for collection of the X-ray data set.

**Table 1** Selected bond lengths (Å) and bond angles (°) for **4b**·4CH<sub>2</sub>Cl<sub>2</sub>

Pt(1)-P(1)	2.2939(7)	Pt(1)-P(2)	2.2661(7)
Pt(2)-P(3)	2.2767(7)	Pt(2)-P(4)	2.2913(8)
Pt(1)-S(1)	2.3675(7)	Pt(1)-S(2)	2.3232(7)
Pt(2)-S(1)	2.3797(7)	Pt(2)-S(2)	2.3519(7)
Hg(1)-C(1)	2.080(3)	Hg(1)-S(1)	2.3990(7)
Hg(1)-S(2)	2.9473(7)	C(1)-C(4)	1.399(4)
C(1)-C(2)	1.402(4)	C(2)-C(4')	1.406(4)
C(2)-C(3)	1.515(4)	C(4)-C(5)	1.509(4)
P(1)-Pt(1)-P(2)	98.84(3)	P(1)-Pt(1)-S(1)	87.60(2)
P(2)-Pt(1)-S(2)	90.62(3)	S(1)-Pt(1)-S(2)	83.31(2)
P(3)-Pt(2)-P(4)	102.29(3)	P(3)-Pt(2)-S(2)	83.66(3)
P(4)-Pt(2)-S(1)	91.65(3)	S(1)-Pt(2)-S(2)	82.44(2)
Pt(1)-S(1)-Pt(2)	86.16(2)	Pt(1)-S(2)-Pt(2)	87.82(2)
Pt(1)-S(1)-Hg(1)	94.56(3)	Pt(1)-S(2)-Hg(1)	82.36(2)
Pt(2)-S(1)-Hg(1)	79.97(2)	Pt(2)-S(2)-Hg(1)	69.794(18)
S(1)-Hg(1)-S(2)	70.50(2)	S(1)-Hg(1)-C(1)	173.49(9)
S(2)-Hg(1)-C(1)	112.13(8)	Hg(1)-C(1)-C(2)	119.4(2)
Hg(1)-C(1)-C(4)	119.3(2)		

**Table 2** Crystal, collection and refinement data for **4b**·4CH<sub>2</sub>Cl<sub>2</sub>*Crystal data*

Empirical formula	C <sub>206</sub> H <sub>180</sub> B <sub>2</sub> Cl <sub>8</sub> Hg <sub>2</sub> P <sub>8</sub> Pt <sub>4</sub> S <sub>4</sub>
Formula weight	4518.26
Crystal system	Monoclinic
Space group	P2 <sub>1</sub> /c

*Unit cell dimensions*





<i>a</i> (Å)	13.5909(1)
<i>b</i> (Å)	40.2714(4)
<i>c</i> (Å)	16.3746(2)
$\beta$ (°)	95.575(1)
<i>V</i> (Å <sup>3</sup> )	8919.83(16)
<i>Z</i>	2
<i>D</i> <sub>c</sub> (g cm <sup>-3</sup> )	1.682


*Data collection*

Crystal size (mm)	0.28 x 0.26 x 0.26
Radiation, wavelength (Å)	Mo-K $\alpha$ , $\lambda$ = 0.71073
Temperature (K)	89(2)
$\theta$ range for data collection (°)	1.81 to 32.62
Reflections collected	180006
Independent reflections	32403 [R <sub>int</sub> = 0.0407]
Absorption coefficient (mm <sup>-1</sup> )	5.133
T <sub>max,min</sub>	1.00, 0.69
<i>F</i> (000)	4436

*Structure analysis and refinement*

Solution by:	Direct methods
Refinement method	Full-matrix least-squares on <i>F</i> <sup>2</sup>
Data / restraints / parameters	32403 / 0 / 1056
Goodness-of-fit on <i>F</i> <sup>2</sup>	1.095
Final R indices [ <i>I</i> > 2 $\sigma$ ( <i>I</i> )]	<i>R</i> <sub>1</sub> 0.0328
	<i>wR</i> <sub>2</sub> 0.0662
R indices (all data)	<i>R</i> <sub>1</sub> 0.0465
	<i>wR</i> <sub>2</sub> 0.0697
Largest difference peak (e Å <sup>-3</sup> )	3.122
Largest difference hole (e Å <sup>-3</sup> )	-1.449





**Figure 1** Molecular structure of the cation of **4b**·4CH<sub>2</sub>Cl<sub>2</sub> with the atom numbering scheme. Phenyl rings of the triphenylphosphine ligands have been omitted for clarity.

## References

- <sup>1</sup> S.-W. A. Fong and T. S. A. Hor, *J. Chem. Soc., Dalton Trans.*, 1999, 639.
- <sup>2</sup> V. W.-W. Yam, K.-L. Yu, E. C.-C. Cheng, P. K.-Y. Yeung, K.-K. Cheung and N. Zhu, *Chem. Eur. J.*, 2002, **8**, 4122.
- <sup>3</sup> P. González-Duarte, A. Lledós and R. Mas-Ballesté, *Eur. J. Inorg. Chem.*, 2004, 3585.
- <sup>4</sup> H. Brunner, M. Weber and M. Zabel, *J. Organomet. Chem.*, 2003, **684**, 6.
- <sup>5</sup> Z. Li, K. F. Mok and T. S. A. Hor, *J. Organomet. Chem.*, 2003, **682**, 73.
- <sup>6</sup> Z. Li, W. Zhang, H. Liu, K. F. Mok and T. S. A. Hor, *Inorg. Chem.*, 2003, **42**, 8481.
- <sup>7</sup> R. Mas-Ballesté, W. Clegg, A. Lledós and P. González-Duarte, *Eur. J. Inorg. Chem.*, 2004, 3223.
- <sup>8</sup> S.-W. A. Fong, W. T. Yap, J. J. Vittal, T. S. A. Hor, W. Henderson, A. G. Oliver and C. E. F. Rickard, *J. Chem. Soc., Dalton Trans.*, 2001, 1986.
- <sup>9</sup> X. Xu, S.-W. A. Fong, Z. Li, Z.-H. Loh, F. Zhao, J. J. Vittal, W. Henderson, S.-B. Khoo and T. S. A. Hor, *Inorg. Chem.*, 2002, **41**, 6838.
- <sup>10</sup> W. Henderson, B. K. Nicholson, S. M. Devoy and T. S. A. Hor, *Inorg. Chim. Acta*, in press.
- <sup>11</sup> R. J. Bertino and G. B. Deacon, *J. Organomet. Chem.*, 1974, **67**, C61.
- <sup>12</sup> D. Grdenić, B. Kamenar, B. Korpar-Čolig, M. Sikirica and G. Jovanovski, *J. Chem. Soc., Chem. Comm.*, 1974, 646.
- <sup>13</sup> S. A. Kur, M. J. Heeg and C. H. Winter, *Organometallics*, 1994, **13**, 1865.
- <sup>14</sup> Y.-H. Han, M. J. Heeg and C. H. Winter, *Organometallics*, 1994, **13**, 3009
- <sup>15</sup> N. E. Leadbeater and M. Marco, *J. Org. Chem.*, 2003, **68**, 5660.
- <sup>16</sup> M. Ueda and N. Miyaura, *J. Organomet. Chem.*, 2000, **595**, 31.

- 
- <sup>17</sup> M. Aresta, E. Quaranta and I. Tommasi, *New J. Chem.*, 1997, **21**, 595.
- <sup>18</sup> A. Sladek, S. Hofreiter, M. Paul and H. Schmidbaur, *J. Organomet. Chem.*, 1995, **501**, 47.
- <sup>19</sup> Y. Cai, S. Monsalud and K. G. Furton, *Chromatographia*, 2000, **52**, 82.
- <sup>20</sup> P. Grinberg, R. C. Campos, Z. Mester and R. E. Sturgeon, *J. Analyt. Atom. Spectrom.*, 2003, **18**, 902.
- <sup>21</sup> S. Mishra, R. M. Tripathi, S. Bhalke, V. K. Shukla and V. D. Puranik, *Anal. Chim. Acta*, 2005, **551**, 192.
- <sup>22</sup> Y. Cai, S. Monsalud, R. Jaffé and R. D. Jones, *J. Chromatogr. A*, 2000, **876**, 147.
- <sup>23</sup> C. E. Briant, T. S. A. Hor, N. D. Howells and D. M. P. Mingos, *J. Chem. Soc., Chem. Commun.*, 1983, 1118.
- <sup>24</sup> Z. Li, X. Xu, S. B. Khoo, K. F. Mok and T. S. A. Hor, *J. Chem. Soc., Dalton Trans.*, 2000, 2901.
- <sup>25</sup> R. Ugo, G. La Monica, S. Cenini, A. Segre and F. Conti, *J. Chem. Soc. A*, 1971, 522.
- <sup>26</sup> H. Sawatzky and G. F. Wright, *Can. J. Chem.*, 1958, **36**, 1555.