

Role of Tephra in Dating Polynesian Settlement and Impact, New Zealand

D.J. Lowe^{1*} and R.M. Newnham²

¹Department of Earth Sciences, University of Waikato, Private Bag 3105, Hamilton, New Zealand (d.lowe@waikato.ac.nz)

²School of Geography, University of Plymouth, Plymouth, Devon PL4 8AA, U.K.

*INQUA Sub-Commission for Tephrochronology and Volcanism (SCOTAV)

Introduction

Tephrochronology in its original sense is the use of tephra layers as time-stratigraphic marker beds to establish numerical or relative ages (Lowe and Hunt, 2001). Tephra layers have been described and studied in New Zealand for more than 160 years (the German naturalist and surgeon Ernst Dieffenbach described 'recognizable' tephra sections in his 1843 book *Travels in New Zealand*), and the first isopach map, showing fallout from the deadly plinian basaltic eruption of Mt Tarawera on 10 June 1886, was published in 1888 (Lowe, 1990; Lowe *et al.*, 2002). More recently, a wide range of tephra-related paleoenvironmental research has been undertaken (*e.g.*, Lowe and Newnham, 1999; Newnham and Lowe, 1999; Newnham *et al.*, 1999, 2004; Shane, 2000), including new advances in the role of tephra in linking and dating sites containing evidence for abrupt climatic change (*e.g.*, Newnham and Lowe, 2000; Newnham *et al.*, 2003). Here we focus on the use of tephrochronology in dating the arrival and impacts of the first humans in New Zealand, a difficult problem for which this technique has proven to be of critical importance.

Heated debate

The timing of initial settlement of New Zealand has been the subject of heated debate. An early but transient contact *c.* AD 50 to 150, based on Pacific rat-bone (*Rattus exulans*) dates obtained from natural sites, was proposed by Holdaway (1996) on the premise that the rats, an introduced predator to New Zealand, accompanied the early Polynesians as a food source or as stowaways (Matisoo-Smith, 2002; Matisoo-Smith and Robins, 2004). This proposal seemingly supported Sutton (1987, 1994), who first suggested, on the basis of small-scale but short-lived disturbance evident in pollen records, that 'archaeologically invisible' Polynesian sailors may have reached New Zealand around this time. However, the reliability of the early rat-bone dates has been disputed, especially as aberrant rat-bone dates were reported from several archeological sites (Anderson, 1996, 2000, 2004; Higham and Petchey, 2000; Higham *et al.*, 2004), and dates on rat-nibbled land snail shells (*Placostylus ambagiosus*) and rat-nibbled seeds both suggested instead that the Pacific rat became established after *c.* AD 1250 (Brook, 2000; Wilmshurst and Higham, 2004). Moreover, the early rat-bone dates at one of Holdaway's (1996) sites in the South Island have not been duplicated (Holdaway *et al.*, 2002; Anderson and Higham, 2004).

Using tephra layers to date archeological and paleoenvironmental sites

Because tephras provide essentially instantaneous chronostratigraphic marker horizons, or *isochrons*, that can be correlated between sites independently of radiometric dating, they provide a way of circumventing the various interpretative difficulties associated with radiocarbon dating very recent (last millennia) archeological and paleoenvironmental (natural) sites. Because tephra deposits are found in both archeological and natural sites, they have the capacity for linking such sites in an unambiguous manner unparalleled by other dating or correlative techniques (Lowe *et al.*, 2000).

Direct links between early Polynesians and their descendants (Maori) and tephra in New Zealand are associated with three different eruptive centres on North Island (Lowe *et al.*, 2002).

- (1) Human footprints and other artifacts are buried beneath and within basaltic ash erupted from Rangitoto Island volcano, near Auckland, at *c.* AD 1400 (Fig. 1).
- (2) The remains of Maori cooking stones (*umu*) aged *c.* AD 1450–1500 lie sandwiched between tephra on the slopes of the andesitic stratovolcano of Taranaki (Mt Egmont) in western North Island (Fig. 2).
- (3) The key event for dating Polynesian settlement in New Zealand was the eruption of Kaharoa Tephra, a geochemically distinctive, rhyolitic tephra layer originating from Mt Tarawera volcano in central North Island near Rotorua (Lowe *et al.*, 2000). Widely dispersed over > 30,000 km² of northern and eastern North Island, Kaharoa Tephra provides a unique ‘settlement layer’ (*landnámslag*) (*cf.* Wastegard *et al.*, 2003; Sveinbjornsdottir *et al.*, 2004) in northern New Zealand. Difficult to date accurately by radiocarbon alone because of wiggles in the calibration curves (Lowe *et al.*, 1998), we derived a wiggle-match date of AD 1314 ± 12 for the Kaharoa Tephra eruption—the main plinian, tephra-fall-producing phase of which occurred in the austral winter (Hogg *et al.*, 2003)—using a carbonized log of *Phyllocladus* spp.. We used the Southern Hemisphere calibration curves of Hogg *et al.* (2002), thus avoiding the interhemispheric offset problem, and confirmed our date’s likelihood using Bayesian statistics (Buck *et al.*, 2003).

Numerous archeological sites in eastern and northern North Island contain the Kaharoa Tephra datum (Fig. 3; Lowe *et al.*, 2000), and the absence of artifacts or cultural remains reported beneath it thus far indicates that these sites must be younger than *c.* AD 1314. In the same region, nearly 20 pollen profiles obtained from peat or lake deposits contain both Kaharoa Tephra and palynological indicators for the onset of significant deforestation, in the form of both marked and sustained rises in bracken (*Pteridium*) spores and charcoal, and a concomitant decline in tall forest trees (Newnham *et al.*, 1998; Horrocks *et al.*, 2001, 2004; McGlone and Jones, 2004). Unprecedented in the Holocene record, these palynological changes are inferred to be the result of initial and repeated firing by early Maori (Ogden *et al.*, 1998; McGlone and Wilmshurst, 1999; *cf.* Flenley and Todd, 2001). In a few profiles, the sustained rises in bracken and charcoal occur well after the Kaharoa Tephra datum but in the others, they occur close to the time of its deposition. In four pollen profiles, the earliest sustained rises are recorded in sediments just a few centimetres below the Kaharoa Tephra layer, probably ≤ 50 years before the eruption of *c.* AD 1314 (Lowe *et al.*, in press). A similar pattern is evident from independent opal phytolith data in tephra-soil sequences in the Bay of Plenty (Kondo *et al.*, 1994; Sase and Hosono, 1996). Thus, it is likely that the Kaharoa eruption was witnessed by a small number of very early Maori (an argument supported by oral tradition) and that archeological sites containing artifacts just beneath the Kaharoa Tephra may yet be found.

Conclusions

Taken together, both the archeological and palynological evidence, constrained by the *c.* AD 1314 Kaharoa Tephra datum, suggests that the earliest environmental impacts associated with initial Polynesian settlement in northern New Zealand (North Island) occurred between *c.* AD 1250 and 1300. This is coincident with the earliest-known settlement dates from archeological remains on both North and South islands, with dates obtained from rat-nibbled snail shells and seeds, and with reliably-dated deforestation signals (Figs. 4 & 5). The fact that the maximum date for the onset of deforestation is similar to dates obtained from the oldest-known archeological sites (*e.g.*, Wairau Bar), implies that the onset of forest burning was more-or-less contemporaneous with initial settlement (Lowe *et al.*, in press). It remains feasible that earlier settlement sites still await discovery beyond the

fall-out zone of macroscopic Kaharoa Tephra and that there might have been an earlier transient contact. If such transient contact occurred, it currently remains invisible in the archeological record and is indistinguishable from natural background events in the palynological record (Lowe *et al.*, in press).

Acknowledgements

Many colleagues (including those with differing viewpoints) have worked with us over the past decade to help solve the problem of timing of Polynesian settlement and impact in New Zealand and we acknowledge all their contributions. Drs Reg Nichol and Brent Alloway kindly provided photographs, Dr Phil Moore alerted us to the Waihi Beach peat sections, Will Esler told us about Dieffenbach's writings, and Betty-Ann Kamp drafted Figure 4. We especially thank the editors for their encouragement and technical support in preparing our article.

References

- Hogg, A.G., Higham, T.F.G., Lowe, D.J., Palmer, J., Reimer, P. and Newnham, R.M. 2003: A wiggle-match date for Polynesian settlement of New Zealand. *Antiquity*, **77**, 116-125.
- Higham, T.F.G., Hedges, R.E.M., Anderson, A.J., Bronk Ramsey, C. and Fankhauser, B. 2004: Problems associated with the AMS dating of small bone samples: the question of the arrival of Polynesian rats to New Zealand. *Radiocarbon*, **46**, 207-218.
- Lowe, D.J., Newnham, R.M., McFadgen, B.G. and Higham, T.F.G. 2000: Tephra and New Zealand archaeology. *Journal of Archaeological Science*, **27**, 859-870.
- Lowe, D.J., Newnham, R.M. and McCraw, J.D. 2002: Volcanism and early Maori society in New Zealand. In: *Natural Disasters and Cultural Change*; Torrence, R. and Grattan, J. (eds), Routledge, 126-161.
- Wilmshurst, J.M. and Higham, T.F.G. 2004: Using rat-gnawed seeds to independently date the arrival of Pacific rats and humans in New Zealand. *The Holocene*, **14**, 801-806.

For full references please see below (or web site: www.pages-igbp.org/products/newsletters/ref2004_3.html).

Captions to figures

Fig. 1: Human footprint in basaltic Rangitoto Tephra, erupted *c.* AD 1400, Auckland, northern North Island (Nichol, 1982). The age of the eruption is derived from multiple radiocarbon, paleomagnetic, thermoluminescence, and obsidian hydration dates (Lowe *et al.*, 2000). Photo courtesy of Reg Nichol.

Fig. 2: Early Maori cooking stones (*umu*) sandwiched between andesitic tephra layers on Taranaki volcano, western North Island. At this site, the stones are overlain directly by Burrell Ash and underlain by Waiweranui Lapilli, dating the *umu* to *c.* AD 1500 or a little before (Alloway *et al.*, 1990; Lowe *et al.*, 2000). First recognized by Oliver (1931), the oldest *umu* on Taranaki is dated at *c.* AD 1450. Photo courtesy of Brent Alloway.

Fig. 3: Distal rhyolitic Kaharoa Tephra showing up as a prominent, 5-cm-thick marker layer in shallow peat deposits at Waihi Beach, western Bay of Plenty, eastern North Island. Taupo Tephra (erupted *c.* AD 232; Sparks *et al.* 1995; Lowe and de Lange, 2000) occurs also in the section, faintly visible as fine lapilli on the 'corner' in the middle of the photo, a few centimetres above the pale muds near the water table. The peat above Kaharoa datum is darker than below because it contains abundant charcoal from Polynesian burning, which is documented by pollen analysis in this area (Newnham *et al.*, 1995). Cutting tool handle is ~30 cm long. Photo: David Lowe.

Fig. 4: Summary of stratigraphy and ages of tephras, erupted from five volcanic centres since *c.* AD 232 (left side of diagram), and their relationship with archaeological and deforestation signals in northern and eastern North Island (right) (after Lowe *et al.*, 2000, 2002). The Kaharoa Tephra provides a settlement datum, or *landnám*slag, for inferred human-induced burning and deforestation

in much of northern and eastern North Island (*e.g.*, Newnham *et al.*, 1998; Horrocks *et al.*, 2001). It matches the earliest settlement dates of *c.* AD 1250–1300 from many sites containing archaeological remains (*e.g.*, Anderson, 1991; Higham and Hogg, 1997; McFadgen, 2003; Higham *et al.*, 2004), including the ancient Wairau Bar artifacts and skeletons (Higham *et al.*, 1999) and the tropical pearl lure at Tairua (Schmidt and Higham, 1998) (2 sigma error ranges), the oldest known rat-nibbled snail shells and seeds (Brook, 2000; Wilmshurst and Higham, 2004), and the earliest reliable dates for sustained deforestation elsewhere in the New Zealand archipelago (Ogden *et al.*, 1998; McGlone and Wilmshurst, 1999). The zone depicting possible early transient human contact is based on Holdaway (1996, 1999). Dutchman, Abel Tasman, was probably the first European to visit New Zealand (AD 1642).

Fig. 5: Archeological excavation of an early Maori village (*kainga*) site on dunes at Papamoa, coastal Bay of Plenty, eastern North Island, with the Kaharoa Tephra (dated at AD 1314 ± 12 by Hogg *et al.*, 2003) forming a prominent white marker layer in peat. Photo: David Lowe.

Full references

- Alloway, B.V., Fyfe, R. and Neall, V.E., 1990: Further evidence of early Maori occupation on the flanks of Egmont volcano. *Archaeology in New Zealand*, **33**, 40-48.
- Anderson, A.J., 1991: The chronology of colonisation in New Zealand. *Antiquity*, **65**, 767-795.
- Anderson, A.J., 1996: Was *Rattus exulans* in New Zealand 2000 years ago? AMS radiocarbon ages from Shag River Mouth. *Archaeology in Oceania*, **31**, 178-184.
- Anderson, A.J., 2000: Differential reliability of ¹⁴C AMS ages of *Rattus exulans* bone gelatin in south Pacific prehistory. *Journal of the Royal Society of New Zealand*, **30**, 243-261.
- Anderson, A.J., 2004: The age discomformity in AMS radiocarbon results on *Rattus exulans* bone. *New Zealand Journal of Archaeology*, **24** (2002), 149-156.
- Anderson, A.J. and Higham, T.F.G., 2004: The age of rat introduction in New Zealand: further evidence from Earthquakes #1, North Otago. *New Zealand Journal of Archaeology*, **24** (2002), 135-147.
- Brook, F.J., 2000: Prehistoric predation of the landsnail *Placostylus ambagiosus* Suter (Stylommatophora: Bulimulidae), and evidence for the timing of establishment of rats in northernmost New Zealand. *Journal of the Royal Society of New Zealand*, **30**, 227-241.
- Buck, C.E., Higham, T.F.G. and Lowe, D.J., 2003: Bayesian tools for tephrochronology. *The Holocene*, **13**, 639-647.
- Flenley, J.R. and Todd, A., 2001: A genome or memome: the cause of the rise in bracken fern at around 700 BP in New Zealand. *Proceedings of the 7th Australasian Archaeometry Conference*, 141-145.
- Higham, T.F.G. and A.G. Hogg, 1997: Evidence for late Polynesian colonisation of New Zealand: University of Waikato radiocarbon measurements. *Radiocarbon*, **39**, 149-192.
- Higham, T.F.G. and F.J. Petchey, 2000: On the reliability of archaeological rat bone for radiocarbon dating in New Zealand. *Journal of the Royal Society of New Zealand*, **30**, 399-409.
- Higham, T.F.G., Anderson, A.J. and Jacomb, C. 1999: Dating the first New Zealanders: the chronology of Wairau Bar. *Antiquity*, **73**, 420-427.
- Higham, T.F.G., Hedges, R.E.M., Anderson, A.J., Bronk Ramsey, C. and Fankhauser, B., 2004: Problems associated with the AMS dating of small bone samples: the question of the arrival of Polynesian rats to New Zealand. *Radiocarbon*, **46**, 207-218.
- Hogg, A.G., McCormac, F.G., Higham, T.F.G., Reimer, P.J., Baillie, M.G.L., Palmer, J.G. and Stuiver, M., 2002. High-precision ¹⁴C measurements of contemporaneous tree-ring dated wood from the British Isles and New Zealand: AD 1850 – 950. *Radiocarbon*, **44**, 633-640.

- Hogg, A.G., Higham, T.F.G., Lowe, D.J., Palmer, J., Reimer, P. and Newnham, R.M., 2003: A wiggle-match date for Polynesian settlement of New Zealand. *Antiquity*, **77**, 116-125.
- Holdaway, R.N., 1996: Arrival of rats in New Zealand. *Nature*, **384**, 225-226.
- Holdaway, R.N., 1999: A spatio-temporal model for the invasion of the New Zealand archipelago by the Pacific rat *Rattus exulans*. *Journal of the Royal Society of New Zealand*, **29**, 91-105.
- Holdaway, R.N., Roberts, R.G., Beavan-Athfield, N.R., Olley, J.M. and Worthy, T.H., 2002: Optical dating of quartz sediments and accelerator mass spectrometry ¹⁴C dating of bone gelatin and moa eggshell: a comparison of age estimates for non-archaeological deposits in New Zealand. *Journal of the Royal Society of New Zealand*, **32**, 463-505.
- Horrocks, M., Deng, Y., Ogden, J., Alloway, B.V., Nichols, S. and Sutton, D.G., 2001: High spatial resolution of pollen and charcoal in relation to the c. 600 year BP Kaharoa Tephra: implications for Polynesian settlement of Great Barrier Island New Zealand. *Journal of Archaeological Science*, **28**, 153-168.
- Horrocks, M., Shane, P.A.R., Barber, I.G., D'Costa, D.M. and Nichol, S.L., 2004: Microbotanical remains reveal Polynesian agriculture and mixed cropping in early New Zealand. *Review of Palaeobotany and Palynology*, **131**, 147-157.
- Kondo, R., Childs, C.W. and Atkinson, I., 1994: *Opal Phytoliths in New Zealand*. Manaaki Whenua Press, 85p.
- Lowe, D.J., 1990: Tephra studies in New Zealand: an historical review. *Journal of the Royal Society of New Zealand*, **20**, 119-150.
- Lowe, D.J. and de Lange, W.P., 2000: Volcano-meteorological tsunamis, the c. AD 200 Taupo eruption (New Zealand) and the possibility of a global tsunami. *The Holocene*, **10**, 401-407.
- Lowe, D.J. Hunt, J.B., 2001: A summary of terminology used in tephra-related studies. In: *Tephra: Chronology, Archaeology*; Juvigné, E.T. and Raynal, J.-P. (eds), *Les Dossiers de l'Archéologie*, **1**, 17-22.
- Lowe, D.J. and Newnham, R.M., 1999: Advances in Quaternary tephrostratigraphy and tephrochronology in New Zealand. *Quaternary Australasia*, **17**, 12-19.
- Lowe, D.J., McFadgen, B.G., Higham, T.F.G., Hogg, A.G., Froggatt, P.C. and Nairn, I.A., 1998: Radiocarbon age of the Kaharoa Tephra, a key marker for late-Holocene stratigraphy and archaeology in New Zealand. *The Holocene*, **8**, 499-507.
- Lowe, D.J., Newnham, R.M., McFadgen, B.G. and Higham, T.F.G., 2000: Tephra and New Zealand archaeology. *Journal of Archaeological Science*, **27**, 859-870.
- Lowe, D.J., Newnham, R.M. and McCraw, J.D., 2002: Volcanism and early Maori society in New Zealand. In: *Natural Disasters and Cultural Change*; Torrence, R. and Grattan, J. (eds), Routledge, 126-161.
- Lowe, D.J., Newnham, R.M., Higham, T.F.G., Wilmshurst, J.M., McGlone, M.S. and Hogg, A.G., in press: Dating earliest human impact and settlement in New Zealand. In: *Origins of the First New Zealanders 2nd edition*; Sutton, D.G. (ed), Auckland University Press.
- Matisoo-Smith, E., 2002: Something old, something new: do genetic studies of contemporary populations reliably represent prehistoric populations of Pacific *Rattus exulans*? *Human Biology*, **74**, 489-496.
- Matisoo-Smith, E. and Robins, J.H. 2004: Origin and dispersals of Pacific peoples: Evidence from mtDNA phylogenies of the Pacific rat. *Proceedings of the National Academy of Sciences*, **101**, 9167-9172.
- McFadgen, B.G., 2003: Pre-European archaeology of the coast. In: *The New Zealand Coast*; Goff, J.G., Nichol, S.L. and Rouse, H.L. (eds), Dunmore Press, 265-296.
- McGlone, M.S. and Jones, K.L., 2004: The impact of Polynesian settlement on the vegetation of the coastal Bay of Plenty. In: *Kohika*; Irwin, G. (ed), Auckland University Press, 20-44.
- McGlone, M.S. and Wilmshurst, J.M., 1999: Dating initial Maori environmental impact in New Zealand, *Quaternary International*, **59**, 5-16.

- Newnham, R.M. and Lowe, D.J., 1999: Testing the synchronicity of pollen signals using tephrostratigraphy. *Global and Planetary Change*, **21**, 113-128.
- Newnham, R.M. and Lowe, D.J., 2000: Fine-resolution pollen record of late-glacial climate reversal from New Zealand. *Geology*, **28**, 759-762.
- Newnham, R.M., Lowe, D.J. and Wigley, G.N.A., 1995: Late Holocene palynology and palaeovegetation of tephra-bearing mires at Papamoa and Waihi Beach, western Bay of Plenty, North Island, New Zealand. *Journal of the Royal Society of New Zealand*, **25**, 283-300.
- Newnham, R. M., Lowe, D.J. and Williams, P.W., 1999: Quaternary environmental change in New Zealand: a review. *Progress in Physical Geography*, **23**, 567-610.
- Newnham, R.M., Lowe, D.J., McGlone, M.S., Wilmshurst, J.M. and Higham, T.F.G., 1998: The Kaharoa Tephra as a critical datum for earliest human impact in northern New Zealand. *Journal of Archaeological Science*, **25**, 533-544.
- Newnham, R.M., Eden, D.N., Lowe, D.J. and Hendy, C.H., 2003: Rerewhakaaitu Tephra, a land-sea marker for the Last Termination in New Zealand, with implications for global climate change. *Quaternary Science Reviews*, **22**, 289-308.
- Newnham, R.M., Lowe, D.J., Green, J.D., Turner, G.M., Harper, M.A., McGlone, M.S., Stout, S.L., Horie, S. and Froggatt, P.C., 2004: A discontinuous c. 80 ka record of Late Quaternary environmental change from Lake Omapere, Northland, New Zealand. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **207**, 165-198.
- Nichol, R., 1982: Fossilised human footprints in Rangitoto Ash on Motutapu Island. *Geological Society of New Zealand Newsletter*, **51**, 11-13.
- Ogden, J., Basher, L.R. and McGlone, M.S., 1998: Fire, forest regeneration and links with early human habitation: evidence from New Zealand. *Annals of Botany*, **81**, 687-696.
- Oliver, W.R.B., 1931: An ancient Maori oven on Mount Egmont. *Journal of the Polynesian Society*, **40**, 73-79.
- Sase, T. and Hosono, M., 1996: Vegetation histories of Holocene volcanic ash soils in Japan and New Zealand – relationship between genesis of melanic volcanic ash soils and human impact. *Earth Science (Chikyu Kagaku)*, **50**, 466-482.
- Schmidt, M. and Higham, T.F.G., 1998: Sources of New Zealand's East Polynesian culture revisited: the radiocarbon chronology of the Tairua archaeological site, New Zealand. *Journal of the Polynesian Society*, **107**, 395-403.
- Shane, P.A.R., 2000: Tephrochronology: a New Zealand case study. *Earth-Science Reviews*, **49**, 223-259.
- Sparks, R.J., Melhuish, W.H., McKee, J.W.A., Ogden, J. and Palmer, J.G., 1995: ¹⁴C calibration in the Southern Hemisphere and the date of the last Taupo eruption: evidence from tree-ring sequences. *Radiocarbon*, **37**, 155-163.
- Sutton, D.G., 1987: A paradigmatic shift in Polynesian prehistory: implications for New Zealand. *New Zealand Journal of Archaeology*, **9**, 135-155.
- Sutton, D.G., 1994: Conclusions: origins. In: *Origins of the First New Zealanders*; Sutton, D.G. (ed), Auckland University Press, 243-258.
- Sveinbjörnsdóttir, A.E., Heinemeier, J. and Gudmundsson, G., 2004: ¹⁴C dating of settlement of Iceland. *Radiocarbon*, **46**, 387-394.
- Wastegard, S., Hall, V.A., Hannon, G.E., van den Bogaard, C., Pilcher, J.R., Sigurgeirsson, M.A. and Hermanns-Audardottir, M., 2003: Rhyolitic tephra horizons in northwestern Europe and Iceland from the AD 700s-800s: a potential alternative for dating first human impact. *The Holocene*, **13**, 277-283.
- Wilmshurst, J.M. and Higham, T.F.G., 2004: Using rat-gnawed seeds to independently date the arrival of Pacific rats and humans to New Zealand. *The Holocene*, **14**, 801-806.



Fig.1



Fig.2



Fig.3




Fig.4



Fig. 5