

INTRODUCTION TO TEPHRA-DERIVED SOILS AND FARMING, WAIKATO-BAY OF PLENTY, NORTH ISLAND, NEW ZEALAND



NEW ZEALAND WINTERIM 2019

*University of Wisconsin–Platteville, USA, and University of
Waikato, New Zealand, 15-17 January 2019*

David J. Lowe and Megan R. Balks
School of Science (Earth Sciences)
University of Waikato, Hamilton



UNIVERSITY OF WISCONSIN
PLATTEVILLE

Note about ages: Ma = millions of years ago; ka = thousands of years ago; BP = before present, 'present' being defined as AD [CE] 1950 for radiocarbon-derived ages. Radiocarbon (^{14}C) ages are given in calendar or calibrated (cal) years BP unless noted otherwise, or as calendrical dates (e.g. AD 232). The Holocene encompasses the last 11,700 cal years (Walker et al., 2009).

Photo on front cover (by Maria Lowe): Mt Pirongia and tephra-draped landscape, Waikato region

Citation: Lowe, D.J., Balks, M.R. 2019. Introduction to tephra-derived soils and farming, Waikato-Bay of Plenty, North Island, New Zealand. University of Wisconsin-Platteville Education Abroad Program: Winterim Field Trip (15-17 January, 2019). *School of Science (Earth Sciences), University of Waikato, Hamilton*. 82 pp.

Provisional itinerary

Tuesday 15 January 2019: Farming and conservation

8.00 am Depart Bryant Hall

8.00-8.45 am Drive to Ormsby Farm, Pekanui Rd, Ngutunui (Pirongia Mtn)

Meet Megan at Ngutunui School

9.00-10.30 am Keith and Margaret Ormsby dairy farm, Pekanui Rd

10.30-11.00 am Drive to Otorohanga

11.00-12.00 Otorohanga Kiwi house and bird sanctuary

12.00-12.45 Lunch at Ronnies Café, Otorohanga

12.45-1.30 Drive to Garland Farm, Rahiri Rd, Roto-o-rangi (Maungatautari Mtn)

1.30-3.30 Bill and Sue Garland sheep and beef farm

3.30-4.15 Special afternoon tea at nearby farm, Cambridge Rd (US connection)

4.15-4.30 Drive to Leamington section

4.30-5.00 Horotiu soil on late Quaternary tephra mantle (allophanic) on volcanogenic alluvium, Leamington (Vitric Hapludand)

5.00-5.30 Return to Bryant Hall, Knighton Rd, Hamilton

Wednesday 16 January 2019: Rotorua geysers and tephras

8.30 am Depart Bryant Hall

8.30-10.00 Drive to Te Puia, Hemo Rd, Rotorua

10.00 Arrival at Te Puia

10.15 Concert

11.00 Guided tour

12.15-1.00 Lunch at Te Puia or local eatery (e.g. BP Connect)

1.00-1.30 Drive to Brett Rd, Lake Rerewhakaaitu

1.30-2.30 Rotomahana soil on layered Holocene tephras, buried soils (Typic Udivitrand)

2.30-2.45 Circumnavigate Lake Rerewhakaaitu (view Mt Tarawera)

2.45-3.15 Drive to Okareka Loop Rd tephra section

3.15-3.45 Okareka Loop Rd tephra section

3.45-4.30 Redwood forest (Long Mile Drive)

4.30-6.00 Return to Bryant Hall, Knighton Rd, Hamilton

Thursday 17 January 2019: Waikato peats and gardens

8.30 am Departure

8.30-8.45 Drive to Raynes Rd

8.45-9.30 Kainui soil, late Quaternary tephra mantle (halloysitic) on buried clayey paleosol (Typic Kandiodult)

9.30-9.45 Drive to Moanatuatua bog and blueberry orchard, Jary Rd, Ohaupo

9.45-10.15 Rukuhia soil on peat with thin tephra layer (Taupo Tephra 232 ± 10 CE) (Sapric or Typic Haplohemist)

10.15-10.20 Drive to The Orchard Café, Blueberry Country processing sheds, Jary Rd

10.30-11.15 Talk about blueberry operation by Warrick Macdonald (general manager, Blueberry Country)

11.15-11.45 Drive to Pirongia village

11.45-12.30 Lunch at bakery in Pirongia

12.30-12.45 Drive to Grey Rd and visitor centre, Mt Pirongia

12.45-1.30 Mangakara forest walk, Mt Pirongia

1.30-2.15 Drive to Hamilton Gardens, Cobham Drive

2.15-3.00 Te Parapara garden (human-modified soils) (Tamahere soils), Hamilton Gardens with talk by Wiremu Puke

3.00-4.30 Free time, Hamilton Gardens

4.30-5.00 Return to Bryant Hall, Knighton Rd, Hamilton

Introduction to volcanism and its products

Volcanoes are not only conical mountains, but span a very wide range of landforms. A volcano is any landform that results from molten rock (magma) being erupted at the Earth's surface. The size and shape of a volcano reflect how often it erupts, the sizes and types of eruptions, and the composition of the magma it produces. Magmas contain almost all of Earth's known chemical elements but typically they comprise only nine – silicon, oxygen, aluminium, magnesium, iron, calcium, sodium, potassium and titanium – that sum 98% or more by weight. Oxygen and Si together are generally the most abundant elements, making up 48–76 % by weight of most magmas. The chemistry of magma, especially Si content, is important for influencing the way it erupts, and three main magma types, and resulting volcanic rocks, are identified on the basis of their chemical composition. A fourth type, dacite, is usually grouped with andesites but is intermediate between andesite and rhyolite in composition (Smith et al., 2007).

- *Basalt* is rich in Fe and Mg and low in Si and erupts at very high temperatures (~1100–1200 °C) as a very fluid magma. Basalt magma with very little gas cools to form dark black, dense lava, but where magma erupts with lots of gas it cools to form ragged scoria or ash.

- *Rhyolite* magma is rich in Si, K and Na and erupts at temperatures between 700–850 °C as an extremely viscous magma. Rhyolite magma containing lots of gas bubbles cools to form pumice, but if the magma contains little gas it may form obsidian glass.

- *Andesite* (also *dacite*) magma is intermediate in composition and physical properties. Erupting at ~800–1000 °C it is more viscous than basalt, but much less viscous than rhyolite. Andesite magma cools to form dark grey lava if gas-poor or scoria or ash if gas-rich.

Volcanic eruptions vary remarkably in style and size. The smallest may just dust the summit of a volcano with volcanic ash, whereas the largest are capable of creating entirely new landscapes across whole countries. Although there is great diversity of volcanic behaviour, eruptions can generally be classed as one of two broad types – either explosive or effusive.

Explosive eruptions

Explosive eruptions are caused by gases violently ripping apart and shattering magma into pieces which cool and solidify into pumice (if the magma is rhyolite), scoria (if the magma is andesite or basalt), and volcanic ash (sand and dust-sized fragments mainly of volcanic glass). Explosive eruptions take place in one of two ways. The first involves gas that is contained within the magma at depth. As magma rises to the surface the drop in pressure causes these gases to bubble and expand violently. Close to the surface magma becomes like the foam that explodes out of champagne or a can of fizzy drink when first opened. The second type of explosive eruptions occurs when magma contacts water, either beneath the ground (at an aquifer) or at the land surface (either a lake or the sea). The very high temperature of magma (700–1200 °C) means that it instantly boils the water, causing violent steam explosions, which blow apart the magma, eject liquid water and steam, and rip up the ground containing the water. These types of eruptions are called phreatomagmatic eruptions.

In both types of explosive eruptions, the shattered magma and other rock fragments are carried violently into the air by the uprush of expanding gases, before settling back to the ground to form a pyroclastic (literally 'fiery fragmental') deposit. The more explosive the eruption the more energetic this roaring stream of gas, and the higher the tephra and rock fragments will be carried into the air. The higher the material goes, the further from the volcano it will be blown by the wind, and so there is a close correlation or relationship between the energy of an eruption and how far the eruption products are spread (Smith et al., 2007).

Effusive eruptions

Effusive eruptions occur when magma contains little gas and so cannot behave explosively, but instead quietly extrudes at the surface forming flows of lava. The very fluid basalt lavas can flow long distances forming extensive, almost flat volcanoes, whereas very viscous rhyolite lava piles up around the vent, like toothpaste squeezed from a tube, forming large, steep-sided mounds called domes.

Three different types of volcanoes in North Island

Although North Island's active volcanoes look quite different from one another, all can be grouped into one of three main landform types:

- classic cones or stratovolcanoes (e.g. Mt Taranaki, Mt Ruapehu)
- volcanic fields (e.g. Auckland Volcanic Field, South Auckland Volcanic Field); and
- caldera collapse craters (e.g. Rotorua caldera, Taupo caldera, Haroharo caldera)

Each of these has obvious landforms and the violence and styles of eruptions are unique to each. These differences reflect the type of magma erupted: basalt at the volcanic fields, andesite at the cone volcanoes, and rhyolite at the calderas. During our trip, we will see mainly rhyolitic or dacitic deposits and landforms

Caldera volcanoes and eruptions

A caldera is a large, deep crater resulting from the caving in of a rhyolite magma chamber. In places the caldera walls can be seen as steep cliffs, but many are difficult to observe in the landscape because they may be filled in with erupted material or covered by water. Calderas have extremely violent origins and form when a vast amount of viscous rhyolite magma, bubbling with gas, erupts explosively from a magma chamber that may only be a few kilometres beneath the ground. During these eruptions, so much magma is erupted that the chamber empties, leaving the ground above it unsupported. This area collapses, dropping like a piston, to form a wide, deep depression. Lake Taupo, the area to the east of Rotorua known as Okataina, and Mayor Island or Tuhua, are three examples of recently active caldera volcanoes in New Zealand. Some older calderas exist in the area between Taupo and Rotorua and on the Coromandel Peninsula (see Fig. 2 below).

Rhyolite calderas may be active for several hundred thousand years, but large eruptions are rare, with typically thousands of years between events. Caldera collapse is not the only effect on the landscape arising from these large explosive eruptions. Huge quantities of pumice, ash and gas are pumped into the atmosphere, and through a combination of heat and momentum, a roiling column of this material may rise to over 50 km above the caldera. From this height, ash and especially aerosols – gases and tiny drops of acid – can spread around the globe, affecting the world's climate for several years. Closer to the caldera the landscape may be buried by metres of pumice. The most devastating process, however, occurs when this column of material falls back to earth like a fountain, then surges out in all directions from the caldera as a hurricane-like billowing, ground-hugging flow of hot pumice, ash and gas. These pyroclastic flows or 'density currents' can travel over 100 km at the speed of a racing car, leaving behind a layer of volcanic (pyroclastic) debris that might be more than 100 metres deep. Some flows are so hot (600-700 °C) and thick that the ash and pumice fragments weld back together, forming solid rock known as partially or densely welded ignimbrite (Smith et al., 2007).

Dome building

Eruptions from rhyolite volcanoes are not always so explosively catastrophic. A small amount of rhyolite magma may remain after a caldera eruption, which is exhausted of all gas and so can only ooze from the volcano slowly, often along the faults and fissures opened up by earlier caldera collapse. The very high viscosity means the silica-rich rhyolite lava will not flow far, and instead, piles up around the vent, like toothpaste squeezed from a tube, to form a steep-sided dome or dome complexes. These domes are prominent landscape features amongst the caldera volcanoes of the Taupo Volcanic Zone (TVZ) (Fig. 1). Mokoia Island and Mt Ngongotaha are rhyolite lava domes erupted within Rotorua caldera, and Mt Tarawera comprises a collection of lava domes erupted c. AD 1314 within Haroharo/Okataina caldera.

Defining tephra, ash, and lapilli

'Tephra' comes from a Greek word *tephra* meaning ashes, and is an all-encompassing term for the explosively erupted, loose, pyroclastic (fragmental) products of volcanic eruptions (Lowe, 2011). It includes all grain sizes ranging from the finest dust to blocks the size of cars. 'Ash' is not a burnt residue. Rather, it consists of particles <2 mm in diameter including rock particles (lithics), pumice, mineral grains (crystals), and glass shards. Grains 2–64 mm in size are called lapilli (lapillus for single grain), and particles >64 mm are called blocks if they are sharp and angular in shape, or bombs if they are partly rounded or smooth in shape.



Table 1 Tephra-related nomenclature in brief (from Lowe, 2011).

Term	Definition
Tephra	All the explosively-erupted, unconsolidated pyroclastic products of a volcanic eruption (Greek <i>tephra</i> , 'ashes')
Cryptotephra	Tephra-derived glass-shard or crystal concentration, or both, preserved in sediment (including ice) or soil and not visible as a layer to the naked eye (Greek <i>kryptein</i> , 'to hide')
Tephrostratigraphy	Study of sequences of tephra layers and associated deposits, their distribution and stratigraphic relationships, and their relative and numerical ages. Involves defining, describing, characterizing, and dating tephra layers in the field and laboratory
Tephrochronology <i>(sensu stricto)</i>	Use of tephra layers as isochrons (time-parallel marker beds) to connect and synchronize sequences and to transfer relative or numerical ages to them using stratigraphy and other tools. An age-equivalent dating method
Tephrochronology <i>(sensu lato)</i>	All aspects of tephra studies and their application
Tephrochronometry	Obtaining a numerical age or date for a tephra layer

Quaternary volcanism in Taupo Volcanic Zone

The highest concentration of Quaternary volcanic activity in New Zealand occurs in the TVZ. This narrow band of cone and caldera volcanoes extends from Ruapehu in the south, over 240 kilometres to Whakaari (White Island) in the north, with the Taupo, Okataina, Rotorua and other calderas nestled between. Some of the planet's largest and most violent volcanic eruptions have occurred from this zone, as well as New Zealand's most recent small eruptions (Ruapehu 1995-1996, Whakaari 2000, Te Maari craters on Tongariro 2012). Volcanism occurs in the TVZ, and at Taranaki, because of subduction of the Pacific tectonic plate beneath the North Island. As this plate descends and is heated, water and other fluids are boiled off and stream into the mantle rocks under the North Island. These fluids cause chemical changes that enable the otherwise solid rock of the mantle to melt, forming basaltic magma. This magma rises until, because of its higher density, it gets trapped underneath the continental crust of the North Island. Here the very hot basalt magma acts like a gigantic blow-torch, melting the crust and mixing with it to form andesite magma, which is then erupted as cone volcanoes. Where enough melting of the continental crust occurs, rhyolite magma forms, generating caldera volcanoes (Smith et al., 2007).




Fig. 1 Simplified North Island plate tectonic setting and main volcanic centres and caldera volcanoes active in the Quaternary (based on Leonard et al., 2010). North Island lies entirely on the Australian Plate. Diagram at right courtesy of Adrian Pittari.




Fig. 2 Volcanic centres and the ages of activity and rocks (including welded ignimbrites) of central North Island. Diagram courtesy of Roger Briggs (after Briggs et al., 2005).

TVZ

The TVZ comprises three distinct parts (Fig. 1). A southern part, dominated by andesite cones, includes the active Ruapehu and Tongariro volcanoes and the probably extinct Pihanga and Tihia-Kakaramea cones. A northern part, which is also dominated by andesite stratovolcanoes, includes the active Whakaari (White Is.) and mid-Holocene Putauaki (Edgecumbe), and the much older, deeply eroded cones of Motuhora (Whale Is.) and Manawahe. These two stratovolcano clusters book-end the third and largest central part of the zone, extending from Turangi north to the Rotorua lakes district. This area is dominated by rhyolite calderas, including the highly active Taupo and Okataina volcanoes (Figs. 1-2), and older calderas including Mangakino, Kapenga, Whakamaru, Reporoa, Rotorua, and newly-identified Ohakuri caldera (Gravely et al., 2007). The origins and extent of Rotorua caldera are debated. Large explosive eruptions over the last 2 million years or so from this nested collection of rhyolite volcanoes have produced a huge volume of pyroclastic deposits, which when loose are called tephra (Table 1), and many of the older volcanoes cannot be seen in the landscape because of burial underneath hundreds of metres of volcanic material from more recent eruptions. The products of these caldera eruptions are most obvious as the extensive plateaux flanking the western and eastern sides of the TVZ, which erosion reveals to be made up of many layers or sheets of ignimbrite, pumice, and tephra fallout layers. However, caldera eruption products are found far beyond the more obviously volcanic landscape of the central North Island. If we consider a volcano as including all the material erupted from it, then in a sense the entire area from Auckland to Hawkes Bay is part of a huge caldera volcano centred on TVZ.

Introduction to Taupo Volcanic Centre and the Taupo eruption

Taupo volcano (Fig. 2) is the most frequently active and productive rhyolite volcano on Earth. Activity began after the eruption of the widespread and voluminous c. 340 ka Whakamaru group ignimbrites, including widespread Rangitawa tephra from Whakamaru caldera. Modern activity began c. 50 ka with nearly 40 eruptions recognised (Fig. 3). These were overwhelmingly pyroclastic (>95%) and from vents mostly now concealed beneath Lake Taupo (Wilson et al., 2006). Pyroclastic deposits exposed in the Taupo-Maroa area represent 11 eruptions from c. 50 ka to c. 25 cal ka, and then the phreatomagmatic Kawakawa/Oruanui eruption occurred at c. 25 cal ka. This super-eruption was extremely voluminous with a total dense-rock equivalent (DRE) volume of ~530 km³ (equivalent to nearly 1200 km³ as bulk pyroclastic material) (Wilson et al., 2006), and it generated one of the most widespread and stratigraphically useful tephra-fall deposits in the New Zealand region (Wilson, 2001; Wilson et al., 2006; Lowe et al., 2008, 2013; Holt et al., 2010). Caldera collapse associated with this eruption generated most of the modern outline of the basin now partly filled by Lake Taupo, and much of the central North Island landscape was changed as a consequence of the eruption (Manville, 2002; Manville and Wilson, 2004; Wilson et al., 2009).

Since the Kawakawa eruption (also known as the Oruanui eruption, hence now referred to as Kawakawa/Oruanui event), there have been a further 28 eruptions, all but three taking place in the last 12,000 cal years. The most recent (apart from lava extrusion of eruption Z – regarded in some papers as part of the Taupo eruption sequence: Houghton et al., 2015) was the extremely powerful ‘ultraplinian’ caldera-forming Taupo eruption event (Wilson and Walker, 1985; Wilson, 1993) (Table 2). Wilson (1993) erected a volcanological nomenclature for these 28 events beginning with ψ (the oldest) followed by Ω , A, ..., and Z. Other names are also used for some of them (Froggatt and Lowe, 1990).




Fig. 3. Interfingering stratigraphic relationships, ages, and volumes (as non-vesiculated, void-free magma or dense-rock equivalent, DRE; multiply by ~ 3 to obtain approximate bulk volumes) of tephras erupted from Okataina and Taupo caldera volcanoes in North Island, New Zealand, since c. 50 cal ka (based on Wilson et al., 2009; Leonard et al., 2010; Danišik et al., 2012). Eruptions from Maroa and Mayor Island volcanic centres are also recorded (Lowe et al., 2015). Earthquake Flat tephra (EFT) (7 km^3 DRE) was erupted from the Kapenga caldera volcano immediately after the Rotoiti/Rotoehu eruption. Ages on around 20 other widespread tephras erupted since 30,000 cal yr BP were revised by Lowe et al. (2013) using a Bayesian age modelling approach.

Table 2 Summary of tephra names and ages and other information for 28 Taupo eruptives post-dating the Kawakawa/Oruanui eruption c. 25 cal ka (after Wilson, 1993, 1994). Note that ages for some eruptives have been modified since this table was published (Lowe et al., 2013)

Previous published tephra formation name	Volcanological name	Adopted age (years BP) ¹⁴ C time scale	Adopted age (years BP) calibrated timescale	Bulk volume in km ³ as used by Wilson (1993)	Other volume estimates, in km ³	Eruptive activity		Pyroclastic flow(s)	Lava extrusion
						'dry' fall	'wet' fall		
(not defined)	Eruption Z	-	1740	0.28	-	-	-	-	yes
Taupo Tephra	Unit Y	1850	1770	44.75	65, 105	yes	yes	yes	-
Mapara Tephra	Unit X	2150	2150	0.8	0.65, 2, 6	yes	yes	-	(possible)
(not recorded)	Unit W	(2650)	2750	0.023	-	-	yes	-	yes
Whakaipo Tephra	Unit V	2700	2800	0.8	0.8, 1.5, 2, 6	yes	yes	-	(possible)
(not recorded)	Unit U	(2750)	2850	0.2	-	-	yes	-	-
(not recorded)	Unit T	(3000)	3200	0.08	-	-	yes	-	(possible)
Waimihia Tephra	Unit S	3300	3550	16.9	15, 17, 19, 29	yes	-	yes	-
Hinemaiaia Tephra	Unit R	3950	4450	0.05	-	-	yes	-	-
	Unit Q	(4050)	4550	0.15	-	-	-	yes	(possible)
	Unit P	(4100)	4750	0.05	-	-	yes	yes	(possible)
	Unit O	(4150)	4800	0.05	-	-	yes	yes	(possible)
	Unit N	4200	4850	0.15	(3)	-	yes	yes	(possible)
	Unit M	(4500)	5250	0.2	-	-	yes	(possible)	(possible)
	Unit L	4550	5300	0.07	-	-	-	yes	yes
	Unit K	(4600)	5350	0.35	-	-	yes	yes	(possible)
	Unit J	(4620)	5370	0.015	-	-	-	yes	(possible)
Motutere Tephra	Unit I	(5200)	5950	0.02	-	-	-	yes	(possible)
	Unit H	(5300)	6050	0.2	(0.5, 1)	-	yes	yes	(possible)
(not recorded)	Unit G	(5800)	6650	0.5	-	-	yes	yes	(possible)
(not recorded)	Unit F	(6150)	7050	0.12	-	-	-	yes	yes
Opepe Tephra	Unit E	(9050)	9950	4.8	4, 5, 12	yes	yes	yes	-
Poronui Tephra	Unit D	(9780)	11 380	0.2	(3, 3.5, 7)	-	yes	-	yes
(not recorded)	Unit C	(9800)	11 400	0.75	-	-	yes	yes	(possible)
Karapiti Tephra	Unit B	10 100	11 800	1.4	2, 5, 6	yes	yes	-	-
(not recorded)	Unit A	(c. 14 200)	(c. 17 000)	(0.01)	-	-	-	yes	(possible)
(not recorded)	Unit Ω	(c. 15 600)	(c. 18 800)	0.1	-	-	yes	yes	-
(not recorded)	Unit Ψ	(c. 17 200)	(c. 20 500)	(0.05)	-	-	yes	yes	(possible)
Kawakawa Tephra	Oruanui ignimbrite	22 600	24 000-	300	-	-	-	yes	-
	Oruanui fall deposit		26 500	c. 500	-	-	yes	-	-

Taupo eruption

The so-called Taupo eruption (eruption Y) took place in late summer to early autumn (typically late March to early April) on the basis of fruit and seeds preserved in a buried forest at Pureora (Fig. 4) and the lack of an outer latewood ring (Clarkson et al., 1988; Palmer et al., 1988; see Lowe and King, 2015). The eruption year was AD 232 ± 10 AD based on dendrochronology and wiggle-match dating by Hogg et al. (2012, 2019). A total eruptive bulk volume was estimated at ~105 km³ (~30 km³ DRE).




Fig. 4 Distribution of Taupo ignimbrite radially around Lake Taupo and tephra fallout isopachs (in cm) derived from the Taupo eruption (after Hogg et al., 2012; Houghton et al., 2014; Wilson and Leonard, 2015).




Fig. 5 Summary of the stratigraphy of pyroclastic deposits of the Taupo eruption (eruption Y of Wilson, 1993). Graphs at right indicate qualitative changes in the inferred degree of magma-water interaction and the magma discharge rate (which becomes ‘extreme’ during emplacement of Subunit Y6, Taupo ignimbrite) during the eruption. E = erosion horizons formed by running water; S = erosion horizons from shearing beneath the fast-moving pyroclastic flow that deposited the Taupo ignimbrite (after Houghton and Wilson, 1986; Houghton et al., 2015) (note: subunit Y7 = eruption Z in Table 2).

The eruption was complex, generating three phreatomagmatic (‘wet’) and two plinian (‘dry’) fall units, a multi-flow intraplinian ignimbrite, and at the eruption climax, the extremely violently emplaced Taupo ignimbrite (Subunit Y6) (Fig. 5). The duration of the entire eruption episode was of the order of tens of hours with a break after Y3 (probably < 3 weeks). The height of the main ultra-plinian phase (Subunit Y5) eruption column has been estimated at 50-55 km although Houghton et al. (2014) suggested that the eruption be ‘downgraded’ to a plinian event with an eruption column height ~35-40 km; this phase lasted for ~6 to 17 hours (Walker, 1980). The ignimbritic material was emplaced cataclysmically over about 400 seconds (~7 mins) by an extremely energetic pyroclastic flow (also called a pyroclastic density current or PDC) moving at 200-300 m/s over a near-circular area (~80 km radius) of c. 20,000 km² around the vents (Wilson, 1985, 1993; Wilson and Walker, 1985; Smith and Houghton, 1995). Its temperature was about 400-500° C at ~50 km from vent (within ~30-40 km of the vent it was 150-300° C) (McClelland et al., 2004; Hudspith et al., 2009). The non-welded ignimbrite was spread thinly over the landscape to generate an archetypal ‘low aspect ratio’ ignimbrite (Walker and Wilson, 1985). Ignimbrite-veener deposits (IVD) are generally 0.5 to 1 m-thick and drape topographic highs; valley-pond ignimbrites (VPI) infill valleys and depressions to depths of 5 to 60 m (Manville, 2001a). Because of its extreme violence and energy release ($\geq 150 \pm 50$ megaton TNT explosive yield, cf. Hiroshima bomb 0.015 Mt), and by analogy with the 1883 Krakatau event, it is likely that the ignimbrite-emplacement phase generated a volcano-meteorological tsunami that may have reached coastal areas worldwide (Lowe and de Lange, 2000). The emplacement of the ignimbrite destroyed all forests in its path (about 1 km³ of timber), and then ignimbrite contains numerous charred logs and charcoal, many of the logs notably being orientated radially around the vents (Froggatt et al., 1981). Yet the forests recovered within 100-200 years (Clarkson et al., 1992, 1995; Wilmschurst and McGlone, 1996; Lowe and King, 2015).

The wide variation in eruption styles and dynamics relate to variations in discharge rate and the degree of interaction between the magma and water in the proto-Lake Taupo (Wilson and Walker, 1985; Wilson, 1993, 1994; Houghton et al., 2010). Much of Lake Taupo was expelled, evaporated, or drained into a caldera-collapse

structure beneath the current lake floor during the eruption. Afterwards it refilled over approximately 15 to 40 years, reaching a height of ~400 m, about 30-40 m above its present level (357 m) to form a semi-continuous, wave-cut bench and highstand shoreline deposits (Manville et al., 1999, 2007, 2009). Catastrophic failure of a pumiceous pyroclastic dam led to the reestablishment of the Waikato River and the release of ~20 km³ of water in a single phase, the peak discharge being 20,000–40,000 m³/sec, equivalent to the Mississippi River in flood (Manville et al., 1999). The break-out flood deposits can be traced 220 km downstream of Lake Taupo (Manville et al., 2007) including on the low terrace that supports the Hamilton Gardens.

Introduction to Okataina Volcanic Centre and Kaharoa and Tarawera eruptions

The Okataina Volcanic Centre (Figs. 1-2) comprises two main parts, the northern Haroharo Volcanic Complex (and Okareka embayment), and the southern Tarawera Volcanic Complex (centred on Mt Tarawera). Since c. 50 cal ka there have been at least 25 eruptions from Okataina (Fig. 4) and the most important tephras since c. 25,000 cal years ago are listed in Table 3 (below).

Kaharoa eruption

Most of us are aware of the basaltic Tarawera eruption on 10th June 1886: the high toll on life (~120 people), landscape devastation, and loss of the Pink and White Terraces. But this was not the first time that Mt Tarawera produced an eruption of importance both to volcanology and human history. The Kaharoa eruption occurred at Mt Tarawera in the winter of 1314 AD (± 12 years) (Hogg et al., 2003; Sahetapy-Engel et al., 2014). The importance of the Kaharoa eruption is at least threefold (Lowe and Pittari, 2014). (1) It is the most recent rhyolite eruption in New Zealand, and the largest New Zealand eruption volumetrically of the last millennium. (2) The Kaharoa tephra is an important marker horizon in late Holocene stratigraphy and geoarchaeology (Vucetich and Pullar, 1964; Lowe et al. 1998, 2000), and in particular helps to constrain the timing of settlement of early Polynesians in North Island, which occurred in c. AD 1280, a few decades before the eruption (Newnham et al. 1998; Wilmschurst et al., 2008). (3) There is a link between the soils that developed on the Kaharoa tephra, the animal ‘wasting’ disease known as ‘bush sickness’, and the birth of the government soil survey group as an independent organisation (Tonkin, 2012).




Fig. 6. Isopach map of Kaharoa tephra fallout blown firstly southeastward then northwestward to generate the lobate fallout pattern. The tephra isochron has been extended using cryptotephra occurrences in peats and lake sediments the Waikato and Auckland regions (after Lowe et al., 1998).

Table 3 Summary of main rhyolitic tephras deposited in the Rotorua-Galatea region since c. 25,400 cal yr BP.

Name (source)*	Date or age [†]	Description
Tarawera Tephra (Tr) (Tarawera)	10 June 1886	Comprises basaltic scoria (Tarawera Scoria) with occasional rhyolite clasts and/or fine greyish brown 'muddy' ash (Rotomahana Mud). Mud was dispersed more widely than the scoria.
Kaharoa Tephra (Ka) (Tarawera)	1314 ± 12 AD (636 ± 12 cal yr BP)	Fine to coarse white to grey ash, with occasional dense pumice, rhyolite, obsidian and basalt lapilli. Contains abundant biotite.
Taupo Tephra (also known as Unit Y) (Tp) (Taupo)	232 ± 10 AD (1718 ± 10 cal yr BP)	Creamy coloured coarse ash with plentiful shower-bedded pumice lapilli (crushable). Ignimbrite unit always associated with charcoal fragments.
Whakatane Tephra (Wk) (Haroharo)	5526 ± 145 cal yr BP	Shower-bedded pale yellow coarse ash, overlying a fine to coarse rhyolitic (pale grey) ash. Rich in cummingtonite. Reddish-brown uppermost horizon (sometimes with basaltic Rotokawau tephra c. 4 cal ka).
Mamaku Tephra (Ma) (Haroharo)	7940 ± 257 cal yr BP	Loose, coarse yellowish-brown pumice ash grading into a weakly shower-bedded coarse ash/lapilli.
Rotoma Tephra (Rm) (Haroharo)	9423 ± 120 cal yr BP	Shower-bedded fine grey to yellowish brown ash with coarse ash layers, cummingtonite. Marked by a dark Ah horizon at top, sometimes with charcoal, or podzolised.
Waiohau Tephra (Wh) (Tarawera)	14,009 ± 155 cal yr BP	Grey fine and coarse shower-bedded ash. Distinctive v. fine cream ash layer at the base. Usually has well developed yellowish-brown or greyish upper soil horizon. Deposited a few centuries before late-glacial cool episode (NZce-3) in NZ-CES [‡] .
Rotorua Tephra (Rr) (Okareka embayment)	15,635 ± 412 cal yr BP	Shower-bedded pumiceous yellowish lapilli or blocks (gravel). Occasional rhyolitic lithics. Deposited at start of late-glacial mild episode (NZce-4) in NZ-CES [‡] .
Rerewhakaaitu Tephra (Rk) (Tarawera)	17,496 ± 462 cal yr BP	Yellowish-brown ash grading down into tephric loess. Contains abundant biotite. Marks transition from Last Glacial to post-glacial conditions (Termination I); reforestation occurred soon after deposition.
Okareka Tephra (Ok) (Tarawera)	21,858 ± 290 cal yr BP	Yellowish brown ash contains abundant biotite. Typically encased in yellowish to olive brown tephric loess. Deposited just before stadial A (NZce-6) NZ-CES [‡] .
Te Rere Tephra (Te) (Haroharo/Okareka)	25,171 ± 964 cal yr BP	Yellowish-brown ash (typically encased in yellowish to olive brown tephric loess).
Kawakawa Tephra (Kk) (also known as Oruanui) (Taupo)	25,358 ± 162 cal yr BP	Olive brown to pale yellowish brown ash (typically encased in yellowish to olive brown tephric loess). Deposited just before interstadial D (NZce-9) in NZ-CES [‡] .

*Terminology is based mainly on Froggatt and Lowe (1990). Descriptions generalised because character may differ from proximal to distal locations and from site to site. The region has received additional distal tephras from Taupo and Tuhua volcanic centres, and has been dusted regularly with andesitic tephra fallout from numerous eruptions at Tongariro Volcanic Centre and Egmont/Taranaki Volcano, most recently in the 1995-96 Ruapehu eruptions.

[†]Ages are given in calibrated or calendar (cal) years (95% probability range) before present (BP). Ages are based on Bayesian probability age modelling: Lowe et al. (2008, 2013). Calendar dates for the Kaharoa and Taupo eruptions have been determined by dendrochronology and ¹⁴C wiggle-match dating (Hogg et al., 2003, 2012, 2019).

[‡]NZ climate event stratigraphy of Barrell et al. (2013)

Tarawera eruption

The Tarawera eruption of 10 June, 1886, was the biggest and most destructive eruption in New Zealand during the historical (European) period. It was a basaltic rather than rhyolitic event, but was nevertheless very explosive: the resulting scoria fall ('Tarawera Scoria') has a dispersal similar in extent to that of the Vesuvius 79 AD pumice fall and is one of the few known examples of a basaltic deposit of plinian type from a fissure source (Walker et al., 1984). The eruption cored out a series of craters in a 7-km-long fissure through the antecedent rhyolite domes (including those emplaced during the Kaharoa event) of Mt Tarawera, and then generated more craters along an 8-km-long southwest extension of the fissure across the Rotomahana basin (which contained two shallow lakes and large silica sinter aprons, the 'Pink' and the 'White' terraces, associated with extensive hydrothermal activity) to Waimangu. Narratives (summarized authoritatively by Keam, 1988) indicate that after a series of precursory earthquakes from ~12.30 am, the eruption began at Ruawahia Dome at about 2.00 am on 10 June, 1886, and then gradually extended both northeastward and southwestward. At ~2.10 am the eruption intensified with the ascent of a tephra plume from the vicinity of Ruawahia Dome up to ~9.5 km. By 2.30 am craters along the whole length of the fissure were erupting, with the Rotomahana extension beginning to erupt possibly at ~3.20 am. By 3.30 am, craters along the entire 17 km-length of the fissure from Wahanga to Waimangu were in eruption. This paroxysmal stage of the eruption was over by 6.00 am when most activity ceased. See also recent papers by de Ronde et al. (2016a, 2016b), Keam (2016), and Lorrey and Woolley (2018).




Fig. 7. Isopach map of 1886 Tarawera scoria fallout (in cm). x = location where scoria occurs mixed with Rotomahana Mud, but does not form a discrete layer (from Walker et al., 1984).

The erupted products were exclusively pyroclastic (no lava flows were generated, although basalt dikes were emplaced). The total volume (as deposited) of Tarawera Scoria is ~2 km³ (Walker et al., 1984). The eruption along the Rotomahana and Waimangu extension was mainly phreatomagmatic (interaction between basalt magma and hydrothermal water) and phreatic. The explosive expansion of superheated water fragmented the country rock containing the hydrothermal system, plus subordinate lake sediment, to produce surge beds and fall deposits ('Rotomahana Mud') that rained out over much of the Bay of Plenty and beyond (~0.5 km³ as deposited). Near Rotomahana, the surge beds were emplaced violently by hot and fast-moving turbulent pyroclastic surges or density currents up to ~6 km from source (Nairn, 1979) (Fig. 8).

Lightning during the eruption set fire to a house in Te Wairoa and to the forest on the north shore of Lake Tarawera; strong winds flattened many trees at Lake Tikitapu; and suffocating gases and falling mud and ash made breathing difficult at Te Wairoa, where most buildings were buried or collapsed under the weight of ~1 m of mudfall. A notable exception was *Hinemihi*, a large meeting house where most survivors were sheltered, because wooden forms for seating guests during Maori concerts were used to prop up the roof.

All but seven of the 108 known fatalities arising from the Tarawera eruption were Maori (the true number of deaths may have been ~120, but the oft-cited ~150 is erroneous: Lowe et al., 2001). The majority of deaths were the result of the Rotomahana explosions, especially the lethal, scorching pyroclastic surges and blasts. Clearly the event had a profound impact on Maori (and others) in the Te Wairoa and Rotomahana area especially, but trauma was felt throughout the extensive fallout zone in the Bay of Plenty and eastern North Island (Keam, 1988). For example, some groups of Maori in the region of the Rangitaiki and Tarawera rivers, north of Tarawera, became refugees at Matata. Although they had escaped with their lives and without serious injury, their possessions were buried by ~15–30 cm of tephra (some were retrievable by excavation), many potato pits were lost and those with livestock had no feed for them and so many starved (Keam 1988; Lowe et al., 2002). These people were eventually resettled in 1903–1905. The plight of these and other Maori seem minor in comparison with the difficulties of those from Te Wairoa-Rotomahana: apart from the lives lost, all possessions had been buried and many crushed. Among livestock, most smaller animals were killed, but dogs, pigs, cattle and horses that survived wandered loose and starving. The main livelihood of the region, tourism, had been destroyed, literally overnight. Whilst Maori continued to participate in the tourist trade, its control effectively moved into European hands from 1894 with the opening of the railway line to Rotorua. However, perhaps the biggest societal impact, according to Keam (1988), was the loss of land. For thirty years, Maori groups in the region had been generally secure in possession of their land and property. In previous times, under the old order, the prospect had always existed that a group might lose homes and land through warfare, but by the time of the Tarawera eruption, the people, long-established traders with European settlers, had become accustomed to a newfound security.

The eruption rather than warfare (against which there could at least be retaliatory or conciliatory action to make-good losses) had destroyed that security and dispossessed the people of the land, which they had prized most. Offers of resettlement for the surviving group, mainly the Tuhourangi subtribe or clan (*hapu*), were received from various parts of central and eastern North Island and beyond, but most settled at Whakarewarewa and Ngapuna, both near Rotorua. Eventually gifts of land were formally ratified and provided a home for most of the Tuhourangi people. Other Tuhourangi settled for a time in the Bay of Plenty and Coromandel. After 30-50 years almost all the refugees or their descendants had returned to Whakarewarewa or Ngapuna and the gifted land was returned to the donors (Keam, 1988; Lowe et al., 2002; see also Cashman et al., 2008).




Fig. 8. Map of Tarawera area showing locations of the main craters of the 10 June 1886 fissure eruption across Tarawera Volcanic Complex, Rotomahana Crater (including pre-eruption lakes Rotomahana and Rotomakariri), and Waimangu craters (after Lowe et al., 2002). Locations of villages and associated fatalities (numbers in parentheses) are based on Keam (1988) (there was an additional death at an unknown locality). Fatalities were all Maori apart from six Europeans at Te Wairoa and one European and three (part) Maori at Waingongongo. On the night of the eruption nearly half of Te Arika's 27 residents were camped at Pink Terrace (Otukupuarangi). Inset shows eastern North Island and documented limits of tephra fallout from the eruption (based on maps by A.P.W. Thomas, 1888). Ash fell on several ships at sea, the farthest being *Julia Pryce* (c. 300 km) and S.S. *Waimea* (c. 1000 km) north of North Island (Keam, 1988).

Distribution of main soil-forming tephras

The thickest tephra sequences occur downwind of the TVZ in the Rotorua-Taupo area, Bay of Plenty, East Coast-Poverty Bay, and Hawke's Bay (Fig. 9). Moderately thick deposits are found in Wanganui-Taranaki, King Country-Waikato-Coromandel, and Auckland regions. Fewer tephra layers occur in other parts of the North Island. Only a handful of tephras has been recognised so far in the South Island. They include Rangitawa Tephra, erupted c. 340,000 years ago from Whakamaru volcano, and Kawakawa Tephra, erupted c. 25,400 cal years ago from Taupo volcano.




Fig. 9 Map showing plate tectonic setting, the main volcanic centres that produced parent materials for many tephra-derived soils, and the general dispersal of tephra on North Island. EG, Egmont/Taranaki volcano; TG, Tongariro Volcanic Centre; TP, Taupo Volcanic Centre; OK, Okataina Volcanic Centre (includes Mt Tarawera and Haroharo volcanic complexes); TU, Tuhua Volcanic Centre (Mayor Is.); W, Whakaari (White Is.) (from Lowe and Palmer, 2005).

Tephra-derived soils of New Zealand

The relatively young landscapes of much of central North Island, especially the Bay of Plenty-Taupo-Tongariro area, are predominantly of volcanic origin. Consequently, the nature and distribution of the soils contained in these landscapes are very strongly influenced by the volcanic history as well as faulting (tectonism) and reworking by water and wind. Variations in the age, thickness, and mineralogical composition of the tephra deposits in which the soils are forming have significantly contributed to the soil patterns. Post-depositional

erosion and reworking of tephra deposits have helped to shape the landscape and, in turn, have created 'genetic' links of varying strengths between the soils and landforms of the region. To some extent, climatic and indigenous vegetation gradients have also been imprinted upon the soils and their distribution pattern (Molloy and Christie, 1998).

Classification of tephra-derived soils

Soils formed from tephra deposits are represented by five orders of the New Zealand Soil Classification (NZSC) which reflect increasing age and development: (Tephric) Recent Soils (~1% of North Island soils), Pumice Soils (~15%), Allophanic Soils (~12%), and Granular Soils (~3%) and (rarely) Ultic Soils (Lowe and Palmer, 2005; Hewitt, 2010; see also table below for relative abundances of New Zealand soils).




Fig. 10 Major taxonomic development paths of New Zealand soils (after A.E. Hewitt in Molloy and Christie, 1998).

Tephric Recent Soils (Entisols)

These soils occur on very young, mainly andesitic or basaltic eruptives (~100–600 years old) close to the volcanic centres of Taranaki, Tongariro-Ruapehu, Rangitoto Island, and Mt Tarawera. They are typically gravelly or sandy and have few nutrients – the result of insufficient time for weathering to release them – and cannot store much water.




Fig. 11 Relative abundances of soil orders in New Zealand (after A.E. Hewitt in Molloy and Christie, 1998). Allophanic and Pumice Soils (Andisols) together cover about 25% of the North Island.

Pumice Soils (mainly Vitrand)

Covering a large swath of the central and eastern North Island, these shallow soils are made up of coarse rhyolitic pumice deposits derived mainly from the Taupo (c. 232 AD) and Kaharoa eruptions (c. 1314 AD). The young and weakly developed soils formed from these deposits (~700–1800 years old) are extremely deficient in many elements essential for animal health including copper, selenium, and (especially) cobalt. The Californian Monterey pine (*Pinus radiata*) grew fast and vigorously on the problematic Pumice Soils, partly by tapping into nutrients and moisture in the paleosols and soil horizons formed in tephras buried beneath them (see notes below about Co deficiency and plantation forestry).

Allophanic Soils (mainly Udands)

These deep, versatile soils are formed typically on accumulating sequences of thin, fine-grained interfingering tephra layers from both rhyolitic and andesitic volcanoes, and occur in the Ohakune-Waiouru area, Taranaki, King Country-Waikato, and western Bay of Plenty-Coromandel. Small patches are found on basaltic scoria cones in Auckland-Northland. Most Allophanic Soils have taken between 10,000 and 25,000 years to form, with some as old as c. 50,000 years, and clearly are irreplaceable, yet they are undervalued by most people. Their name comes from the tiny nanocrystalline clay mineral formed in them, allophane, which dominates their physical and chemical properties because of its positive charge and huge surface area: a single teaspoon (about 5 g) of allophane has the surface area of a rugby field (areas range from 250 m² g⁻¹ to as much as 1125 m² g⁻¹ (Lowe and Palmer, 2005; Neall, 2006; McDaniel et al., 2012; Huang et al., 2016a).

Table 4. Abundances of 12 soil orders of the world (of Soil Taxonomy) vs abundances in New Zealand (rank = relative abundance) (after Lowe et al., 2000a).

Order	World ¹		New Zealand ²		Main NZSC order(s)
	Land area % (ice-free)	Rank	Land area %	Rank	
Alfisols	9.7	4	9.9	4	Pallic Soils
Andisols	0.7	12	12.9	3	Allophanic, Pumice, Recent Soils
Aridisols	12.0	2	0.9	9	Semiarid Soils
Entisols	16.2	1	7.4	5	Recent, Gley, Raw, Anthropic Soils
Gelisols	8.6	5	0 ³	12	—
Histosols	1.2	11	0.9	8	Organic Soils
Inceptisols	9.8	3	47.4	1	Brown, Gley, Pallic, Recent Soils
Mollisols	6.9	8	1.2	7	Melanic Soils
Oxisols	7.5	7	0.2	10	Oxidic Soils
Spodosols	2.6	9	13.1	2	Podzol Soils
Ultisols	8.5	6	4.2	6	Ultic, Granular Soils
Vertisols	2.4	10	0.1	11	Melanic Soils
(Non-soils)	(13.9)		(2.0)		(Raw Soils)

¹After Soil Survey Staff (1999).

²Correlations with NZSC based on Hewitt (1998, p.10-14); land area percentages are approximate and based on 1: 1 000 000 maps published by Landcare Research in 1995.

³Gelisols (on frost-churned materials underlain by permafrost) probably occur in NZ's Ross Dependency, Antarctica




Fig. 12 Distribution of four main groupings of tephra-derived soils in North Island (after Kirkpatrick, 1999, based on Rijkse and Hewitt, 1995; modified from Lowe and Palmer, 2005).




Fig. 13 General subdivision of the main groupings of Andisols in North Island into six zones according to their multilayered soil character (soil stratigraphy), the primary compositions of component tephras, and approximate ages of the 1-m deep soil profiles (from Lowe and Palmer, 2005). Ages on Kaharoa and Taupo tephras from Hogg et al. (2003, 2012). Note: 'Aggregating' should read 'Aggrading'.

Pumice Soils, “bush sickness”, and plantation forestry

Inherently low in Co, these pumiceous parent tephras led to low Co levels in soils and herbage, and ultimately to a deficiency in ruminant animals (sheep and cows) that developed a serious and commonly fatal wasting disease known as “bush sickness” (Lowe and Palmer, 2005; Neall, 2006) (Fig. 14). This term referred to a variety of symptoms exhibited by livestock stemming from their inability to produce vitamin B₁₂ (cobalamine, discovered in 1948), of which Co is an essential component for red blood cell production (Cornforth, 1998; Hendy, 2008).




Fig. 14 'Bush-sick' cow on Taupo soil at Ngaroma, South Waikato (from Grange and Taylor, 1932)

Identified in 1911, bush sickness became an increasingly urgent problem in the late 1920s when many farms on Pumice Soils, settled after WW I, were abandoned as the depression hit. The association of bush sickness with the Pumice Soils was recognized by Les Grange (who began mapping volcanic deposits in 1926) in the early 1930s (Grange and Taylor, 1932; Grange, 1937). Acquired Co deficiency also occurred in some other soils where high rainfall and strong leaching were the cause. Veterinarians were unable to diagnose any diseases, and it was thought that some property of the soil was responsible. Analytical techniques of the day (early 1900s) were too insensitive to show what this might be, but trial and error showed that iron ore ("limonite") from some, but not all, sources provided relief as a stock 'lick' (Hendy, 2008). There was some rivalry in this work. Elsa Kidson (Cawthron Institute) and K.J. McNaught (Department of Agriculture) each developed essentially the same methods for finding trace amounts of cobalt. They published a series of papers on cobalt levels in rocks, soils, and pastures, while pointing out each other's errors (Tonkin, 2012).

Initial research was directed at trying to supplement the animals' diets with iron. In 1934, R.E.R. Grimmett and F.B. Shorland (senior chemists at the Department of Agriculture) found that the iron ore which gave the best results contained significant amounts of cobalt, and went against popular wisdom by dosing animals with cobalt, with spectacular results. Australians Underwood and Filmer (1935) confirmed the association. Grimmett and Shorland then developed cobaltised superphosphate fertiliser, which has been applied to the affected area ever since at a rate of a few grams per hectare, and has resulted in the addition of about 250,000 ha of productive farmland to New Zealand's stock. Possibly, this one discovery has paid for all of the scientific research ever carried out in New Zealand (Hendy, 2008). In addition, Grange's insight resulted in soil survey becoming a separate, independent branch of the Department of Scientific and Industrial Research (DSIR) in 1936 rather than effectively an 'add-on' to the Geological Survey Branch as it had been (Lowe, 1990; Tonkin, 2012). Grange was the first director of Soil Survey (and later a director of Geological Survey); it was renamed Soil Bureau in 1945 until morphing into Landcare Research in July, 1992.




Fig. 15 Map of North Island showing single isopachs of tephras deposited from Taupo and Kaharoa eruptions, soils on which were deficient in Co which resulted in 'bush sickness' (from Lowe and Palmer, 2005). Co deficiency also occurred in soils in some other areas under high rainfall. Cartoon by geochemist Dr Nick Kim.



Fig. 16 Landscape at Litchfield, midway between Putaruru and Tokoroa, underlain by shallow Taupo soils on Taupo tephra. The Co deficiency here was the first to be rectified – on the nearby farm of K.S. Cox.

Partly as a result of these agricultural problems, a forestry industry became established, the mainstay tree being *Pinus radiata* from California (known as Monterey pine). *P. radiata* has been very successful, it matures rapidly in about 25 years on average, and can grow on 'clapped out' soils with little problem. Its roots can easily punch through surficial pumice deposits to reach nutrients and water in buried soil horizons. The soils to the east of SH1 between Putaruru and Tokoroa are now growing their 5th crop (rotation) of *P. radiata*. Palmer et al. (2005) used a P-based nutrient model to establish that *P. radiata* was growing sustainably after two rotations on Podzol Soils (Andic Haplohumods) developed on Taupo tephra in elevated areas on the southern Mamaku Plateau to the east of Tokoroa. Models to predict *Pinus radiata* productivity throughout New Zealand were developed by Watt et al. (2010) and various spatial prediction techniques for developing *Pinus radiata* productivity surfaces across New Zealand were compared by Palmer et al. (2010).

Erodibility

The soils developed in loose Taupo tephra deposits around Lake Taupo are susceptible to severe gully erosion. The erodibility of the soils under three classes of vegetation – pasture grasses, ungrazed grass, and scrub – was assessed by Selby and Hoskins (1973) with the use of a specially designed flume that simulated gully erosion. Erosion was found to be significantly higher beneath scrub than beneath grass vegetation when the same erosive stress was applied in each case. Under natural conditions, however, greater runoff, and therefore greater erosive power, always occurs on pasture grasses than under scrub. In spite of the greater inherent erodibility of scrub, actual erosion is therefore greater on pasture lands (Selby and Hoskins, 1973). Under pasture grasses, plant roots and organic matter bind the soil particles and inhibit erosion. This effect is less under scrub. Gully erosion is attributable largely to the low density and easy entrainment of pumice particles. 'Retirement' of such gullies with tree planting has been successful in stabilising them.

Classification

Andisols are one of 12 soil orders in the international soil classification system *Soil Taxonomy* (Soil Survey Staff, 1999). They are often deep soils commonly with depositional stratification developing mainly from ash, pumice, cinders (scoria), or other explosively erupted, pyroclastic volcanic material (referred to collectively as tephra) and volcanoclastic or reworked materials. Andisols occur much less commonly on lavas. Unlike many other soils, Andisol profiles commonly undergo upbuilding pedogenesis as younger tephra materials are deposited on top of older ones, as described below. The resulting profile character is determined by the interplay between the rate at which tephra is added to the land surface and classical topdown processes that form soil horizons. Therefore, understanding Andisol genesis usually requires a stratigraphic approach combined with an appreciation of buried soil horizons and polygenesis (see notes below) (McDaniel et al., 2012). The NZSC splits Andisols into Allophanic Soils (mainly Udands) and Pumice Soils (mainly Vitrandis). Vitrandis are defined as slightly weathered, coarse-textured, glass-dominated Andisol with <15% 1500 kPa water retention on air-dried samples and <30% kPa water retention on undried samples (Soil Survey Staff, 2014).

Andisols are defined by characteristic andic soil properties that include physical, chemical, and mineralogical properties that are fundamentally different from those of soils of other orders (andic properties are termed allophanic soil materials in NZSC). The coarser fractions of Andisols are often dominated by volcanic glass. This glass weathers relatively quickly to yield a fine colloidal or nanoscale fraction (1–100 nm) dominated by short-range-order materials composed of 'active' Al, Si, Fe, and organic matter, especially humus. Previously described erroneously as 'amorphous', short-range order materials comprise extremely tiny but structured nanominerals, referred to nowadays as nanocrystalline, the main ones being allophane and ferrihydrite. Another colloidal constituent, imogolite, comprises long filamental tubes and therefore has both short- and long-range order (Churchman and Lowe, 2012). The nanominerals, chiefly allophane, ferrihydrite, and also Al- or Fe-humus complexes, are responsible for many of the unique properties exhibited by Andisols (McDaniel et al., 2012). The formation of allophane and other minerals including halloysite are described by Churchman and Lowe (2012). Some properties of buried soil horizons may be altered via diagenesis.

Allophane and its formation

Allophane is a nanocrystalline aluminosilicate comprising tiny, hollow spherules (sometimes called nanoballs) ~ 3.5 to 5.0 nm in diameter and with a chemical composition $(1-2)\text{SiO}_2 \cdot \text{Al}_2\text{O}_3 \cdot (2-3)\text{H}_2\text{O}$ (Abidin et al., 2007; Huang et al., 2016a; Theng, 2019) (Figs. 17-19). It provides many tephra-derived soils including Allophanic Soils and Pumice Soils with many of their unique chemical and physical properties (McDaniel et al., 2012; Yuan and Wada, 2012). With its small size, extreme surface area (up to $\sim 1200 \text{ m}^2 \text{ g}^{-1}$) (Allbrook, 1983, 1985; Parfitt, 2009; Yuan and Theng, 2012; Huang et al., 2016a), and variable surface-charge characteristics that arise via $(\text{OH})\text{Al}(\text{OH}_2)$ groups at wall perforations of its outer gibbsitic octahedral sheet $[\text{Al}(\text{OH})_3]$, allophane has strong affinity for water, metal cations, anions, organic molecules and DNA (Harsh, 2012; Huang et al., 2014, 2016a, 2016b).

The essential conditions for the formation of allophane are the activity of silicic acid in the soil solution, the availability of Al species, and the opportunity for co-precipitation (Fig. 21). These conditions are controlled largely by the leaching regime, the organic cycle, and pH, which, in turn, are potentially influenced by numerous environmental factors including rainfall, drainage, depth of burial, parent tephra composition and accumulation rate, dust accession, type of vegetation and supply of humic substances, and human activities (such as burning vegetative cover), together with thermodynamic and kinetic factors (McDaniel et al., 2012). Availability of Al, derived mainly from the dissolution of glass or feldspars, is assumed to be unlimited in this model, though potentially more is available from andesitic and especially basaltic tephtras than rhyolitic tephtras. In contrast, in pedogenic environments rich in organic matter and with pHs ≤ 5 , humus effectively competes for dissolved Al, leaving little Al available for co-precipitation with Si to form allophane or halloysite. In these environments (such as in parts of Japan), Al-humus complexes are formed instead of allophane (McDaniel et al., 2012).




Fig. 17 Diagram of imogolite nanotubes and Al-rich allophane nanospheres, which have similar structures at the atomic scale (from McDaniel et al., 2012).




Fig. 18 Micrographs of (A) allophane (showing hollow spherules) and (B) imogolite (external diameter of nanotubes is ~2 nm) (from McDaniel et al., 2012, after Parfitt, 1990).




Fig. 19 Various volcanic glass compositions and dissolution of Al and Si and their reprecipitation to form allophane spherules or 'nanoballs' (from McDaniel et al., 2012, after Hiradate and Wada, 2005).

In New Zealand, both mineralogical and soil-solution studies on soils derived from tephras extending across a rainfall gradient showed that rainfall, coupled with through-profile drainage, helps govern Si concentration [Si] in soil solution and thus the likelihood of allophane being formed or not (Parfitt et al., 1983; Singleton et al., 1989; Parfitt, 1990, 2009; Churchman and Lowe, 2012). The Si leaching model is summarized as follows: where

[Si] is less than ~10 ppm (mg l^{-1}), allophane is formed; where [Si] is greater than ~10 ppm, halloysite is formed. If [Si] is close to ~10 ppm then either allophane or halloysite may predominate. A profile throughflow threshold of approximately 250 mm per year of drainage water likely controls [Si] – less than ~250 mm means that the loss of Si is insufficient for Al-rich allophane to form and halloysite (or Si-rich allophane) forms instead (Parfitt et al., 1984; Lowe, 1986; McDaniel et al., 2012).




Fig. 20 Simplified allophane-halloysite rainfall Si-leaching model (from McDaniel et al., 2012).




Fig. 21 Environmental influences and controls that govern the critical conditions leading to the formation of different clays from the weathering of tephra, and the likely occurrence, or not, of Andisols (mainly Allophanic and Pumice Soils in NZSC) as a result (from Churchman and Lowe, 2012). ± indicates that the clay mentioned may also be present.

Why Andisols (Allophanic Soils) are rich in carbon

Studies by Huang et al. (2016a) have determined, using an experimental approach with salmon-sperm DNA and artificial allophane, why Andisols may contain large amounts of organic matter (up to ~8–12% organic carbon). The formation of very stable allophane nanoaggregates and microaggregates (see Fig. 40) enables up to 28 $\mu\text{g mg}^{-1}$ of DNA to be adsorbed (~80% of total) within tiny spaces (nanopores) between allophane spherules and allophane nanoaggregates (as “physical adsorption”), giving a total of 34 $\mu\text{g mg}^{-1}$ of DNA adsorbed by the allophane. The stability of the allophane–DNA nano- and microaggregates likely prevents encapsulated DNA from exposure to oxidants, and DNA within small pores between allophane spherules and nanoaggregates may not be accessible to enzymes or microbes, hence enabling DNA protection and preservation in such materials. By implication, substantial organic carbon is therefore likely to be sequestered and protected in allophanic soils (Andisols) in the same way as demonstrated for DNA, that is, predominantly by encapsulation within a tortuous “nanolabyrinthic” network of nanopores and submicropores amidst stable nanoaggregates and microaggregates, rather than by chemisorption alone.

Forming a soil whilst tephras accumulate: geological vs pedological processes

A distinctive feature of many tephra-derived soils is the multilayered nature of their profiles which attests to building up the landscape via the deposition of tephras from numerous eruption. After establishing the stratigraphy (geological layering) of the sequences, the interplay of topdown pedogenesis (soil formation) and the accumulation to tephras on the land surface can be considered (upbuilding pedogenesis).

Topdown pedogenesis is the ‘classical’ formation of *soil horizons* in a profile through various processes that gradually deepen the profile as a downward moving ‘front’ on a pre-existing parent material on a stable land surface with *nil or negligible additions* to the surface. Soil formation proceeds by effectively modifying pre-existing parent materials to a greater or lesser extent according to a range of factors that dictate a range of processes. However, in many landscapes, such as those of alluvial plains, or where tephras or loess are deposited, aggrading parent materials are very common. The evolution of soils in such landscapes therefore has an additional complexity because the impact from topdown processes is modified by the rates at which new materials are added to the landsurface via geological processes (Almond and Tonkin, 1999). The resultant soils are formed by upbuilding pedogenesis.

Upbuilding pedogenesis is the ongoing formation of soil *via topdown processes* whilst tephras or loess (or alluvium, colluvium) *are concomitantly added* to the land/soil surface as normal geological processes (Lowe and Tonkin, 2010; McDaniel et al., 2012). The resultant soils may show distinctive layering and buried horizons (sometimes referred to as paleosols), forming *multi-layered profiles*. The frequency and thickness of tephra accumulation (and other factors) determine how much impact topdown processes have on the ensuing profile character, and if *developmental* or *retardant* upbuilding, or both, will take place. These terms were coined by Johnson and Watson-Stegner (1987) and Johnson et al. (1990) as part of their dynamic-rate model of soil evolution whereby soils are envisaged to evolve by ‘ebb and flow’ through time (Schaetzl and Anderson, 2005).

- **Developmental upbuilding** occurs when the rate of addition of tephra or loess to the land is incremental and sufficiently slow so that topdown pedogenesis effectively keeps pace as the land gradually rises (a corollary is that each part of the profile has been an A horizon at one time) (Fig. 22).
- **Retardant upbuilding** occurs when a relatively thick layer of tephra (or alluvium, colluvium) is instantaneously added to the surface, or the rate of accumulation of thinner additions is fast, so that the original soil is rapidly buried (overwhelmed), and thus becomes a *buried horizon cut off and isolated* from the new land surface in which pedogenesis begins anew (Fig. 23).

We may see examples of soils dominated by developmental (Tirau) and retardant (Rotomahana) upbuilding.




Fig. 22 Model of upbuilding pedogenesis in tephra deposits and the formation of a multilayered profile (photo at right) over c. 25,000 years in South Waikato. The underlying initial surface is marked here by a thick pinkish-grey layer of Kawakawa Tephra (c. 25,400 cal yr BP). In phase 1, thin, distal tephras accumulate slowly whilst topdown processes imprint weak horizonation features on them as the land surface gradually rises (developmental upbuilding). In phase 2, the sudden deposition of a tephra layer ~0.5 m thick, Taupo ignimbrite here (c. AD 232), from a particularly powerful eruption buries the antecedent soil, isolating it from most surface processes so that topdown processes begin anew on the freshly deposited tephra (retardant upbuilding). In phase 3, incremental tephra deposition on the new soil continues and developmental upbuilding resumes (after McDaniel et al., 2012).




Fig. 23 Very clear example of a multilayered soil formed through retardant upbuilding at Brett Rd near Mt Tarawera. After each tephra is deposited, soil begins to develop until it is buried by another tephra and topdown pedogenesis begins operating in the fresh deposit. The buried 'mini' soil profile on Whakatane tephra (4Bwb, 4C horizons) reflects ~3800 years of pedogenesis, the amount of time it was at the land surface before burial by Taupo tephra; that on Taupo tephra (3ABb, 3BCb, 3C) reflects ~1100 years; that on Kaharoa tephra (2Ahb, 2Bwb, 2BCb) reflects ~570 years; and the topmost (surface) mini profile (Ap, BC(x), C) reflects ~120 years of pedogenesis on hydrothermally altered, mud-rich tephra deposited in AD 1886 by the Tarawera eruption (from McDaniel et al., 2012).

Buried brown rather than black 'topsoil' horizons

Buried 'A' horizons in New Zealand on tephra tend to be brownish rather than dark or black (and hence often have AB or Bw rather than Ah notations) and there is debate as to the reason for this. One suggestion is that in New Zealand they have largely been developed under podocarp-broadleaved forest until very recent times (last c. 700 years) and that such soils, especially Allophanic Soils, tend have brownish rather than dark A horizons anyway (this applies in USA for Andisols under conifers: P. McDaniel, pers. comm., 2008). Alternatively, or in addition, once 'A' horizons are buried then they may, depending on depth of burial, become isolated from the organic cycle and hence no longer receive new organic matter to maintain their darkness via melanisation (Fig. 25).




Fig. 24 Idealised models of buried soil horizons at different depths and how they may be impacted by surficial (topdown) processes (after Schaetzl and Thompson, 2015).




Fig. 25 Comparison of humic acids extracted mainly from surface horizons, and some subsurface (buried) A horizons, on tephra-derived soils in Japan and New Zealand (redrawn after Yamamoto et al., 1989). The 'melanic' status has been included. The carbon content of each soil and that of the associated humic acid were correlated ($r = 0.685$) for the Japanese soils but not for New Zealand soils (Yamamoto et al., 1989). The buried soil horizon on Kaharoa tephra is melanic-like and contains charcoal probably from Polynesian burning.

Residual colours after removing organic matter from A horizons by H₂O₂ or burning in the lab are similar to those of buried horizons on the tephras (P.J. Tonkin pers. comm., 2006). In some cases the depositional (burial) event may ‘scalp’ the topsoils (e.g., during emplacement of the Taupo ignimbrite), leaving effectively subsoils to represent the antecedent (now paleo) land surface. Forest fires following eruption events may also partially ‘bleach’ upper horizons (Wilson, 1994), and the effects of podzolisation (acid leaching), giving rise to bleached ‘E’ (albic) horizons over dark brown or reddish-brown podzolic-B (spodic) horizons are also evident in soils on Holocene tephras including on Taupo, Whakatane, and Rotoma tephras (Lowe et al., 2012).

Impacts of deforestation and burning, and the role of bracken fern, on the properties of surficial or buried soil A-horizons

Bracken fern (*Pteridium* spp.) is an aggressive plant that commonly invades disturbed sites. Its success as an invader is attributable, in part, to its ability to produce abundant growth, both below ground in the form of rhizomes and fine roots and above ground as fronds and stems (Table 5; Lowe and McDaniel, 2010). This biomass production has been shown to affect numerous soil properties. For many years, the black A horizons often found on Pumice Soils have been attributed ‘to bracken fern, which replaced much of the forest’. Analyses of humus and phytoliths in the A horizons of soils developed especially on Kaharoa and Taupo tephras in central North Island (buried beneath 1886 Tarawera eruptives in the parts of Bay of Plenty) showed that type-A humic acids predominated and that fernland and grassland had replaced the pre-existing forests (Fig. 25) (Birrell et al., 1971; Sase et al., 1988; Hosono et al., 1991; Sase and Hosono, 1996). Pollen, phytolith and associated studies, together with tephrochronology, have shown that human-induced deforestation by burning began in New Zealand soon after Polynesian settlers arrived (Newnham et al., 1998; McGlone and Wilmshurst, 1999; McWethy et al., 2009, 2014). The repeated burning resulted in the formation of extensive fernlands (McGlone et al., 2005).

Table 5 Bracken biomass comparisons – New Zealand and northern Idaho, USA*

Location	Rhizome biomass		Frond biomass	
	Mean	Range	Mean	Range
	(kg m ⁻²)		(kg m ⁻²)	
Nelson, New Zealand	--	7.08 (max.)	--	1.41 (max.)
New Zealand (23 stands)	2.92	0.91-5.19	--	--
Idaho, USA (9 stands)	1.96	1.14-2.54	0.52	0.27-0.89

*From Lowe and McDaniel (2010)

In northern Idaho, USA, establishment of bracken is associated with the conversion of allophanic to non-allophanic Andisol mineralogy (Johnson-Maynard et al., 1997). These changes include increased soil carbon, darker soil colours, lower pH, and increased organic forms of active Al. In Japan, a major part of soil organic carbon in Andisols and associated soils comprises humic acids, which are characterized by their stability and aromatic (humified) structure (Shoji et al., 1993; Hiradate et al., 2004). These features arise from the presence of labile and active metals, chiefly Al and Fe, supplied by the weathering of glass and other tephric materials, which are able to bind humic substances through strong coordination bonding to form macromolecules of Al- and Fe-humic acid complexes very resistant to degradation or leaching (Hiradate et al., 2004).

The resultant very dark or commonly black A horizons, typically melanic horizons in *Soil Taxonomy* (Table 6), can contain up to 15% organic carbon (Hiradate et al., 2004). The Japanese pampas grass or ‘susuki’, *Miscanthis*

sinensis, a C4 plant with large amounts of root residues, has been maintained artificially for millennia in ancient Japan to provide roofing material, straw bags, and fodder for grazing. Burning the grass in early spring before germination was common practice (buds of the pampas grass are several centimetres underground and so are not damaged) (Lowe and McDaniel, 2010). During burning, large amounts of very dark-coloured and charred microparticles were produced and these provided a second source of humic acids. The pampas grass has been regarded as a classic ‘andisolizer’ or ‘melaniser’ (Wada, 1986; Shoji et al., 1990, 1993; Hiradate et al., 2004).

Table 6 Main properties of melanic horizon*

-
- Munsell colour values and chromas of ≤ 2 (dark) throughout
 - Melanic index ≤ 1.70 throughout
 - $\geq 6\%$ organic C as weighted average
 - Andic soil properties
 - ≥ 30 cm thick
-

* After Soil Survey Staff (1999, p.23)

The relative importance of these two mechanisms – (1) stabilization of humic acids by complexing reactions with Al and Fe, or (2) cultivation of *M. sinensis* and its charred derivatives as a major source of carbon in humic acids – was evaluated by Hiradate et al. (2004) using $\delta^{13}\text{C}$ analyses of humic substances in a range of soils in Japan. Hiradate et al. (2004) were able to compare the contributions from C3- or C4-plant- derived carbon because *M. sinensis* was effectively the only C4 plant species in (traditional) Japan. They found that the contribution to humic and fulvic acids of carbon derived from *M. sinensis* ranged from 18 % to 52 %. Their conclusions were that although highly-humified, i.e. dark-coloured, humic acids tended to be derived from C4-plant-derived carbon (ultimately *M. sinensis*), the dominant source of carbon for humic and fulvic acids in many cases was from C3 plants (mainly forest vegetation). Consequently, the overriding importance of active Al and Fe especially on the formation and accumulation of the dark-coloured humic acids in *Kuroboku* soils (Andisols) was reinforced (Hiradate et al., 2004).

In New Zealand the ‘black’ A horizons generally have melanic properties, but do not meet the thickness criterion for melanic epipedons, and hence are regarded as ‘melanic-like’. As noted above, they formed under fernland and tussock grassland and most contain A-type humic acids (\pm charcoal). Non-melanic topsoils are formed under forest (most have non A-type humic acids, including Pg in P-type humic acids from deciduous trees) (Figs. 25, 32). Differences between New Zealand and Japan thus relate largely to differences in human settlement history and impacts, with New Zealand having an exceptionally short prehistory of only c. 700 years (since c. 1280 AD) in comparison with >10,000 years in Japan (Lowe and McDaniel, 2010).

Breaking in the land acre by acre

By Penelope Hall

NGUTUNUI dairy farmer Keith Pairiki Ormsby has worked tirelessly on the land he broke in. And says he couldn't have done it without the help of his parents and his wife Margaret.

"Margaret and I are a team. It's as simple as that."

ONE OF FOUR

Keith dates his farming legacy back to his great grandparents.

Robert Ormsby (1823-1920) who emigrated from Ireland, married Meri Rangihurhia Pianika Takiari (1828-1905) of Ngati Maniapoto in 1848.

Although they weren't farmers, it was Meri's land that has been farmed by their offspring through the generations.

Including their third child, Keith's grandfather, Arthur Ormsby (1853-1926) who married Matire Wright (ca 1855-1935).

And their 13th child was Keith's father, Robert Ormsby (1899-1967) who married Mabel Randall (1904-1988).

Keith was born in 1936 to the Puketotara dairy farmers.

He was the third of Robert and Mabel's four children — Noel then Gloria — followed by younger sister Beverley.

Keith attended Pirongia School and Te Awamutu College but left to go farming once he earned his School Certificate.

He says: "I was always going to be a farmer, so I stayed working on mum and dad's farm after that."

That ended when his parents moved to Pirongia in 1955 and Noel took



over the property.

BUYING A BLOCK

Keith, then aged 18, went off to forge his own future, buying 52.60ha (130ac) of maiden bush on Pekanui Rd, in Ngutunui.

"My parents couldn't help financially, but they went guarantor for the loan and helped me break in the land," he says.

"I'd use the chainsaw, dad would take out the undergrowth, mum would pick up the scraps and burn them and I'd pull up the stumps with the digger.

"When I moved up here I built a wooden lean-to for the fertiliser, with a little room where I lived. It was hard work, but the fact my parents

were helping me, made me stick with it."

Trees were either milled or burned, but Keith retained some bush including everything surrounding the 10 waterways running through the property.

As the land was cleared, he'd add livestock until he had 14 cows, nine heifers and some sheep.

He says: "That's when a couple of uncles came and helped me build my first cow shed in 1958.

"It was a three-bay walk through unit, which still stands on our property today and meant I could convert to milking.

MEETING MARGARET

It was then he met Margaret Keltie, originally from Thames and a sec-



WHEN Keith and Margaret married in 1961 they spent the first winter in the hand-built lean-to's living space at Ngutunui. Margaret says it was tough going.

retary at Te Awamutu legal firm Edmonds Judd and Goldfinch.

Keith says the two were "set up on a blind date" at the Hamilton River Carnival by a good mate.

"She was a challenge, but I thought she was alright."

The couple married in Te Awamutu in May of 1961, settling into married life in the lean-to's living space at Ngutunui, alongside farming tools and equipment.

to page 3



KEITH (left), pictured with his brother Noel, cleared 242.81ha (600ac) of bush from 1955 while retaining native trees surrounding the property's 10 waterways.

from page 2

"Living there through the winter was tough," says Margaret who was "happy" to move into their newly built home on the property in November.

That same year, the couple bought an adjacent 52.60ha (130ac) bush block and Keith continued clearing it between milking twice a day.

Margaret continued working full time until Christmas, before the birth of the first of their

four children — Kaye born in 1962, followed by Brett, Ross and finally Angela, in 1969.

MARRIED LIFE

Keith remembers going out in the morning darkness to milk, spending all day with his father clearing bush, getting the cows in for milking in the late afternoon, then coming home in the dark after the children were in bed.

"It was hard for me to do much milking with four children running around," says Margaret.

"But I reared calves and did all the milking when Keith was haymaking"

The dairy herd increased to about 120 by the early 1970s which is when Keith felt he'd cleared enough land to run some drystock.

"I chose romney perendale sheep because they do well in hill country and friesian bulls because they're the closest to beef bulls.

"Margaret and I would buy calves locally, rear them through to two-year-olds and sell them on as bull beef. America took the bull beef ungraded,


so it was a good way to go financially."

DEVASTATING

In 1984 life took a horrific turn when their 16-year-old son Ross broke his neck in a collapsed scrum during Te Awamutu College's 1st XV rugby trials.

The incident left him a tetraplegic. "It was devastating," says Margaret.

to page 4



DESPITE the challenges of breaking in his Ngutunui farm, Keith says the land means everything to him. On this part of the property there is a clear view of Mt Kakepuku.

Breaking in the land acre by acre

from page 3

"I stayed with Ross for three months in the Ota-ara spinal unit and only went home once in that time. When I did, is when I crashed [mentally].

"I needed to get back up there. I felt as long as I was there with him, it was alright no matter how sick he was.

Margaret says they always looked on the bright side which helped get them through and now "he's absolutely amazing doing things most people don't do.

"He's a bit of a daredevil and lives his life to the fullest".

During the mid-1980s, Keith — by this time in his 60s — wanted to slow down while getting the best financial return so they entered into their first 50/50 sharemilking contract. And it's been that way ever since.

Keith and Margaret bought a second adjoining farm in 2001 to expand their holding to 267.09ha (660ac) total, of which 242.81ha (600ac) is effective rolling contour land milking 580 cows.

Keith says: "Our [current] sharemilker Gra-hame Wallis has done a great job here for the past 10 years.

"I drive around the farm and it's in great order,

so I don't need to be out there anymore."

No longer tied to the farm, the couple have been travelling the world ever since.

"We've been lucky," says Margaret.

"But people don't know what it was like in the early days when we didn't have anything.

"The one thing we always did, was pour all our money back into the farm, otherwise we wouldn't have what we've got today."

REPLANTING THE BUSH

Keith and Margaret are most proud of replac-ing some of the bush that was felled to create

the farm.

So, with the help of their children and grand-children, 27,000 trees and plants were reintro-duced between 2006 –2015.

They have also protected 15ha (37.06ac) with two QEII National Trust covenants.

In 2009, Keith and Margaret were presented with PGG Wrightson Land and Life award for sustainable farm practices and leadership.

And in 2014, they were recipients of Donaghys Farm Stewardship and Waikato River Catchment improvement awards.

to page 5

NKG Farmer Tuesday, December 11, 2018 5



KEITH says he couldn't have made the farm what it is today without Margaret on his "team" for the past 57 years.



A FAMILY get together in 2003, included Matekino (Nig) Roa (left), Mary Armstrong, Keith, Jean Bell, Margaret, Noel Ormsby, Ngaire Ramage and Bill Armstrong. The event was held to bless the adjoining land purchased by Keith and Margaret two years before.

Breaking in the land acre by acre

from page 4

It was about the same time when Keith suffered a heart attack and an angioplasty was performed, which was followed by a second last December.

He says: "I was in mid-sentence when I dropped.

"I died, but a policeman passing by gave me CPR and brought me back to life."

MY VOCATION

Now aged 82, Keith looks back on being a farmer saying, "I never thought about being a farmer, it's just what I did, it was my vocation."

"I don't think about what I've achieved," he

says.

"But when I drive through the farm I know why we're still here more than 60 years later.

"Buying the land was a challenge, breaking it in was a challenge, changing over to dairy was a challenge. It's been many challenges all the way through.

"But I cut it out of the bush.

"It means everything to us, we're connected to it."

So big is Keith's love of the Pekaui Rd prop-erty that one section of bush is known by the family as Pairiki's Last Stand.

"It's the place I want to go when I die."

KEITH and Margaret have been involved in many activities off the farm as well..

Rugby — Keith played and coached for the Pirongia Rugby Club and coached. He was presented a life membership by former Wallaby Nick Farr-Jones at the club's 125-year celebrations in 2001. He also captained the Te Awamutu Repre-sentative team and later played for Waikato Maori Rugby during their Samoa, Fiji tour in 1962.

Horse racing — the Ormsby family have now been part of the Pirongia Boxing Day Races for six generations and Keith, a life member since 2011, counts his mother being pregnant with him at the event as his first time and been there every year since.

Golf — Keith and Margaret have been part of the

Pirongia Golf Club since the 1970s. Keith has also played for the Eagles Golf Society since 1990.

Tennis — the Ngutunui Tennis Club presented them both with life memberships in 1985.

School — Ngutunui School, facing closure due to low student numbers in 2004, brought Keith in as chairman of the board of trustees. He knocked on the doors of new parents in the area asking them to send their children to the school as well as requesting parents who'd taken their children out, to reconsider. Then in 2005 when the Ministry of Education withdrew transport funding Margaret instigated a calf rearing scheme raising more than \$25,000 for the school to buy their own minivan.

Singing — Margaret is a founding and lifetime member of the Rosebush Chirsters

STOP HOROTIU SANDY LOAM, LEAMINGTON

Location BD34 159025, elevation ~60 m asl, rainfall ~12200 mm pa

Beware of traffic

Horotiu sandy loam with parent material of composite, intermixed late Quaternary tephra-fall mantle over volcanogenic alluvium (Hinuera Formation). Discuss origin of Hinuera Formation (deposited c. 20,000 cal. yr BP at this location or soon after). Cores from lakes in the area have provided evidence for continuing incremental accumulation of tephra-fall deposits since deposition of the Hinuera Formation ceased. Note views of terraces along Waikato River.

Soil Taxonomy: *Vitric Hapludand*

NZSC: *Typic Orthic Allophanic Soil*



Mamaku Plateau – development of complex ignimbritic landscape




Fig. 26 Left. General model for landscape inversion in an ignimbrite terrain involving the emplacement of an ignimbrite as valley-pond and veneer deposits, and the subsequent erosion of the ignimbrite to form wide mesas and narrow buttes, and smaller tors, alongside deep valleys (after Hill et al., 1999). **Right.** General development of landforms in an ignimbrite sheet comprising three units with different degrees of welding, forming gorges then widened valleys flanked by mesas and buttes. Tors may originate where fumarolic activity has caused strengthening by secondary mineral deposition and alteration. Modified after M.J. Selby in Healy (1992, p. 261).




Fig. 27 Topographic relief model illustrating some of the physiographic regions covered partly on Wednesday 21 Dec 2016. The 'geomorphic' Taupo Volcanic Zone (TVZ) comprises an area of largely volcanic hills and lakes and covers a similar area (but not exactly equivalent) to the volcanological TVZ (defined by vent locations only) (from Leonard et al., 2010, p. 7).




Fig. 28 Soil-landscape model for the Tirau-Putaruru area and Tapapa. The three main welded ignimbrite units (~1.23 Ma to 0.9 Ma, erupted from Mangakino volcano) form distinctive plateaux in the landscape and are mantled with tephra-fall deposits and subordinate tephric loess. Fluvial terraces are also mantled with tephra. TPA, Taupo Pumice Alluvium (c. 1700 cal yr BP) (diagram modified from McLeod, 1992).



Fig. 29 Photos of Tapapa section (upper, middle, lower) and modern Tirau soil in upper part of sequence. Tephras preserved in nearby Lake Okoroire (~10 km NW of Tapapa) provide a record of tephra fall since c. 20 cal ka (Lowe, 1986, 1988) (from Lowe et al., 2012).

TIRAU SILT LOAM

Grid ref: N66/271251

Location: Hetherington Road, Tirau, on road side
 Aspect: - 1.5 km east of Tirau, 1 km south of cemetery
 Vegetation: Improved pasture species - rye grass, white clover, paspalum
 Land use: Dairying, breeding and fattening sheep and cattle, maize cropping

Altitude (m): 120
 Rainfall (mm): 1400

Slope: 2°
 Landform: Downland
 Drainage class: Well drained
 Parent material: "Tirau Ash" consisting mainly of Taupo Pumice, Rotorua Ash, tephric loess, Kawakawa Formation, Rotoehu Ash, Hamilton Beds

PROFILE DESCRIPTION

Horizon	Depth (cm)	Description
Ap	0-18	very dark brown (10YR 2/2) gritty silt loam; friable; moderately developed medium nut structure breaking to moderately developed fine nut and crumb structure; abundant roots; few fine Taupo lapilli; distinct irregular (worm mixed) boundary,
Bw1	18-30	dark yellowish brown (10-7.5YR 4/4) sandy loam; very friable; moderately developed medium crumb structure; many roots; distinct irregular boundary,
2Bw2	30-51	yellowish brown (10YR 5/6) sandy loam; friable; moderately developed medium nut structure breaking to moderately developed fine nut and crumb structure; many fine and coarse roots; few very thin discontinuous humus coatings on root channels; few very fine lapilli; indistinct smooth boundary,
3Bw3	51-74	yellowish brown (10YR 5/6) greasy sandy loam; friable; moderately developed coarse nut structure breaking to fine nut and crumb structure; many roots; many thin discontinuous strong brown (7.5YR 4/6) coatings (iron) on peds; distinct smooth boundary,
3C	74-100	brown near (10YR 5/4) silt loam; firm; moderately developed medium block structure breaking to fine and medium nut structure; few roots; many fine dark brown (7.5YR 3/4) stains along root channels; few fine lapilli.

Description of Tirau soil at Hetherington Rd, near Tirau (T15/546556) (~18 km from Tapapa) from Parfitt et al. (1981) (see also Bakker et al., 1996). At Tapapa the soil was mapped by McLeod (1992) as part of the Tirau series, which occur on rolling land grading to higher terraces (elevational range 70-400 m) and with a thermic temperature regime. However, Tirau soils have very similar parent materials to soils of the Waiohotu series that occur at usually higher elevations on high-level terraces (plateau surfaces) (elevational range 250-500 m) under higher rainfall (Waiohotu 2000-2400 mm pa, Tirau 1400-2000 mm pa) and mesic temperature regimes (see soil-landscape model p.10). The Waiohotu soils have slightly coarser textures (gritty silt loams vs silt loams), lower pHs (Waiohotu 4.5-5.9, Tirau 5.6-6.4), stronger leaching, and until recently have not normally been used for maize production (M. McLeod pers. comm., 2008). Tapapa is at the thermic/mesic boundary (Aldridge, 1982).

Classification of Tirau silt loam

NZSC: Acidic Orthic Allophanic; Mt; na (Rh and An fines), loamy; m

Soil Taxonomy: Medial, thermic/mesic Typic Hapludands


CHEMISTRY TIRAU SILT LOAM																			
Sample No. SB	Depth (cm)	Hor.	pH			Exchangeable cations (meq/100 g)				Extr. Acidity (pH 8.2)		Acidity-Al (meq/100 g)	ECEC	CEC (meq/100 g)		Base saturation (%)			
			H ₂ O	KCl	ΔpH	NaF	Ca	Mg	K	Na	H (KCl)			Al (KCl)	NH ₄ OAc (pH 7)	Σ Cations (pH 8.2)	Σ bases CEC NH ₄ OAc	Σ bases Σ Cations	
9432																			
A	0-18	Ap	5.6	4.8	-0.8	9.6	12.4	0.90	1.02	0.12		0.15	43.8	43.6	14.6	29.7	58.2	48	25
B	18-30	Bw1	6.2	5.6	-0.6	9.9	4.2	0.48	0.82	0.13		0.00	25.0	25.0	5.6	11.3	30.6	50	18
C	30-51	2Bw2	6.2	5.8	-0.4	9.8	4.1	0.61	0.27	0.19		0.00	23.4	23.4	5.2	10.5	28.6	50	18
D	51-74	3Bw3	6.4	5.5	-0.9	9.5	5.3	0.38	0.16	0.27		0.00	20.3	20.3	6.1	12.5	26.4	49	23
E	74-100	3C	6.4	5.5	-0.9	9.4	6.5	0.45	0.13	0.44		0.03	18.3	18.3	7.5	13.2	25.8	57	29

Sample No. SB	Depth (cm)	Hor.	Total C (%)	Total N (%)	P (mg/100 g)			P Retention (%)	Dithion. cit. (%)		Tamm ox. (%)		Pyrophos. (%)		Reserves (meq/100 g)		Extractable S (ppm)	
					H ₂ SO ₄ (0.5 M)	Inorg.	Org.		Fe	Al	Fe	Al	Si	Fe	Al	K _c		Mg _r
9432																		
A	0-18	Ap	7.9	0.71	129	128	121	88	0.81	0.83	0.59	3.3	1.32	0.17	0.72	0.12	1.9	125
B	18-30	Bw1	2.0	0.16	30	34	28	98	1.00	0.59	0.72	4.1	2.2	0.02	0.22	0.14	1.4	418
C	30-51	2Bw2	1.0	0.09	25	28	16	98	1.00	0.54	0.69	4.0	2.6	0.00	0.15			374
D	51-74	3Bw3	0.5	0.04	13	16	7	91	1.12	0.40	0.41	2.8	1.71	0.00	0.10			157
E	74-100	3C	0.5	0.04	7	9	7	86	1.13	0.36	0.49	2.4	1.29	0.00	0.09			112

PHYSICS										
Hor. Depth (cm)	Hor.	15 bar water Field moist (%)	Air Dry (%)	Core Depth (cm)	Dry bulk density (T/m ³)	Total porosity (%)	Large pores (%)	Field Cap. (at 0.2 bar) (% v/v)	Wilting Pt. (at 15 bar) (% v/v)	Available water (% v/v)
0-18	Ap	31.1	23.3	2-9	0.75					
18-30	Bw1	33.7	15.7	20-27	0.71					
30-51	2Bw2	33.6	16.5	34-41	0.69					
51-74	3Bw3	34.7	22.0	58-65	0.79					
74-100	3C	33.9	20.4	80-87	0.87					

PARTICLE SIZE DISTRIBUTION (<2 mm) TIRAU SILT LOAM									
Sample No. SB	Depth (cm)	Hor.	Sand		Silt	Clay	Fine clay	Fine clay Total clay	Stones (%)
			2-0.1 mm (%)	0.1-0.05 mm (%)	0.05-0.002 mm (%)	<0.002 mm (%)	<0.0002 mm (%)		
9432A	0-18	Ap	18	17	46	19			
B	18-30	Bw1	17	16	56	11			
C	30-51	2Bw2	17	16	54	13			
D	51-74	3Bw3	14	16	50	20			
E	74-100	3C	6	14	62	18			

SB 9432 Tirau Silt Loam



Mineralogy TIRAU

Sample No. SB	Depth (cm)	Hor.	Clay Fraction (%)													
			Mica-Smectite	Mica-Vermiculite	Smectite	Vermiculite	Interlayered Hydrous Micas	Mica	Kaolinite	Halloysite	Gibbsite	Quartz	Cristobalite	Allophane	Feldspar	Anatase
9432A	0-18								8				65			27
B	18-30							6					75			19
C	30-51							4					80			16
D	51-74							9	tr				55			36
E	74-100							21	tr		tr		48			31

Anthropogenic fluorine accumulation in the Waikato and Bay of Plenty regions of New Zealand: comparison of field data with projections

Nicholas D. Kim, Matthew D. Taylor, John J. Drewry

Environmental Earth Sciences (2016) 75: 147 (DOI 10.1007/s12665-015-4897-2)

Long-term annual application of phosphate (P) fertilizers to farmland may be causing worldwide accumulation of fluorine (F) in productive soils, as F concentrations in rock phosphate and P fertilizers are typically ~75–150 times higher than those in recipient soils. Possible impacts of excess anthropogenic F in agricultural soils fall into three categories:

1. Toxicity, where concerns include toxicity to soil organisms and terrestrial wildlife, phytotoxicity, chronic fluorosis in grazing animals, and potential for aluminofluoride complexes to interfere with biochemical signalling pathways. AlF_x complexes are of toxicological interest because they act as phosphate analogues in a variety of enzymes.
2. Alterations of soil chemistry and function. Excess F may reduce organic matter turnover (Rao and Pal 1978) and phosphatase activity accelerate aluminosilicate weathering, induce formation of AlF_x complexes in pore-water, and alter uptake of F, Al, AlF_x and other F-complexed elements in crops and animals.
3. Wider environmental impacts, including potential for contamination of ground-water, toxicity to aquatic ecosystems, and perturbation of the atmospheric F cycle from fluorinated gases released from treated soils or during fertilizer manufacture. Volatile F-containing species are effective greenhouse gases.

We report results of sampling across two large regions of New Zealand involving soils of 298 native and farmed properties. This has enabled the first wide-scale retrospective estimates of anthropogenic F enrichment in pastoral and horticultural soils. Results validate earlier projections: F accumulation in farmed soils has been comparatively rapid (mean increase 2.1 % per year) and widespread. Over 50 years, average total F concentrations in surface soils have doubled from ~220 to 440 mg/kg. Thresholds protective against chronic fluorosis in grazing animals are being substantively passed, indicating that land may be rendered unsuitable for pastoral production in the mid-term future: when sampled 44 % of dairy farm soil samples exceeded 500 mg/kg total F, and 10 % exceeded 650 mg/kg. Potential for F toxicity to plants and wildlife appear to be significant and marginal, respectively, at current soil F concentrations. Relationships between F and other elements suggest P fertilizers are also a source of several elements geochemically linked to F, and that anthropogenic F modifies soil aluminium (Al) chemistry, potentially opening a broad-scale Al exposure pathway to pollinating insects.

Table 3 Modern (mod) and estimated natural background (bkd) mean concentrations (mg/kg) of F ordered by soil order and land-use category

Soil order	Pastoral soils						Horticultural soils						All farmed soils	
	Dairy		Sheep and beef		All Pastoral ^a		Arable		Orchard		All hort		N	mod/bkd
	N	mod/bkd	N	mod/bkd	N	mod/bkd	N	mod/bkd	N	mod/bkd	N	mod/bkd		
Allophanic	32	554/242	25	444/228	57	505/236	12	483/235	13	533/215	25	509/225	82	506/233
Brown	10	382/216	10	407/215	21	387/215	2	485/208	0	–	2	485/208	23	395/214
Gley	29	467/229	6	457/238	35	465/231	5	361/236	0	–	5	361/236	40	452/231
Granular	9	481/204	14	336/212	25	403/210	7	377/208	1	680/208	8	415/208	33	406/210
Organic	11	434/51	9	346/51	20	394/51	0	–	1	119/51	1	119/51	21	381/51
Podzol	0	–	3	351/254	3	351/254	0	–	0	–	0	–	3	351/254
Pumice	22	467/248	12	392/246	35	437/247	1	491/208	2	345/254	3	394/239	38	434/247
Recent	7	433/234	4	292/235	11	377/233	4	330/243	2	245/231	6	302/239	17	351/235
Ultic	0	–	7	349/247	7	349/247	0	–	0	–	0	–	7	349/247
All orders	120	481/217	90	389/212	214	440/215	31	420/227	19	469/212	50	438/221	264	440/216

^a Totals include four pastoral sites that are neither dairy nor sheep and beef (one brown, two granular, one pumice)

Comparing volcanic glass shards in unfertilised and fertilised Andisols derived from rhyolitic tephra, New Zealand: evidence for accelerated weathering and implications for land management

Matthew D. Taylor, David J. Lowe, Peter Hardi, Geerd A. Smidt, Ewald Schnug

Geoderma 271 (2016) 91–98 (<http://dx.doi.org/10.1016/j.geoderma.2016.01.035>)

Enhanced weathering associated with the use of phosphate fertilisers has been identified in some of the major farming areas of NZ and an evaluation of its effects on soil properties and the implications for soil management are needed. We assessed changes in the character of volcanic glass shards in topsoils (0–10 cm depth) of two tephra-derived Andisols of differing ages (Udivitrand, Hapludand), and with and without long-term fertilisation, to test if fertilising accelerates the weathering of soil constituents. Using visual assessment based on scanning electron microscopy and electron microprobe analyses of glass shards from samples from paired sites, we showed that the average sizes of the shards and the sharpness of the glass-shard edges (angularity) diminished with the age of the soil, and that these decreases were more marked in the soils that had been fertilised. Silica polymorphs were observed only in the older soil (Hapludand) that had been fertilised. The addition of phosphate-containing fertiliser enhanced the dissolution of volcanic glass, consistent with phosphoric acid and F⁻-induced dealumination and desilication, and thus soil weathering has been accelerated. The Al and Si may subsequently coprecipitate as secondary minerals, such as allophane. However, silica polymorphs may form where Al activity is low. The occurrence of neogenic silica in topsoils (despite high annual rainfall that should result in desilication of the topsoil and reprecipitation of silica deeper in the soil) is explained by the seasonality of fertiliser application and the spring–summer climate, where the soil remains moist for periods sufficiently long enough for the dissolution of volcanic glass (and possibly of other aluminosilicates), the formation of Al–humus complexes, and the subsequent precipitation of silica in dry periods.

Table 3

Comparison of glass-shard sizes for samples from fertilised and unfertilised soils, and roundness classes (range 0 to 6) of Powers (1953), and presence of neogenic silica polymorphs.

	Taupo sandy loam (Udivitrand)				Tirau silt loam (Hapludand)			
	Average µm	Median µm	Range µm	n	Average µm	Median µm	Range µm	n
Unfertilised	39	33	12–231	104	37	25	22–252	101
Fertilised	31	23	7–246	102	24	22	8–94	100
Roundness	Mainly classes 0 & 1 (very angular to angular) with class 1 predominant in fertilised soil				Mainly classes 2 & 3 (subangular to subrounded) with class 3, and pitting, predominant in fertilised soil			
Neogenic SiO ₂ polymorph	Not present in either unfertilised or fertilised soils				Present in fertilised soil only			

Implications for soil management and health emerge because of the accelerated weathering. Some trace elements, such as Co, are already at levels below optimum for animal health in these soils and increased leaching of these could reduce the efficiency of corrective fertiliser additions. Increased direct dosing of animals with essential trace elements may be required in the future. Other implications for soil management include changes in soil permeability and water storage capacity. Loss of mass may lead to increased porosity of the soil and enhanced permeability, if it is undisturbed by further cultivation. Conversely, precipitation of clays within pore spaces may reduce soil permeability but increase water storage. Changes in porosity are also likely to cause changes in surface area and capacity to store cations and anions. More broadly, there may be a significant feedback mechanism between climate and weathering processes. Enhanced weathering could result in enhanced removal of CO₂ from the atmosphere, as soil carbon derived ultimately from photosynthesis is converted into dissolved HCO₃⁻. In addition, weathering releases divalent cations to the ocean via groundwater and riverine transport. Once there, divalent cations promoting the drawdown of CO₂ from the atmosphere by the precipitation and subsequent burial of carbonate mineral. If the enhancement of weathering is significant, it may provide some mitigation for climate change.

LAKE OKARO AND WETLANDS, OKARO RD

Location U16 177171, elevation 420 m asl, rainfall ~1500 mm pa

Geology and soils in Lake Okaro area

Lake Okaro was formed c. 700 years ago, probably during the Kaharoa eruption (1314 ± 12 AD), by a phreatic (i.e. steam) eruption (Lowe and Green, 1992; Lowe, 2006). The geology and soils of the area were described by Cross (1963), and soil units were recorded on the Rerewhakaaitu sheet (scale 1: 31,680) by Vucetich and Cross (1978) (also in Vucetich and Wells, 1978). Soils on the lake's western margins were mapped mainly as Rotomahana silt loam, mottled phase. Studies of rill erosion on the hills to the north of Lake Okaro, and on mechanisms and effects of the phreatic and secondary hydrothermal eruptions that formed the lake, were undertaken by Hardy (2005). The rilling, well exposed between Waimangu and Lake Okaro, is formed entirely in the ~1 m-thick mantle of Rotomahana Mud deposited in the area on 10 June 1886. The rill formation was rapid, beginning within days of the eruption and ending within a few years. The reason the rill erosion ceased is unknown but is possibly related to the permeability of underlying layers (Hardy, 2005). Rills are defined as centimetre to decimetre scale erosion channels cut by water running down a slope. Gullies are defined on a metre scale. In the USA, rills are described as small, runoff channels that can be obliterated by conventional tillage whereas gullies are large, runoff channels that cannot be obliterated by conventional tillage (Schoeneberger et al., 2002).

Lake Okaro and water quality (from Özkundakci et al. 2008)

Lake Okaro has an area of 0.28 km² and maximum depth of 18 m. It is monomictic and is typically thermally stratified for eight months of the year, from October until June. Limnological records for Lake Okaro extend back to 1955 (Jolly, 1977) when no cyanobacteria were observed in the lake. By contrast, a persistent cyanobacterial bloom was observed in 1963 (Forsyth et al., 1988). Today, Lake Okaro is the most eutrophic lake in the Rotorua region as a result of excess nutrients which promote primary production and algae growth. To meet statutory obligations, Environment Bay of Plenty, as part of its Land and Water Plan, planned the following restorative measures: (i) a phosphorus-absorbent lake-bed cap; (ii) a constructed wetland to remove primarily N from stream flows (iii); the protection of all riparian margins in the catchment; and (iv) the introduction of best management practices to reduce N leaching from land-use over time. The Regional Water and Land Plan for the Bay of Plenty sets water quality targets for the Rotorua lakes on the basis of trophic level index (TLI) (Burns et al., 1997). TLI values are determined annually from annual mean surface water concentrations of chlorophyll *a*, total nitrogen (TN), total phosphorus (TP), and Secchi-disk depth (SD). In the Lake Okaro Action Plan (2006), the TLI goal for the lake is 5.0. Between 1991 and 2008 the TLI of Lake Okaro ranged between 5.09 and 5.94, which classifies the lake as supereutrophic (TLI 5.0–5.99). Considerable experimental work on lake restoration has been carried out on Lake Okaro (see LERNZ website and publications by Univ. of Waikato staff).



Fig. 30 Wetland soon after completion at Lake Okaro (2005) (LERNZ website)

STOP – ROTOMAHANA SILT LOAM, BRETT RD

Location V16 146168, 38° 17.905 S, 176° 28.823 E; elevation ~454 m




Fig. 31 Modern soil, buried soil horizons and tephra layers at Brett Rd section. Ages from Lowe et al. (2013). Rotomahana Mud was erupted from Rotomahana Crater and deposited through the air on 10 June 1886, i.e. it is a tephra-fall deposit, albeit an unusual one compositionally, containing ~20% clay with a distinctive mineralogy (Kirkman, 1976), described below.



Fig. 32 Aerial view of landscape around Lake Rerewhakaaitu showing landforms. *Photo:* D.J. Lowe. Lake Rerewhakaaitu lies in a shallow basin impounded initially by pyroclastic deposits of the Waiohau eruption ca. 14,000 cal. yr BP, with further damming to present levels by Kaharoa pyroclastic deposits in ~1314 AD. It has an area of 7.4 km² and a maximum depth of 15 m. Awaatua basin is a small explosion crater ca. 10,000 years old (Nairn, 2002).

Soils and land use in the Waio tapu region (including Rerewhakaaitu district) were described by Vucetich and Wells (1978). The Rerewhakaaitu area was surveyed by Derek Cross (scale 1: 31,680). Analytical data for the Rotomahana soil are available in Parfitt et al. (1981; data below) and in Part 3 of 'Soils of New Zealand' (1968) (pp.122-123) from a site ~400 m west of the Waimangu Tearooms. Phytolith and humus analyses were undertaken at that site by Hosono et al. (1991); the clay minerals were examined by Kirkman (1976), who identified allophane, interstratified montmorillonite/mica, montmorillonite, mica, kaolinite, and a zeolite mineral, mordenite.

Notes from Mac Pacey, farmer on 'The Mud', Tulachard Farm, Brett Rd

Advantages: Naturally good fertility. High in Ca, Mg and K. Mainly fertilise with P and S. Good pH. Holds on better than surrounding country in dry periods. We stay green and do not have plant deaths in dry periods; therefore when it rains we recover quite quickly. When conditions are good we grow grass!

Disadvantages: When it rains and the soil is sodden the pasture can turn to mud very quickly. To overcome this we have built a feed pad to feed cows and stand them on in particularly wet conditions. Once it stops raining we can go back on the pasture within 24 hours with care. 'Mud' is high in Mb which causes Cu deficiency in the cattle so we have to supplement with Cu.

Note from soil surveyor Wim Rijkse

Rotomahana coarse sandy loam (near Waimangu Tearooms) is moderately to slightly acid and weakly leached (base saturations 71%, 65%, 94%). Exchangeable cations (Ca, Mg, K and Na) are of medium to high levels. Reserve potassium and magnesium of the Rotomahana Mud are of medium to high value. Extractable phosphorus is low.

Soil profile description by Haydon Jones

Waikato Regional Council, Hamilton

Rotomahana reference data

- Soil name:
 - Series: Rotomahana
 - Type: silt loam
- Soil classification:
NZSC: Buried-pumice Tephric Recent; Mt; na (Rh fines and Ba 'fines'/stones); loamy; m/r
Soil Taxonomy: Fine-loamy/medial, mixed/glassy, active, mesic Typic Udivitrands

Site data

- Location:
 - Map reference: NZMS 260 V16 2814530 6316815
 - Word description: road cutting on Brett Road (western side of Lake Rerewhakaaitu) about 215 m north of reserve/boat ramp
- Elevation: 450 m
- Rainfall: ~1500 mm pa
- Temperature:
- Geomorphic position: Profile on a 6° convex/linear slope with 125° aspect contained within a gentle slope in easy rolling country
- Erosion/deposition: Nil
- Vegetation: Pasture, grasses
- Parent material: Rotomahana Mud over a thin layer of Tarawera Scoria with underlying buried soil horizons formed in various tephra deposits (Kaharoa, Taupo, Whakatane, and Rotoma tephra) (see photos of stratigraphy for tephra ages or see summary p. 27)
- Drainage class: Well drained

Soil data

Note: "gravels" = "stones".

Ap

0-11 cm Very dark greyish brown to dark brown (10YR 3/2.5) silt loam; slightly sticky; moderately plastic; peds weak and friable; weakly pedal; common fine polyhedral peds; non allophanic; abrupt smooth boundary. [Rotomahana]

BC(x)

11-36 cm Greyish brown (2.5Y 5/2) silt loam; very few extremely fine faint brown (7.5YR 5/4) mottles; slightly to moderately sticky; very plastic; peds slightly firm and brittle; very high penetration resistance; weakly pedal; common fine to coarse blocky peds; non allophanic; abrupt smooth boundary. [Rotomahana]

Cu

36-39 cm Greyish brown (2.5Y 5/2) extremely gravelly sand with fine to medium sub-rounded slightly weathered scoria gravels; non sticky; non plastic; apedal single grain; non allophanic; abrupt smooth boundary. [Tarawera]

2bAh

39-46 cm Black (2.5Y 2.5/1) sandy loam; slightly sticky; non plastic; peds weak and friable; apedal earthy; profuse very fine polyhedral peds; weakly allophanic; distinct smooth boundary. [Kaharoa]

2bBw

46-68 cm Dark greyish brown to brown (10YR 4/2.5) very slightly gravelly loamy sand with fine sub-rounded slightly weathered pumice gravels; slightly sticky; non plastic; soil very weak and very friable; apedal single grain; moderately allophanic; indistinct wavy boundary. [Kaharoa]

2bBC 68-79 cm	Light olive brown to light yellowish brown (2.5Y 5.5/3) very slightly gravelly loamy sand with fine sub-rounded slightly weathered pumice gravels; very few extremely fine distinct brown (7.5YR 4/4) mottles; slightly sticky; non plastic; soil weak and friable; apedal massive; strongly allophanic; distinct wavy (occluded) boundary. [Kaharoa]
3bAB 79-101 cm	Very dark grey (2.5Y 3/1) very slightly gravelly sandy loam with fine sub-rounded slightly weathered pumice gravels; slightly sticky; slightly plastic; peds weak and friable; apedal earthy; profuse very fine polyhedral peds; strongly allophanic; distinct irregular (occluded) boundary. [Taupo]
3bBC 101-138 cm	Olive brown to light olive brown (2.5Y 4.5/3) very slightly gravelly (= not stony) loamy sand with fine sub-rounded slightly weathered pumice gravels; slightly sticky; non plastic; soil very weak and friable; apedal massive; strongly allophanic; distinct wavy boundary. [Taupo]
3Cu 138-151 cm	Pale yellow (2.5Y 7/3) extremely gravelly (= extremely stony*) sand with fine to medium sub-angular slightly weathered pumice gravels; non sticky; non plastic; apedal single grain; very weakly allophanic; distinct wavy boundary. [Taupo]
4bBw 151-173 cm	Very dark greyish brown to dark brown (10YR 3/2.5) slightly gravelly sandy clay loam with fine sub-rounded slightly weathered pumice gravels; moderately sticky; slightly plastic; peds weak and friable; apedal earthy; profuse very fine polyhedral peds; strongly allophanic; indistinct irregular (occluded) boundary. [Whakatane]
4bBC 173-199 cm	Yellowish brown (10YR 5/6) very gravelly sand with medium sub-rounded slightly weathered pumice gravels; non sticky; non plastic; apedal single grain; strongly allophanic; distinct irregular boundary. [Whakatane]
5bAh 199-225 cm	Dark greyish brown to brown (10YR 4/2.5) very slightly gravelly sandy loam with fine sub-rounded slightly weathered pumice gravels; slightly sticky; non plastic; peds very weak and friable; apedal earthy; profuse very fine polyhedral peds; moderately allophanic; indistinct wavy boundary. [Rotoma]
5bBC 225-255 cm	No data. [Rotoma]
5Cu 255 cm –on	No data. [Rotoma]

* “Extremely gravelly or stony” implies the horizon is clast-supported hence there is no need to use term ‘loose’.

ROTOMAHANA LOAM

Location: Waimangu Road, 400 m west along from tearoom,
east side of road reserve
Aspect: West near top of ridge Altitude (m): 400 Rainfall (mm): 125
Vegetation: Bracken fern with mahoe, coprosma, kamahi, lupin
Land use: Dairying and semi-intensive sheep farming

Grid ref: N85/845863
Slope: 2° Landform: Ridge
Drainage class: Moderately well drained
Parent material: Rotomahana Mud, hydrothermally altered rhyolitic ejecta from Lake Rotomahana (erupted in 1886)

Profile	Depth (cm)	PROFILE DESCRIPTION
Ah	0-7	very dark brown (10YR 2/2) loam; slightly sticky non plastic; slightly greasy; friable; moderately developed fine nut structure; many fine and medium roots; distinct smooth boundary,
C1	7-50	light olive grey (5Y 6/2) silt loam; friable; sticky slightly plastic; weakly developed medium block structure breaking to crumb structure; few fine roots; indistinct irregular boundary,
C2	50-67	light olive grey (5Y 6/2) silt loam; slightly firm; slightly sticky non plastic; weakly developed medium block structure breaking to crumb structure; few fine grey (10YR 6/1) basaltic lapilli; few dark brown root channels; distinct regular boundary,
Cg	67-78	pale olive (5Y 6/3) sandy loam; slightly firm; weakly developed medium block breaking to crumb structure; many indistinct yellowish brown and dark yellowish brown (10YR 5/6 & 4/6) fine and medium mottles; few medium dark brown (7.5YR 3/2) root channels; sharp smooth boundary,
2Ah	78-87	black (7.5YR 2/0) sandy loam; friable; weakly developed medium nut structure; few roots; distinct smooth boundary,
2Bw	87-96	dark brown (10YR 3/3) sandy loam; friable; weakly developed fine nut and crumb structure; few medium (2-6 mm) lapilli; locally a 4 cm pale brown (10YR 6/3) gritty sandy loam layer (E hor.) overlies this horizon; distinct irregular boundary,
2Bs	96-105	strong brown (7.5YR 4/6) pumice gravel; loose; single grain; lapilli 2-10 cm; sharp smooth boundary,

Profile	Depth (cm)	PROFILE DESCRIPTION
3Ah	105+	on yellowish brown (10YR 5/4) sandy loam; friable; weakly developed nut and crumb structure; many fine (2-4 mm) lapilli; few fine roots.

CLASSIFICATION - N.Z. Genetic: Recent soil (ash)

Taxonomy: Typic Udorthent, fine loamy, mixed, mesic

CHEMISTRY ROTOMAHANA LOAM

Sample No. SB	Depth (cm)	Hor.	pH				Exchangeable cations (meq/100 g)					Extr. Acidity (pH 8.2)	Acidity-Al (meq/100 g)	ECEC	CEC (meq/100 g)		Base saturation (%)		
			H ₂ O	KCl	ΔpH	NaF	Ca	Mg	K	Na	H				Al	NH ₄ OAc (pH 7)	Σ Cations (pH 8.2)	Σ bases CEC NH ₄ OAc	Σ bases Σ Cations
9581 A	0-7	Ah	5.7	4.7	-1.0	7.8	14.7	4.6	2.03	0.33		0.22	29.6	29.4	21.8	28.2	51.3	77	42
B	7-25	C1	6.0	4.5	-1.5	8.8	5.9	2.76	1.81	0.49		0.14	7.5	7.4	11.3	12.3	18.5	89	59
C	25-50	C1	6.7	4.8	-1.9	8.5	8.3	3.11	1.31	1.23		0.08	5.5	5.4	14.1	13.7	19.5	(100)	72
D	50-67	C2	6.9	5.0	-1.9	8.5	8.2	3.05	0.97	1.56		0.08	4.4	4.3	14.0	13.7	18.2	(100)	76
E	67-78	Cg	6.9	5.2	-1.7	8.5	10.2	3.6	1.40	2.13		0.10	5.6	5.5	17.4	17.2	22.9	(100)	76
F	78-87	2Ah	6.5	5.5	-1.0	8.8	37	6.6	0.40	0.42		0.31	27.8	27.5	44.7	46.2	72.2	96	61
G	87-96	2Bw	6.5	5.5	-1.0	10.2	11.0	2.11	0.21	0.24		0.49	16.4	15.9	13.8	15.1	30.0	90	46
H	96-105	2Bs	6.4	5.3	-1.1	10.6	7.3	1.24	0.25	0.52		0.81	17.9	17.1	9.3	12.2	27.2	76	34


Sample No. SB	Depth (cm)	Hor.	Total C (%)	Total N (%)	P (mg/100 g)			P Retention (%)	Dithion. cit. (%)		Tamm ox. (%)			Pyrophos. (%)		Reserves (meq/100 g)		Extractable S (ppm)
					H ₂ SO ₄ (0.5 M)	Inorg.	Org.		Fe	Al	Fe	Al	Si	Fe	Al	K _c	Mg _r	
9581 A	0-7	Ah	7.1	0.17	16	21	31	24	0.76	0.19	0.46	0.22	0.06	0.19	0.09	0.64	17	14
B	7-25	C1	0.6	0.12	17	23	6	17	0.77	0.19	0.34	0.14	0.10	0.05	0.01	0.73	26	3
C	25-50	C1	0.2	0.06	18	23	2	11	0.71	0.07	0.30	0.08	0.10	0.01	0.00	0.74	24	3
D	50-67	C2	0.2	0.05	18	24	1	10	0.72	0.07	0.27	0.08	0.09	0.01	0.00	0.81	22	2
E	67-78	Cg	0.3	0.06	18	24	2	13	0.99	0.14	0.35	0.10	0.09	0.03	0.00	0.72	19	7
F	78-87	2Ah	9.5	0.80	4	8	38	42	0.56	0.33	0.29	0.31	0.07	0.18	0.23	0.16	2.0	2
G	87-96	2Bw	3.7	0.41	4	11	26	45	0.58	0.37	0.32	0.49	0.04	0.25	0.31	0.12	1.1	4
H	96-105	2Bs	2.7	0.31	7	12	21	60	0.72	0.84	0.45	0.81	0.34	0.33	0.34	0.08	0.7	4

PARTICLE SIZE DISTRIBUTION (<2 mm) Rotomahana Loam

Sample No. SB	Depth (cm)	Hor.	Sand		Silt	Clay	Fine clay		Stones (%)
			2-0.1 mm (%)	0.1-0.05 mm (%)	0.05-0.002 mm (%)	<0.002 mm (%)	<0.0002 mm (%)	Fine clay Total clay (%)	
9581A	0-7	Ah	27	11	43	19			(4)
B	7-25	C1	18	10	48	24			(<1)
C	25-50	C1	16	8	46	30			(1)
D	50-67	C2	22	8	41	29			(<1)
E	67-78	Cg	18	11	42	29			
F	78-87	2Ah	25	14	45	16			2
G	87-96	2Bw	22	13	57	8			9
H	96-105	2Bs	50	11	34	5			50

PHYSICS

Hor. Depth (cm)	Hor.	15 bar Field moist (%)	water Air Dry (%)	Core Depth (cm)	Dry bulk density (T/m ³)	Total porosity (%)	Large pores (%)	Field Cap. (at 0.2 bar) (% v/v)	Wilting Pt. (at 15 bar) (% v/v)	Available water (% v/v)
0-7	Ah	25.0	16.7							
7-25	C1	14.0	12.1							
25-50	C1	14.3	12.4							
50-67	C2	14.0	12.3							
67-78	Cg	15.0	12.8							
78-87	2Ah	21.2	16.0							
87-96	2Bw	15.5	7.3							
96-105	2Bs	12.4	7.4							



Mineralogy Rotomahana

Sample No. SB	Depth (cm)	Hor.	Clay Fraction (%)										Sand Fraction (%)																		
			Mica-Smectite	Mica-Vermiculite	Vermiculite	Interlayered Hydrous Micas	Mica	Kaolinite	Halloysite	Gibbsite	Quartz	Cristobalite	Allophane	Feldspar	Anatase	Volcanic glass	Quartz	Feldspar (acid)	Andesine	Glass	Muscovite	Biotite	Hornblende	Augite	Hypersthene	Epidote	Zoisite	Apatite	Magnetite	Plant opal	
9581A	0-7	Ah			41	10		28	10			1	10																		
B	7-25	C1			40	6		26	17				11																		
C	25-50	C1			40	6		38	14				12																		
D	50-67	C2			41	4		24	16				15																		
E	67-78	Cg			56	4		14	11				10																		
F	78-87	2Ah			20			27					14			32															
G	87-96	2Bw			17			26					28			42															
H	96-105	2Bs						10								61															




Fig. 33 Relationship between vegetation, tephra age (note ages are given in ¹⁴C ka (not calibrated)) and human impact (from Watanabe and Sakagami, 1999 – after Sase and Hosono, 1996). Note dates for initial human impact vary in Japan. Initial settlement of New Zealand occurred c. 1280 AD, a few decades prior to the Kaharoa eruption.




Fig. 34 Evidence of environmental change since c. 25,000 cal years BP based on phytolith and clay mineral data from analysis of buried soil horizons on rhyolitic tephras at Te Ngae, near Rotorua (from Churchman and Lowe, 2012, after Newnham et al., 1999; based on Green, 1987; Sase et al., 1988; Hodder et al., 1990; Kondo et al., 1994).

Extracting DNA from allophanic paleosols on tephras for paleoenvironmental reconstruction: new method and application, NZ

Huang, Y-T., Lowe, D.J., Zhang, H., Cursons, R., Young, J.M., Churchman, G.J., Schipper, L.A., Rawlence, N.J., Wood, J.R., Cooper, A. 2016. A new method to extract and purify DNA from allophanic soils and paleosols, and potential for paleoenvironmental reconstruction and other applications. *Geoderma* 247, 114-125.

Soils developed from late-Quaternary tephras are commonly dominated by the nanocrystalline aluminosilicate, allophane, and are called Andisols. They contain large stores of organic matter and are potential reservoirs for DNA. However, DNA recovery from Andisols has been difficult because of strong chemical bonding between DNA and allophane and organic matter, and also because up to 80% of DNA can be physically protected in nanolabyrinthic networks of nanopores in allophane nano/microaggregates. We have developed a new two-step DNA isolation method for Andisols and allophanic paleosols, including those low in clay, which circumvents these problems. The method centres on (1) using a buffer that releases mainly microbial/fungal DNA and unbound DNA and prevents re-adsorption of DNA on allophanic materials, and (2) novel application of acidified ammonium oxalate (Tamm's reagent) to dissolve the allophane and thus release DNA which had been both chemically-bound and encased within nanopores. Sequencing of PCR products obtained from a buried allophanic paleosol at 2.2-m depth on a Holocene tephra near Mt Tarawera yielded endemic and exotic plants that differed from the European grasses growing currently on the soil's surface. This difference suggests that the DNA extraction method is able to access (paleo)environmental DNA derived from previous vegetation cover. Our method hence may be applied to Andisols and allophane-bearing paleosols, offering (1) a means to isolate paleoenvironmental DNA and thus facilitate reconstruction of past environments in volcanic landscapes (datable using tephrochronology), (2) a new way to evaluate biodiversity in such soils/paleosols, and (3) possible application to soil forensic analysis.




Fig. 35 (a) Synthetic allophane and (b, c) its spherical morphology seen using TEM (from Huang et al., 2016b)




Fig. 36 **A** Direct chemical adsorption of DNA on allophane; **B** indirect chemical adsorption of DNA on organic matter-rich allophane; **C** 'physical' adsorption of DNA (i.e. protection) in the nanopores of allophane nano- and micro-aggregates (from Huang et al., 2016a)

Extracting DNA from allophanic paleosols on tephras for palaeo-environmental reconstruction: new method and application, NZ

MARSDEN FUND
TE PŪTEA RANGAHAU
A MARSDEN

David J. Lowe, Y.-T. Huang, H. Zhang, R. Cursons, J. Young, G.J. Churchman, L.A. Schipper, N.J. Rawlence, J.R. Wood, A. Cooper

1. Introduction: (dis)solving the almost impossible

- Ancient DNA (aDNA) is usually extracted from *discrete* fossil samples of bone, teeth, eggshell, leaves, etc, and aDNA has become a **powerful tool** for Quaternary studies including (e.g.) palaeoecology, archaeology, biogeography, and human evolution and disease
- *Disseminated* environmental DNA (e.g. from ice, sediment, soil, tephra, dental calculus), defined as **palaeoenvironmental DNA (PalEnDNA)**, is also used in Quaternary studies and archaeology (e.g. Rawlence et al. 2014, *J. Quat. Sci.* 29, 610-626).
- Allophanic paleosols on tephras sequester much carbon (incl. ~80% physically protected in stable nanoaggregates), hence provide a potential reservoir for PalEnDNA. But DNA extraction from such paleosols is very difficult because of (i) strong chemical bonding between allophane and DNA, and (ii) inaccessibility of DNA encased in nanoaggregates. **Our new method bypasses these difficulties.**




Fig. 1 A Direct chemical adsorption of DNA on allophane; B indirect chemical adsorption of DNA on organic matter-rich allophane; C 'physical' adsorption of DNA (i.e. protection) in the nanopores of allophane nano- and micro-aggregates

2. Objectives

- Develop an effective method for DNA extraction from (i) synthetic allophane and (ii) natural allophanic soils or paleosols
- Search for possible PalEnDNA from buried allophanic paleosols on Holocene tephras




Fig. 2 (a) Synthetic allophane and (b, c) its spherical morphology seen using TEM

3. Results

- Synthetic allophane and salmon-sperm DNA provided proof-of-principle before trialling natural allophane; humic acids add complexity
- Our two-step method includes novel use of ammonium oxalate to *dissolve* allophane to release **all DNA including that entrapped in nanoaggregates** (unlike previous methods):
Step 1 releases microbial & unbound DNA (likely modern);
Step 2 releases chemically bound and physically adsorbed DNA (likely to be pre-modern, potentially aDNA)

3. Results contd.




Fig. 3 Summary of two-step DNA extraction from natural allophanic soils/paleosols




Fig. 4 A DNA extracted from paleosols on Taupo (TP), Whakatane (WK), Rotoma (RM) tephras; B amplifiable DNA fragments from Rotoma paleosol (Table 1) with bacterial, animal, and plant primers: 16s, bacterial; Lep, insect; rbcl, *trnL*, plant



Fig. 5 Sampling for possible PalEnDNA in buried allophanic paleosols on tephras deposited since ~12,000 yr BP



Fig. 6 Paleosols on dated tephras near Mt Tarawera

Modern plants at land surface = European grasses
Such vegetation at this site dates to c. 1900 AD or earlier

Table 1 New Zealand endemic plants detected in buried soil on Rotoma tephra using our extraction/purification protocols & polymerase chain reaction (PCR) with selected primers

Family	Plant taxa	Primer pair and % similarity to reference
Myrtaceae	<i>Leptospermum scoparium</i>	<i>trnL</i> g-h (98-99%) and <i>rbcl</i> (99%)
Araliaceae	<i>Hydrocotyle</i>	<i>rbcl</i> (97%)
	Other genera (multiple)	<i>trnL</i> c-h (99%) and <i>trnL</i> g-h (100%)
Griselinaceae	<i>Griselinia lucida</i>	<i>rbcl</i> (98%)
Podocarpaceae	<i>Podocarpus</i>	<i>trnL</i> c-h (95-98%)

(Huang et al. 2016 *Geoderma* 274, 114-125)

4. Conclusions

- Our two-step method enables previously inaccessible DNA to be extracted from allophanic soils by releasing DNA entrapped in stable nano/microaggregates
- Gel electrophoresis of extracted DNA, then gel purification, removes sufficient humic material for successful DNA amplification via PCR of multiple gene regions
- DNA extracted from a buried paleosol on Rotoma tephra (~9.4 cal ka) includes native NZ forest spp., different from European adventives on the land surface
- Thus DNA extraction from allophanic paleosols potentially offers a means to isolate PalEnDNA and hence facilitate reconstruction of past environments extensively in volcanic landscapes (datable using tephrochronology)
- Other potential applications include studies on soil biodiversity and soil forensics

Potential impacts of ash fall on dairy farms – Rerewhakaaitu case study by Wilson and Cole (2007)

Nat Hazards (2007) 43:103–128
DOI 10.1007/s11069-007-9111-8

ORIGINAL PAPER

Potential impact of ash eruptions on dairy farms from a study of the effects on a farm in eastern Bay of Plenty, New Zealand; implications for hazard mitigation

Thomas M. Wilson · James W. Cole

Abstract This paper investigates the impact ash fall would have on dairy farming, based on a study of ‘Tulachard’, a dairy farming operation at Rerewhakaaitu, North Island, New Zealand. It includes analysis of the potential effects on the dairy shed and milking machine, electrical supply and distribution, water supply and distribution, tractors and other farm vehicles, farm buildings (haysheds, pump sheds, implement sheds, etc.), milk-tanker access to the farm and critical needs of dairy cows and farm to keep milking. One of the most vulnerable areas identified in the study was the cooling of milk at the milking shed, pending dairy tanker pick-up. The cooling system’s condenser is exposed to the atmosphere and falling ash would make it highly vulnerable. Laboratory testing with wet and dry ash was conducted to determine its resilience to ash ingestion. It was found to perform satisfactorily during dry testing, but during wet testing significant clogging/blocking of the condenser’s radiator occurred, dramatically reducing airflow through the condenser. Specific mitigation recommendations have been developed that include cleaning with compressed air and adapting farm management techniques to lessen usage of the condenser during an ash-fall event. Specific recommendations for management of dairy farm operation are given to mitigate the effects of an ash-fall event.




Fig. 2 The annual farm activities at the farm shows the key vulnerable periods during the year of farming operations. The farm’s most important activities are closest to the centre, with activities of decreasing importance progressively further out from the centre

7 Recommendations

The effect of ash fall will depend on the magnitude, style of eruption, location of active vent zone, vent positions within the active vent zone, local topographic controls and wind directions. It is extremely hard to predict any of these factors for the next eruption, given the variety of styles of eruptions that have affected the farm during the last 22,000 years. Warning time is variable. There may be 1 year to 3 months warning for a rhyolitic eruption, while there may be only 3–4 h for a basaltic eruption.

Key recommendations to minimise the hazard are as follows:

- (1) Long term planning (*periods of quiescence*)
 - Maintain power supply lines so they are in good order.
 - Develop a water supply with large tank storage capacity.
 - Develop a 'feed pad' that is easily cleared of ash for distributing supplementary feed.
 - Ensure roof pitches are greater than 30° and strengthened (perhaps to snow codes) on any new buildings.
 - Ensure the tractor is 4WD and has front bucket and blade attachments.
- (2) Medium term planning (*12 to 3 months out from eruption*)
 - Conduct a vulnerability analysis of equipment and facilities to determine which would be the most affected by ash fall, and which are adequately and inadequately protected (<http://volcanoes.usgs.gov>).
 - Identify appropriate methods of protecting vulnerable equipment and facilities from ash.
 - Off load non-essential stock (i.e. beef cows, sick animals in the herd, dry milkers, old cows).
 - Increase reserves of supplementary feed (i.e. buy in or produce more; alternatively use less by off loading stock).
 - Ensure tractor and milking machine have been serviced recently.
 - Attempt to stockpile tractor engine and milking machine filters (air, oil, pulsators and milk), lubricating oil, brake and hydraulic fluids, and seals.
 - Purchase an air compressor or ensure it is in good working order.
 - Ensure diesel tank is maintained at a high level.
 - Have ladders and brooms (roof cleaning).
 - Ensure sump, drainpipes and drain grills are clear. Make sure sump pump is in good working condition.
 - Be prepared for false alarms; predicting a volcanic eruption is difficult.
- (3) Short term planning (*immediately before an eruption*)
 - Cover all essential equipment (either within sheds or under a covering).
 - Move essential stock (milking herd) close to the milking shed.
 - Store transportable supplementary feed close to areas where it would be distributed (i.e. feed pad, paddocks close to the milking shed).
 - Ensure enclosed water storage tanks are at their maximum (especially if surface water is the farm's water supply).
 - Place ladders for access to key roofs in a secure way for easy safe cleaning. Plan to have a lot of time to clean roofs (prevent injury).
 - *Do not put stock onto the road* in the hope of finding somewhere better as they will hinder emergency service's and evacuee's mobility. They have a greater chance of survival on their farm (Neild et al. 1998).
- Conduct any maintenance on tractor, milking machine and other key machinery (e.g. change filters). If there is time purchase new filters.
- (4) During eruption
 - Keep ash out of buildings, machinery, vehicles, downspouts, water supplies and wastewater systems (e.g. dairy shed drains) as much as possible. The best way to prevent damage is to reduce machinery usage as much as possible, shutting down, closing off or sealing equipment. However, critical farm functions such as milking should be preceded by removal of as much ash as possible before operating equipment.
 - Minimize human exposure to airborne ash by using dust or filter masks and minimising travel.
 - Remove ash from roofs to prevent collapse and on going remobilisation. Take your time whilst cleaning roofs and wear a mask whilst doing so. Try to undertake the activity when you are alert and physically fresh.
 - Plan each day, in terms of what activities are critical, required and optional. These may change each day so be prepared to remain flexible. Develop a priority list of facilities that must be kept operative versus those that can be shut-down during and after ash falls.
 - Stay aware of the condition of your cows; they are the most important part of your farm.
 - Do not put stock onto the road in the hope of finding somewhere better as they will hinder emergency service's and evacuee's mobility. They have a greater chance of survival left on your farm.
 - Prioritize and sequence areas for cleanup (top to bottom).

TEPHRAS AND BURIED SOIL SEQUENCE, MATAHINA GRAVEL, ASH PIT RD

Location V16 188155, 38° 18.568 S, 176° 31.817 E; elevation ~448 m




Fig. 37 Matahina gravel and associated buried soil horizons and tephra layers at Ash Pit Rd section. Ages from Lowe et al. (2013). Note the clear spodic-podzol-profile morphology (distinct E horizon on a Bh horizon, referred to as a spodic or podzolic-B horizon) evident on Taupo tephra, and another weak E horizon beneath Taupo on Whakatane tephra in this photo and one on page above. A blotchy E horizon also occurs beneath Whakatane tephra on Rotoma tephra. A similar sequence occurs at a nearby farm, 'Ashton Dairies', which shows three distinct podzol-profile morphologies (see Fig. 38). The grey 'E' horizons, and associated underlying Bh or Bs horizons, are pedogenic in origin and can easily be misidentified as separate pale tephra layers on buried soils (i.e., 'geology meets pedology') (Lowe et al., 2012).

Note from Wim Rijkse

Matahina gravel is a moderately leached soil with base saturations fluctuating from 71% in Tarawera Tephra, to 51% in the paleosol on Kaharoa Tephra, and 19% in Kaharoa lapilli, to finally 67% in the paleosol on Taupo Tephra ((Vucetich et al., 1960). Exchangeable cations are of low values except sodium which is medium. Available phosphorus presumably is low.

NZSC: Allophanic Orthic Pumice; Mt; na (Ba and Rh fines/stones); loamy/sandy; m

Soil Taxonomy: Ash-skeletal/pumiceous, mixed/glassy, subactive(?), mesic Vitrandic Udorthents




Fig. 38 Summary stratigraphy in the Lake Rerewhakaaitu area at Brett Rd, Ashton Dairies pit, and Rerehakaaitu Rd sections, and a fourth sequence at Maniatutu Rd near Lake Rotoiti (from Lowe et al., 2012, based on Marsden-funded research “New views from old soils”).

Military activities in the Rerewhakaaitu area

Military activities around Rerewhakaaitu, documented by Mossop (1993), began with annual camps by New Zealand field artillery units in the 1920s-1930s. With the onset of World War II, the artillery units became established along the eastern shores of Lake Rerewhakaaitu and near Mt Tarawera. Batteries fired from escarpments along Northern Boundary Road into the lower slopes of Mt Tarawera. The New Zealand Tank Brigade (~1000 men and ~300 vehicles) trained there from May 1943. Three infantry divisions of the US Army (the 25th, 37th, and 43rd), comprising about 15,000 men in total, camped and trained in the area from 1942 to July 1944. The main units were the 43rd 'Winged Victory Division' and to a lesser extent the 25th 'Tropic Lightning Division'. The 25th Division was from Hawaii. It left New Zealand in February 1944. The 43rd Division was raised from Vermont, Connecticut, and Maine. It began training at Rerewhakaaitu on April 1944, and left New Zealand by the end of July. Winter in 1944 was very cold and snowfalls were recorded.

The main training area comprised land to the northeast of Lake Rerewhakaaitu bounded by Mt Tarawera in the north and the edge of the Kaingaroa forest in the south. The only other people living in the district were on Brett Rd, west of the lake, according to Mossop (1993). The main line of fire was towards the east, the so-called 'impact area' bounded today by eastern Ash Pit Rd. The camps were entirely tent-based with no wooden buildings. After the war ended, engineers from the New Zealand Army began the job of disposing of unexploded shells etc, completed by about 1950.

From the 1950s, the area on the northern shores of the lake was used as a camp for compulsory military training by the New Zealand Army (Steve Gilbert pers. comm., 2006). Gilbert also recorded that the road to the top of Mt Tarawera had been constructed by the US Army and lookouts posted there permanently during the war. Today, reminders of the USA presence in the area include Yankee Rd, Republican Rd, and Democrat Rd, although the last has been re-named Rerewhakaaitu Rd on the latest topographic maps. A trig point near the intersection of Ngamotu Rd and Okahu Rd is recorded as 'America NL' (506 m).



Captain Smith (left) and his cannoneers examine the effects of close range fire on a tree in the Rerewhakaaitu area.

Fig. 39 From Griffiths and Schuler (1993)



Fig. 40 Rotorua city and Lake Rotorua inside Rotorua caldera. Multiple volcanic landforms are present.




Fig. 41 (Top) Okataina Volcanic Centre, including the Okataina (or Haroharo) caldera complex, dome complexes and known volcanic vents (after Nairn, 2002; Cole et al., 2010, 2014). **Fig. 42 (Bottom)** Rhyolite domes associated with the Okareka Embayment and their relationship to stops labelled 5 and 5. Lake Tikitapu ('Blue Lake') lies directly to the south.

STOP – RNL PUMICE QUARRY: NEAR-VENT PUMICEOUS TEPHRA

Building the history of eruptions and landscape evolution

Sequence is as follows (Fig. 44 below):

- early tephtras (Te Rere, 25.1 ka; Okareka, 21.8 ka) and thin interbedded loess within the remnant buried hill at the base of the sequence;
- the buried hill itself, which represents a period of erosion that occurred during the last glacial period, and remnants of a (reworked?) tephra draping the hillslope (Rerewhakaaitu, 17.6 ka);
- the thick, bedded pumice lapilli tephra mantling the hill (Rotorua, 15.6 ka), the top of which is modified to form a distinct orange-coloured soil; and
- younger tephtras and soil horizons overlying the buried soil on the Rotorua tephra.

From distal to proximal – getting closer to the source

The thick Rotorua tephra is particularly distinctive here. We have been tracking this tephra all the way from Hamilton to this site, only 4.5 km from its source vent(s). The source vent(s) lie to the southeast beneath the Trig7693 and Middle domes of the Okareka Embayment. We can see from the isopach map (Fig. 43) how the Rotorua tephra thickens closer to the vent(s), and also that much of it was dispersed towards the northwest. The tephra deposits to the northwest represent deposition from an early explosive phase of the Rotorua eruption sequence (phase 1 on Fig. 43) involving a ~20-km-high plinian eruption plume that must have lasted for at least 3.5 hours, but no more than 3-4 days (Kilgour and Smith, 2008). The second phase of the eruption sequence, which lasted for 3-6 years, involved the growth of the Trig7693 and Middle domes and intermittent explosive eruptions that seem to have generated thin tephra deposits blown to the southeast (denoted as phase 2 on Fig. 43).

The Te Rere tephra also comes from nearby source vent(s) in the Okareka Embayment beneath the Northern and Eastern domes. Most of the remaining tephtras seen here are from the Tarawera Volcanic Complex, 15 km to the southeast, except the Rotoma tephra, which was derived from the Rotoma Rhyolite Complex, 25 km to the northeast. The youngest tephra is Rotomahana Mud, deposited during the Tarawera eruption on 10 June 1886, at the land surface.



Fig. 43 Isopach map for the Rotorua tephra (modified from Kilgour and Smith, 2008; Shane et al., 2003) with thicknesses in centimetres. Compositional data are available in Smith et al. (2004).



Fig. 44 Outcrop on Okareka Loop Rd by entrance to RNL Pumice Quarry (stratigraphy partly after Nairn, 1992). Ages on tephras from Lowe et al. (2013). 'Cal yr BP' refers to calendar (calibrated) years before present, 'present' taken as 1950 in radiocarbon-based timescale.

STOP – KAINUI SILT LOAM & NAIKE CLAY [NB: DATA BELOW AT GORDONTON RD SECTION]

Location S14 133859, elevation ~60 m asl, rainfall 1200 mm pa

At this stop are several remarkable features both stratigraphic and pedological, and a ‘two-storied’ soil, the Kainui silt loam alongside (in just a few places) the Naike clay. Both soils are Ultisols. The sequence of tephra beds and buried soil horizons spanning about 1 million years was exposed in 2007 by road works. We will not see this site; rather, an equivalent site on Raynes Rd south of Hamilton (Kainui soil).




Fig. 45 Provisional stratigraphy of Gordonton Rd section. Horization based on Clayden and Hewitt (1989). The Kainui soil is entirely halloysitic because of limited desilication. *Photos:* left, David Lowe; right, Richard McEwan.

Stratigraphy

(1) At the base of the section is the tiny remnant of an ancient landsurface represented by a buried, clay-rich soil >0.78 Ma on the basis of its reversed magnetism and other evidence (Horrocks, 2000; Lowe et al., 2001). Several undifferentiated beds visible beneath it are likely to be c. 1 Ma or older. These old tephra beds at the base of the Gordonton Rd section (northern end) are part of the so-called Kauroa ash bed sequence, the uppermost unit being the Waiterimu ash member or K15 (Ward, 1967; Lowe and Percival, 1993; Lowe et al., 2001).

(2) Overlying K15, with an unconformity, is the white ~0.5-m thick Rangitawa Tephra aged 0.35 Ma, one of the most widespread tephtras in the southwest Pacific region that fell during Marine Oxygen Isotope Stage 10 (OIS 10). Rangitawa Tephra is overlain in turn by a ~3-m thick sequence of weathered, yellowish brown to brown to reddish brown clayey tephra beds and buried soils – the Hamilton ash beds – that represent OIS stages 9 to 5. The uppermost distinctive, dark reddish-brown buried soil, known as the Tikotiko ash member or bed H6/7 (Ward, 1967; see below), represents probably the Last Interglacial (OIS 5) paleo-landsurface (probably equivalent to the Sangamon paleosol/geosol of North America) (Lowe, 2019).

(3) At the top is a thin, silty cover bed mantle of intermixed late Quaternary tephtras about 0.4-0.6 m thick, occasionally overthickened in hollows up to ~1.2 m in thickness, that have accumulated incrementally over the past c. 50,000 years (OIS 3–1). The base of the cover bed has been identified as Rotoehu Ash (c. 50 cal ka)

but it is evident from detailed mineralogical and geochemical studies that the component tephras making up the composite mantle have been strongly intermixed during upbuilding pedogenesis (Lowe, 1981, 1986, 2000). Nearby lakes, formed c. 20 cal ka following deposition of the volcanogenic alluvium of the Hinuera Formation, have preserved >40 multiple, thin, visible tephra layers within their sediments (e.g. Green and Lowe, 1985; Lowe, 1988). These tephras each range in thickness from a few millimetres to several centimetres and in this area amount to an estimated ~37 cm in total thickness (Lowe, 1988). Together with tephras deposited between c. 50 and 20 cal ka (Kawakawa Tephra, aged c. 25.4 ka; Hauparu Tephra, c. 36 ka; Tahuna Tephra, c. 39 ka; and Rotoehu Ash, c. 50 ka) they form the parent material of the upper part of the Kainui soil (Lowe, 1986, 2019). The thin tephras preserved in the lake sediments are both rhyolitic (predominant) and andesitic and were derived from five volcanic centres 70 to 200 km away (Lowe, 1988). There are likely to be numerous very thin (< 1 mm) 'non-visible' tephras or sparse concentration zones of glass shards, known as cryptotephras (Alloway et al., 2007a; Lowe, 2011), in the lake sediments as well as the visible layers. From recent work on lake cores and peat bogs, such cryptotephras are confirmed in the Waikato region (Gehrels et al., 2006, 2008; Loame et al., 2018) and thus probably were assimilated into the Kainui soil as 'dustings' from small-scale eruption plumes.




(Above)

Fig. 46 Lake Maratoto, south of Hamilton, was formed c. 20 cal ka. It contains the parastratotype for the base of the Holocene for Australasia (marked by Konini-bed b tephra) (Walker et al., 2009). *Photo:* David Lowe

(Above right)

Fig. 47 The first core of 33 eventually taken from the lake (April 1979). Grey layer near base is alluvium overlying dark proto-lake lake sediment (near tape head) and pre-lake soil (Green and Lowe, 1985). Part core at right (from Lake Rotongata) shows four visible tephra layers preserved within dark organic-rich lake sediment






Fig. 48 (Left) Central Waikato area showing locations of lakes cored to obtain detailed post-c. 20 cal ka tephra record. **(Right)** Cores from Lake Rotomanuka sliced open to show tephra layers preserved in dark lake sediments (from Lowe, 1988). We have recently re-cored three lakes (Kainui, Rotokauri, and Kainui for studies on cryptotephra (funded by EQC) and tephra 'seismites', i.e., tephra layers that have been liquefied by earthquake activity in the past 20,000 years.

Lake sites are: 1, L. Maratoto; 2, L. Rotomanuka; 3, L. Ngaroto; 4, L. Mangakaware; 5, L. Mangahia; 6, L. Rotoroa; 7, L. Rotokauri (thickness measurements do not include Rerewhakaaitu Ash); 8, L. Kainui; 9, L. Rotokaraka; 10, Leeson's Pond; 11, L. Okoroire; 12, L. Rotongata.

Fig. 49 Thickness and compositional relationships based on measurements of tephra in lake cores. Bars with hatching = total thickness of all tephra \leq c. 20 cal ka; blank bars = rhyolitic tephra; stippled bars = andesitic tephra. Solid bars = actual measurements, dashed bars = thicknesses corrected for compaction and dissemination in sediment cores. Dashed lines = isopachs (in mm) of sum of tephra at subaerial sites deposited since c. 20 cal ka (from Lowe, 1988).




Table 7 Total thicknesses of tephras < c. 20 cal ka in central Waikato area based on lake core measurements and estimated 'dry-land' (subaerial) thicknesses (values in centimetres) (from Lowe, 1988)

Area	Total visible thickness in cores (average)	Compaction - corrected thickness (visible x1.75)	Estimated dissemination thickness (visible x0.1)	Total equivalent dry-land thickness (approx.)
Hamilton–Ohaupo	25	44	2.5	47
Whitikahu–Morrinsville	20	35	2.0	37
Okoroire–Tirau	42	74	4.2	78

* Assumed to represent airfall material only, with no modification to thickness by postdepositional reworking or catchment erosion.





Figure 10.4 Transect across the Hamilton Basin showing the simplified stratigraphy and age of late Quaternary tephra deposits overlying Hamilton Ash on the low hills. Marker beds are distinguishable in the field in the south (Otorohanga and Te Awamutu-Ohaupo profiles) but around Hamilton the tephra deposits, thinning away from their sources, become intermixed to form a composite cover bed with no distinguishable marker beds. Where the cover bed is eroded, exhumed Hamilton Ash occurs at the surface, as for the Huntly profile, but thin late Quaternary tephras are present further north towards Auckland. The inset shows the stratigraphy of prominent tephra layers preserved in lake sediments in a core from Lake Maratoto a 17 000-year-old peaty lake about midway between Ohaupo and Hamilton (see Figure 10.5). It indicates potential contributions of specific tephras as probable parent materials in the upper soil profile (hatching). ka = 1000 years BP. After Lowe (1986).

Fig. 50 (Left) Otorohanga and Ohaupo soils (Hapludands) occur at sites marked by columns 1 and 2, Kainui soils (Udults) are at the site marked by columns 3-5 ('Hamilton' ≅ Gordonton Rd); and Naikie soils (Humults) at site marked by column 6 ('Huntly') (from Selby and Lowe, 1992). Ages in ¹⁴C yr ka. **(Below)** Otorohanga soil, Kakepuku Rd.



Description and data for Kainui silt loam

Hauraki Plains survey (by Malcolm McLeod and Brian Daly)

LAB. Nos. SB9801A-F
 Field Nos. HC12 a-f
 Est./Funct. Mod./Var.

SOIL SET: _____
 CLASSIFICATION (in full)—Common: _____
 Technical: _____
 SURVEY: Hauraki Plains County Collector: MM BD Date: 9-8-82 Pit/Comp. (___ cores)
 LOCATION—County: Hauraki Plains Survey District: H. P. Block: _____ Cult'd/Uncult'd/Virgin
 Orientation: _____ Section: _____ Topdr./Untopdr.
 Exact Location: Top NE corner of paddock Farmer: _____
 Grid Ref. NZMS 1 N52 900052
 Altitude: _____ ft (30 m) Rainfall: 1200 - 1400 mm Raindays (>0.01"): _____ (>0.05"): _____ Mean Annual Temp.: 12.8°C (57°F)

Field No.	Lab. No.	Horiz.	Depth in cm	DESCRIPTION—Colour, Texture, Consistence, Structure, Boundary, etc.
HC12 a	9801 A	Ap	0-16	10YR 3/2 with some 10YR 4/2 worm mixing from below; silt loam; moderately weak; brittle; few fine pores; weak fine root breaking to crumb; many fine roots; distinct irregular boundary.
HC12 b	9801 B	Ew1	16-32	10YR 5/3 silt loam; very weak; brittle; many medium pores; weak fine root breaking to crumb; few fine roots; indistinct wavy boundary.
HC12 c	C	Ew2	32-47	10YR 5/4 clay loam; moderately weak; brittle; many medium & coarse pores; weak fine root structure; few fine roots; indistinct wavy boundary.
HC12 d	D	2bBt(g)	47-86	7.5YR 4/6 with horizontal veins 7.5YR 6/2 plus organic matter down cracks 7.5YR 4/2; clay; moderately firm; brittle; very few medium pores; abundant clayshims; moderately developed medium & coarse blocky breaking to fine blocky; few fine roots down cracks; distinct wavy boundary.
HC12 e	9801 3bBt(f)	E	86-108	7.5YR 4/6 with few medium distinct 10YR 5/8 mottles; clay; brittle; moderately weak; few medium pores; very firm in situ; weak medium blocky; many clayshims; no roots.
	F		@ 180	

DJL note * Colour criteria qualify these upper subsoil horizons as E horizons (Clayden and Hewitt, 1989) but note that there is a lithological discontinuity at 47 cm depth where 2bBt(g) horizon is recognised. McLeod (1992) suggested that the upper subsoil horizons = eluvial horizon.

SITE—Slope: 6 Nature of Slope: _____
 Aspect: SE Exposure: _____
 Microfeatures: _____
 Erosion: Slight creep.
 Disturbance: _____
 Drainage: Moderately well drained

VEGETATION (over sample)—Height: 10cm Main Species: Pasture grasses
 (at site)—Height: 10cm Main Species: _____

History: _____ Type of Vegetation: _____

Root Distribution: _____
 PARENT MATERIAL: 47 cm moderately gravelled late Pleistocene rhyolitic and andesitic volcanic ash over strongly argillised Pleistocene rhyolitic and andesitic volcanic ash.
 Parent Rock: _____

Classification Kainui silt loam

NZSC: Buried-granular Yellow Ultic Soils; tephric; na (mixed Rh >> An fines); silty/clayey; moderate/slow
Soil Taxonomy: Fine-silty over clayey, halloysitic, thermic Typic Kandiodults

The 'organic matter down cracks' noted for 2bBt(g) in above description is likely to be MnO₂ (pyrolusite) coatings, abundant in lower subsoils (near boundary upper/lower parts of profile) of many Kainui soils. A 'Buried-granular' subgroup, rather than Podzolic subgroup, was adopted in 2010 to encompass widespread Kainui soils (Hewitt, 2010).

Moanatuatua bog and the ‘wet-desert’ paradox

The Moanatuatua peat bog is a large raised, restiad (jointed rush) bog up to 12 m deep. It began forming from c. 15,000–13,000 cal. yr BP when regional water tables rose, probably in response to moistening climate (Green and Lowe 1985; Jara et al., 2017).

‘Bogs’ have raised surfaces on a peat substrate; they are entirely rainwater fed, and have low nutrient levels (e.g. Moanatuatua, Rukuhia, Kopouatai bogs).

‘Fens’ have a flattish or slightly depressed surface on a peat and mineral substrate; they are part groundwater/runoff and part rain fed, and have medium nutrient levels (e.g. Whangamarino fen).

‘Swamps’ are groundwater/run-off fed with a mainly mineral (non peat) substrate; they are high in nutrients (Charman 2002; Clarkson 2002).

Note the ‘unusual’ microclimatology of this and other bogs in the Waikato region. These studies show that evaporation rates from *Sporodanthus*-dominated raised bogs such as Moanatuatua and Kopouatai (Hauraki) are 2.74 and 3.01 mm/day, respectively (about twice the rate for *Empodisma*-dominated canopies). These two groups of plants have highly-conservative evaporation regimes that help the bogs maintain wet conditions, hence effects of drought and artificial drainage are lessened. Their nearest relatives occur in dry deserts! The species on this and other northern North Is. bogs was re-identified as *Sporodanthus ferrugineus* (de Lange et al. 1999), a late successional species on raised peat bogs. Within these ecosystems, it is eliminated by fire and re-establishment is from seed subsequently deposited by remnant plants (it takes 5-12 years following fire to re-establish itself fully within bog vegetation) (de Lange et al. 1999). *Empodisma robustum* is also a newly recognized species (Wagstaff and Clarkson, 2012; Clarkson et al. 1999, 2004).

Peat subsidence

Rates of subsidence of three Waikato peats were examined at 119 sites (Pronger et al. 2014). They found that drainage and conversion of peatlands to productive agro-ecosystems has led to ongoing surface subsidence because of densification (shrinkage and consolidation) and oxidation of peat substrates. The average contemporary (2000s–2012) subsidence rate for Waikato peatlands was $19 \pm 2 \text{ mm yr}^{-1}$ ($\pm \text{SE}$), significantly less ($p = 0.01$) than the historic rate of $26 \pm 1 \text{ mm yr}^{-1}$ between the 1920s and 2000s. A reduction in the rate of subsidence through time was attributed to the transition from rapid initial consolidation and shrinkage to slower, long-term, ongoing oxidation. These subsidence rates agree with a literature synthesis of temperate zone subsidence rates reported for similar lengths of time since drainage. A strong nonlinear relationship was found between temperate zone subsidence rates and time since initial peatland drainage: Subsidence (mm yr^{-1}) = $226 \times (\text{years since drained})^{-0.59}$ ($R^2 = 0.88$). This relationship suggests that time since drainage exerts strong control over the rate of peatland subsidence and that ongoing peatland subsidence rates are likely to gradually decline with time in the absence of major land disturbance (Pronger et al. 2014).

Bogs as carbon sinks (Goodrich et al. 2017)

The net ecosystem carbon balance (NECB) was estimated over four years, which included two drought summers, a relatively wet summer, and a meteorologically average summer. In all measurement years, the bog was a substantial sink for carbon, ranging from 134.7 to 216.9 $\text{gC.m}^{-2} \text{ yr}^{-1}$, owing to the large annual net ecosystem production (161.8 to 244.9 $\text{gCO}_2\text{-Cm}^{-2} \text{ yr}^{-1}$). Annual methane fluxes were large relative to most Northern Hemisphere peatlands (14.2 to 21.9 $\text{gCH}_4\text{-Cm}^{-2} \text{ yr}^{-1}$), although summer and autumn emissions were highly sensitive to dry conditions, leading to very predictable seasonality according to water table position.

Note: peatlands make up ~2 to 3% of the Earth’s surface and store a large proportion of soil carbon (~50%) (Goodrich et al. 2017). CH_4 (methane) is a very potent greenhouse gas.

STOP RUKUHIA PEAT, JARY ROAD

Location BE34 080974 Elevation ~ 50 m asl Rainfall ~1400 mm pa

Examine **Rukuhia peat**.

NZSC: *Acid Mesic Organic Soil*

Soil Taxonomy: *Either Sapric or Typic Haplohemists*

Of = fibrous peat or 'fibric soil material' – contains mainly well-preserved plant remains (contains $\geq 75\%$ fibre after rubbing)

Om = semi-fibrous peat or 'mesic soil material' ('hemic soil material' in *Soil Taxonomy*) – contains mainly partially decomposed plant remains

Oh = humified peat or 'sapric soil material' (Gk, *sapros*, rotten) – contains strongly decomposed organic materials remains (contains $< 15\%$ fibre after rubbing).

Where organic surface horizons have been mixed by cultivation, the suffix 'p' is added, e.g. Ofp, Omp, Ohp (Clayden & Hewitt pp.7-8).

von Post scale

Peat may also be described on a 10-point 'von Post scale' (from pioneer Swedish palynologist von Post) used for assessing the degree of decomposition of organic materials. In the field a sample of wet organic material (about the size of small egg) is squeezed in the closed hand and the colour of the liquid that is expressed between the fingers, the proportion of the original sample that is extruded, and the nature of the plant residues are observed (classes H1 to H10) (Milne et al. 1995, p. 54-55; see also McLay et al., 1992):

H1-H3 = 'fibric', squeezed water mainly clear or slightly coloured;

H4-H7 = 'mesic' or 'hemic', squeezed water brown-dk brown, & most peat oozed between fingers;

H8-H10 = 'sapric', no water squeezable, peat oozed between fingers

Organic Soils (Histosols) always key out separately from all other soil orders

Organic Soils have many properties that differ from 'mineral' soils (i.e. soils of the other orders dominated by inorganic material). For example, bulk densities in Organic Soils normally range from ~ 0.03 to 0.4 Mg/m^3 (Hewitt, 2010). Although the moisture content of peat is normally very high, a large proportion of it may be unavailable to plants because it is hydrophobically sorbed by the organic particles at suctions above wilting point. Consequently, strict water table control is crucial to maintain the amount of water available for plant growth (McLay et al., 1992).

Tephra

Note the layer of fine lapilli representing Taupo Tephra at about 0.5 m depth (probably Taupo Plinian, subunit Y5; erupted 232 ± 10 AD) (Hogg et al. 2012), or possibly co-ignimbrite lapilli that fell during and after the ignimbrite phase emplacement (subunit Y6). Other tephra-fall layers have been identified at depth in the peat, including Mamaku, Rotoma, Waiohau, and Rotorua tephra (Lowe 1988; Newnham et al. 1995; Jara et al., 2015). Kaharoa Tephra (1314 ± 12 AD; Hogg et al. 2003) and andesitic material are also present as *cryptotephra* (Gk, *kryptein*, to hide) ('hidden' glass-shard and/or crystal concentrations not sufficiently numerous, or too disseminated, to be visible as a layer in the field: Lowe 2011) near the top of the peat (Gehrels et al., 2006, 2008, 2010).

Blueberries

The Organic Soils here have been developed to grow blueberries (*Vaccinium corymbosum* or *V. ashei*) which favour low pHs (around 4.3 at this site). The bushes were planted in 1982 and include about 13 different varieties. Picking is from mid-December to early April. Much fruit is exported. Note problems and benefits in managing these Organic Soils (McLay et al. 1992; Clarkson et al. 1999).

Main landscape units and geological materials, Hamilton Basin





Fig. 51. Generalised geology-landform relationship (upper) and associated soil pattern (lower) in the Hamilton Basin

General landscape pattern

The Hamilton area is characterised by four main landscape units or landforms:

- Low rolling hills (e.g. University, Hospital, Hamilton East Cemetary, Wintec city campus) – the ‘Hamilton hills’
- Flat alluvial plains (e.g. much Hamilton East) with micro-relief of low mounds (bars) and swales (depressions)
- Low terraces adjacent to the modern Waikato River (e.g. Lower Hamilton Gardens, Memorial Park)
- Gullies cut into the alluvial plain or low terraces (e.g. gully running through Hungerford Cresc into the Gardens)

1. The low rolling hills represent the remnants of a landscape dating back around more than a million years. A drill hole through a hill will typically show the following sequence of deposits (from top down):

- Silty cover bed of post-Hamilton-Ash tephra (volcanic ash) beds from multiple sources; ~0.5 m thick; ≤c. 50 ka
- Red-brown, clayey weathered tephra beds (Hamilton Ash); ~1–3 m thick; top bed c. 50–100 ka, basal c. 340 ka (dark red-brown uppermost buried soil horizon probably represents soil formation during the last interglacial)
- Orange/reddish/cream gravelly alluvial clays (Karapiro Formation); variable thickness (few metres); c. 500 ka
- Very dark reddish brown, clayey weathered tephra beds (part of Kauroa Ash Formation); patchy; c. 1 Ma
- Cream-coloured ignimbrite (deposits from hot, ground-hugging, fast moving pyroclastic flows, also called pyroclastic density currents); up to 10–20 m thick. Three main units: Ongatiti Ig., Rocky Hill Ig., Kidnappers Ig., aged from c. 1.2 to 0.9 Ma, respectively. The sculptures at Gate 1 (by Hungerford Cres/Cobham Drive) are made of Ongatiti Ignimbrite (c. 1.23 Ma) which is quarried as 'Hinuera Stone' in the Hinuera Valley between Piarere and Hinuera (near Matamata).

The earliest of these deposits, the ignimbrites, usually form the 'core' of the Hamilton hills. They and associated alluvium have been partly eroded to form a rolling landscape which has then been blanketed with subsequent tephra beds draping the hills and the valleys. So, the underlying hilly terrain is actually quite an old remnant landscape feature, one of the oldest in New Zealand. Buried soils or soil horizons are common and represent landscape surfaces of the past. The modern soils are developed largely in the uppermost deposits, namely the multiple tephra-derived cover bed and upper Hamilton Ash beds: the main soils, typically sticky and clayey, are referred to as the Kainui and Hamilton series (Kn + Hm). Hamilton soils tend to occur more commonly on shoulder slopes and are less common than soils of the Kainui series.

2. The plains represent alluvium derived ultimately from the mainly volcanic catchments of the central North Island and deposited by ancestral Waipa and then ancestral Waikato river system in a series of depositional episodes over the past c. 100 ka or so. These deposits swept around and over the pre-existing hilly landscape in the Waikato, partly burying it so that today we find just remnants of the hills protruding through the essentially flat-lying alluvial surface (the surface actually has slight surface features, a series of low ridges/bars and swales or depressions; it also slopes very gently in a fan form, the apex at Maungatautari and the toe at Taupiri, ~1 m vertically for every 1 km horizontally). The ancestral Waikato River was predominantly a high energy, braided system that until around 22,000 calendar (cal.) years ago flowed through the Hauraki Plains via the Hinuera Valley. It then switched (avulsed) at Piarere near Karapiro to flow into the Hamilton Basin. The name of the volcanogenic alluvium deposited by the ancient Waipa and Waikato rivers is the Hinuera Formation, and the surface of the plains is called the Hinuera Surface. The deposits of the Hinuera Formation are up to 60 m thick. The latest depositional episode in the Hamilton Basin was between c. 22,000 and 18,000 cal. years ago. Soon after, the ancestral Waikato River began to entrench into its modern channel, occasionally forming terraces at lower levels, as full forest vegetation was re-established over the landscape from c. 17,500 years ago. Thin but numerous tephra layers (each a few millimetres to a few centimetres in thickness) have blanketed much of the Hinuera Surface since it was abandoned c. 18,000 cal. years ago by the entrenching Waikato River. The tephra layers are well preserved in lake sediments (e.g. cores from Lake Rotoroa contain numerous tephra layers) and peat bogs that developed on or alongside the Hinuera deposits. The soil pattern on the ash-draped Hinuera Surface mimics the alluvial depositional environments: well drained soils occur on the slightly raised channel/bar deposits (Horotiu series, H: ashes on coarse alluvium) and poorly drained soils occur on lower-lying overbank flood deposits (Te Kowhai series, TK; Ngaroto series, Nt). In between are the Bruntwood soils (Br, well drained upper, poorly drained lower horizons) and Silverdale soils (S, moderately well drained upper and poorly drained lower horizons).

3. The lowermost terraces adjacent to the modern Waikato River mark deposition from a dramatic breakout flood event about AD 250 AD ago following the latest eruption of Taupo Volcano (in AD 232 ± 10). Huge quantities of pumiceous deposits were swept down the Waikato River, which rose several metres to tens of metres, and then left stranded as terrace deposits adjacent to the main river channel and up tributary valleys or gullies that drained into it. The deposits are known as the Taupo Pumic Alluvium and are up to ~30 m thick.

Soils developed on these materials (Waikato series, W) are weakly formed because of their relatively young age. The Cambridge Golf Course boasts that it was ‘sculptured by the Waikato River 15,000 years ago’. This is untrue: the course is dominated by deposits and paleochannels of the Taupo Pumice Alluvium of c. AD 250, only c. 1750 cal. years ago (the earlier Hinuera Formation materials are well buried underneath or were cannibalized during the Taupo break-out flood event).

4. Gullies are occasionally cut into the Hinuera Surface, usually draining towards the modern Waikato River. Soils of the gully sides, and terrace scarps, are Kikiriroa series (Kk) and soils on the recent alluvium in gully bottoms are Tamahana series (V). We suspect some are on the newly-discovered faults in Hamilton.

Past vegetation and climate

The first European descriptions of the central Waikato area noted that the landscape was covered largely in bracken and scrub. However, pollen and spore studies (palynology) of samples from cores taken from local lakes and peats demonstrated that the region was dominated by grassland from c. 20,000–17,500 cal. years ago when the climate was cool and dry. The region became forested from about 17,500 cal. years ago as climate became warmer and wetter, and forest cover persisted through to about AD 1300 when the prevalent podocarp-hardwood and kauri forests were partly destroyed by burning. Such burning is attributed largely to Polynesian activities. European settlers continued modifying the vegetation pattern.




Fig. 52 Summary of inferred vegetation and climate since c. 20,000 cal. years ago in the Hamilton area, and volcanic ash layers in a core from Lake Rotorua (from Lowe, 2014, after Newnham et al. 1989, 2013; Green and Lowe, 1994). See also more recent paper by Jara et al. (2017).




Fig. 53 Map of main geological units and associated soil series, Hamilton Gardens. See soil-landscape models above for key to abbreviations of names of soil series (H, Br, TK etc).



Fig. 54 'Puke' or small mounds for growing kumara in the unique early Maori garden (called Te Parapara) at the Hamilton Gardens, on the lowest terrace underlain by Taupo Pumice Alluvium near the Waikato River, Hamilton. Participants in the background are being spoken to by the garden's main architect, Wiremu Puke, in early December, 2014. See Gumbley et al. (2004) regarding soil modification in the central Waikato by early Maori.

References

- Abidin, Z., Matsue, N., Henmi, T., 2007. Differential formation of allophane and imogolite: experimental and molecular orbital study. *J. Computer-Aided Mater. Des.* 14, 5–18.
- Allbrook, R.F., 1983. Some physical properties of allophane soils from the North Island, New Zealand. *N. Z. J. Sci.* 26, 481–492.
- Allbrook, R.F., 1985. The effect of allophane on soil properties. *Appl. Clay Sci.* 1, 65–69.
- Almond, P.C., Tonkin, P.J. 1999. Pedogenesis by upbuilding in an extreme leaching and weathering environment, and slow loess accretion, south Westland, New Zealand. *Geoderma* 92, 1–36.
- Barrell, D.J.A., Almond, P.C., Vandergoes, M.J., Lowe, D.J., Newnham, R.M., NZ-INTIMATE members 2013. A composite pollen-based stratotype for inter-regional evaluation of climatic events in New Zealand over the past 30,000 years (NZ-INTIMATE project). *Quaternary Science Reviews* 74, 4–20.
- Birrell, K.S., Pullar, W.A., Heine, J.C. 1971. Pedological, chemical, and physical properties of organic horizons of paleosols underlying the Tarawera Formation. *New Zealand Journal of Science* 14, 187–218.
- Briggs, R.M., Houghton, B.F., McWilliams, M., Wilson, C.J.N. 2005. $^{40}\text{Ar}/^{39}\text{Ar}$ age of silicic volcanic rocks in the Tauranga-Kaimai area, New Zealand: dating the transition between volcanism in the Coromandel Arc and the Taupo Volcanic Zone. *New Zealand Journal of Geology and Geophysics* 48, 459–469.
- Charman, D. 2002. *Peats and Environmental Change*. Wiley, Chichester.
- Churchman, G.J., Lowe, D.J. 2012. Alteration, formation, and occurrence of minerals in soils. In: Huang, P.M., Li, Y., Sumner, M.E. (Eds), *Handbook of Soil Sciences*, 2nd edition. Vol. 1: Properties and Processes. CRC Press (Taylor & Francis), Boca Raton, FL, pp. 20.1–20.72.
- Clarkson, B. 2002. Swamps, fens and bogs. In: Clarkson, B.; Merrett, M.; Downs, T. (eds) "Botany of the Waikato", Waikato Botanical Society, 49–58.
- Clarkson, B.R., Patel, R.N., Clarkson, B.D. 1988. Composition and structure of forest overwhelmed at Pureora, central North Island, New Zealand, during the Taupo eruption (c. A.D. 130). *Journal of the Royal Society of New Zealand* 18, 417–436.

- Clarkson, B.R., Clarkson, B.D., Patel, R.N. 1992. The pre-Taupo eruption (c. AD 130) forest of the Bennydale-Pureora district, central North Island, New Zealand. *Journal of the Royal Society of New Zealand* 22, 61-76.
- Clarkson, B.R., McGlone, M.S., Lowe, D.J., Clarkson, B.D. 1995. Macrofossil and pollen representing forests of the pre-Taupo volcanic eruption (c. 1850 yr BP) era at Pureora and Bennydale, central North Island, New Zealand. *Journal of the Royal Society of New Zealand* 25, 263-281.
- Clarkson, B.R., Thompson, K., Schipper, L.A., McLeod, M. 1999. Moanatuatua Bog – proposed restoration of a New Zealand restiad peat bog ecosystem. Pp. 127-137 in W. Streever (ed) “An international perspective on peatland rehabilitation”. Kluwer, The Netherlands.
- Clarkson, B., Merrett, M., Downs, T. 2002. *Botany of the Waikato*. Waikato Botanica Society, Hamilton. 136 pp.
- Clarkson, B.R.; Schipper, L.A.; Lehmann, A. 2004. Vegetation and peat characteristics in the development of lowland and restiad peat bogs, North Island, New Zealand. *Wetlands* 24, 133-151.
- Clarkson, B.D., Clarkson, B.R., Downs, T.M. 2007. Indigenous vegetation types of Hamilton Ecological District. Centre for Biodiversity and Ecology Research (Hamilton) Contract Research Report 58. 16 pp.
- Clayden, B., Hewitt, A.E. 1994. Horizon notation for New Zealand soils. Manaaki Whenua Press, Lincoln, 30pp.
- Cole, J.W., Spinks, K.D., Deering, C.D., Nairn, I.A., Leonard, G.S. 2010. Volcanic and structural evolution of the Okataina Volcanic Centre; dominantly silicic volcanism associated with the Taupo Rift, New Zealand. *J. Volcanol. Geotherm. Res.* 190, 123–135.
- Cole, J.W., Deering, C.D., Nairn, Burt, R.M., Sewell, S., Shane, P.A.R., Matthews, N.E. 2014. Okataina Volcanic Centre, Taupo Volcanic Zone, New Zealand: A review of volcanism and synchronous pluton development in an active, dominantly silicic caldera system. *Earth-Sci. Rev.* 128, 1-17.
- Cornforth, I. 1998. Practical soil management. Lincoln University Press, Canterbury, and Whitirea Publishing, Wellington, 248pp.
- Danišić, M., Shane, P.A.R., Schmitt, A.K., Hogg, A.G., Santos, G.M., Storm, S., Evans, N.J., Fifield, L.K., Lindsay, J.M. 2012. Re-anchoring the late Pleistocene tephrochronology of New Zealand based on concordant radiocarbon ages and combined $^{238}\text{U}/^{230}\text{Th}$ disequilibrium and (U-Th)/He zircon ages. *Earth and Planetary Science Letters* 349-350, 240-250.
- de Lange, P.J., Heenan, P.B., Clarkson, B.D., Clarkson, B.R. 1999. Taxonomy, ecology, and conservation of *Sporodanthus* (Restionaceae) in New Zealand. *New Zealand Journal of Botany* 37, 413-431.
- de Ronde, C.E.J., Scott, B.J., Leonard, G.S., Calvert, A.T. 2016a. Evolution of the sublacustrine geothermal system at Lake Rotomahana, New Zealand: Effects of the 1886 Tarawera Rift eruption—an introduction. *Journal of Volcanology and Geothermal Research* 314, 1–9.
- de Ronde, C.E.J. et al. 2016b. The Pink and White Terraces of Lake Rotomahana: what was their fate after the 1886 Tarawera Rift eruption? *Journal of Volcanology and Geothermal Research* 314, 126–141.
- Froggatt, P.C., Lowe, D.J. 1990. A review of late Quaternary silicic and some other tephra formations from New Zealand: their stratigraphy, nomenclature, distribution, volume, and age. *New Zealand Journal of Geology and Geophysics* 33, 89-109.
- Froggatt, P.C., Wilson, C.J.N., Walker, G.P.L. 1981. Orientation of logs in the Taupo Ignimbrite as an indicator of flow direction and vent position. *Geology* 9, 109-111.
- Gehrels, M.J., Lowe, D.J., Hazell, Z.J., Newnham, R.M. 2006. A continuous 5300-yr Holocene cryptotephrostratigraphic record from northern New Zealand and implications for tephrochronology and volcanic-hazard assessment. *The Holocene* 16, 173-187.
- Gehrels, M.J., Newnham, R.M., Lowe, D.J., Wynne, S., Hazell, Z.J., Caseldine, C. 2008. Towards rapid assay of cryptotephra in peat cores: review and evaluation of various methods. *Quaternary Int.* 178, 68-84.
- Gehrels, M.J., Lowe, D.J., Newnham, R.M., Hogg, A.G. 2010. Enhanced record of tephra fallout since ~232 AD revealed by cryptotephra studies at Moanatuatua bog near Hamilton: implications for volcanic hazard analysis. Abstracts, GeoNZ 2010 Conference, Auckland. Geosciences Society of New Zealand Miscellaneous Publication 129A, p. 103.
- Goodrich, J.P., Campbell, D.I., Roulet, N.T., Clearwater, M.J., Schipper, L.A. 2015. Overriding control of methane flux temporal variability by water table dynamics in a Southern Hemisphere, raised bog. *Journal of Geophysical Research – Biogeosciences*, 120, DOI: 10.1002/2014JG002844.
- Goodrich, J.P., Campbell, D.I., Schipper, L.A. 2017. Southern Hemisphere bog persists as a strong carbon sink during droughts. *Biogeosciences* 14, 4563-4576.
- Grange, L.I. 1931. Volcanic ash showers. *New Zealand Journal of Science and Technology* 12, 228-240.
- Grange, L.I. 1937. The geology of the Rotorua-Taupo Subdivision. *New Zealand Geological Survey Bulletin* 37.
- Grange, L. I., Taylor, N. H., 1932. The distribution and field characteristics of bush-sick soils. Part 2A. In: *Bush Sickness*. N.Z. Department of Scientific and Industrial Research Bulletin 32, pp. 21-35.
- Gravley, D.M., Wilson, C.J.N., Leonard, G.S., Cole, J.W. 2007. Double trouble: paired ignimbrite eruptions and collateral subsidence in the Taupo Volcanic Zone, New Zealand. *Geological Society of America Bulletin* 119, 18-30.

- Green, J.D., Lowe, D.J. 1985. Stratigraphy and development of c. 17 000 year old Lake Maratoto, North Island, New Zealand, with some inferences about postglacial climatic change. *New Zealand Journal of Geology and Geophysics* 28, 675-699.
- Grimmett, R.E.R., Shorland, F.B. 1934. Some characteristics of "limonite" used in the cure and prevention of bush sickness. *Transactions of the Royal Society of New Zealand* 64, 191-213.
- Gumbley, W., Higham, T.F.G., Lowe, D.J. 2004. Prehistoric horticultural adaptation of soils in the middle Waikato Basin: review and evidence from S14/201 and S14/185, Hamilton. *New Zealand Journal of Archaeology* 25, 5-30.
- Harsh, J., 2012. Poorly crystalline aluminosilicate clay minerals. In: Huang, P.M., Li, Y., Sumner, M.E. (Eds), *Handbook of Soil Science*, 2nd edn, Vol. 1: Properties and processes. CRC Press, Boca Raton, pp. 23.21–23.13.
- Healy, J. 1992. Central volcanic region. In: Soons, J.M., Selby, M.J. (eds), "Landforms of New Zealand, 2nd Edition". Longman Paul, Auckland, pp. 256-286.
- Hendy, C.H. 2008. Bush sickness and the discovery of trace element deficiencies. In: Lowe, D.J. Soils – the most complex ecosystem in the world. "Science on the farm" website, School of Science, University of Waikato, Hamilton. URL: <http://sci.waikato.ac.nz/farm/content/soils.html>.
- Hewitt, A.E. 2010. *New Zealand Soil Classification*, 3rd edition. Landcare Research Science Series No. 1. Manaaki Whenua Press, Landcare Research, Lincoln, 136 pp.
- Hill, R.B., Tonkin, P.J., Almond, P.J. 1999. A revised soil stratigraphic interpretation of the geomorphology and late Quaternary erosion history of the southern Mamaku Plateau. *Geological Society of New Zealand Miscellaneous Publication* 107A, p. 64.
- Hiradate, S., Wada, S-I. 2005. Weathering process of volcanic glass to allophane determined by ²⁷Al and ²⁹Si solid-state NMR. *Clays and Clay Minerals* 53, 401-408.
- Hiradate, S., Nakadai, T., Shindo, H., Yoneyama, T. 2004. Carbon source of humic substances in some Japanese volcanic ash soils determined by carbon stable isotopic ratio, $\delta^{13}\text{C}$. *Geoderma* 119, 133-141.
- Hogg, A.G., Higham, T.F.G., Lowe, D.J., Palmer, J., Reimer, P., Newnham, R.M. 2003. A wiggle-match date for Polynesian settlement of New Zealand. *Antiquity* 77, 116-125.
- Hogg, A.G., Lowe, D.J., Palmer, J.G., Boswijk, G., Bronk Ramsey, C.J. 2012. Revised calendar date for the Taupo eruption derived by ¹⁴C wiggle-matching using a New Zealand kauri ¹⁴C calibration data set. *The Holocene* 22, 439-449.
- Hogg, A.G., Wilson, C.J.N., Lowe, D.J., Turney, C.S.M., White, P., Lorrey, A.M., Manning, S.W., Palmer, J.G., Bury, S., Brown, J., Southon, J., Petchey, F. 2019. Correspondence: The Taupo eruption occurred in 232 ± 10 CE, and not later. *Nature Communications* (submitted 28 Dec 2018, NCOMMS-18-7769176) – available as open access non-peer-reviewed preprint published at EarthArXiv (DOI: [10.31223/osf.io/7r5jp](https://doi.org/10.31223/osf.io/7r5jp)) (or <https://eartharxiv.org/7r5jp/>)
- Holt, K., Wallace, R.C., Neall, V.E., Kohn, B.P., Lowe, D.J. 2010. Quaternary tephra marker beds and their potential for palaeoenvironmental reconstruction on Chatham Islands east of New Zealand, southwest Pacific Ocean. *Journal of Quaternary Science* 25, 1169-1178.
- Holt, K.A., Lowe, D.J., Hogg, A.G., Wallace, R.C. 2011. Distal occurrence of mid-Holocene Whakatane Tephra on the Chatham Islands, New Zealand, and potential for cryptotephra studies. *Quaternary International* 246, 344-351.
- Hosono, M., Oba, Y., Sase, T., Utsugawa, T., Aoki, K. 1991. Holocene volcanic ash soils at Waimangu Road tephra section, North Island, New Zealand – soil formation-vegetation relationship. *The Quaternary Research (Japan)* 30, 91-101.
- Houghton, B.F., Wilson, C.J.N. 1986. Explosive rhyolitic volcanism: the case studies of Mayor Island and Taupo volcanoes. *New Zealand Geological Survey Record*, 12, 33-100.
- Houghton, B.F., Wilson, C.J.N., McWilliams, M.O., Lanphere, M.A., Weaver, S.D., Briggs, R.M., Pringle, M.S. 1995. Chronology and dynamics of a large silicic magmatic system: central Taupo Volcanic Zone, New Zealand. *Geology* 23, 13-16.
- Houghton, B.F., Carey, R.J., Cashman, K.V., Wilson, C.J.N., Hobden, B.J., Hammer, J.E. 2010. Diverse patterns of ascent, degassing, and eruption of rhyolite magma during the 1.8 ka Taupo eruption, New Zealand: evidence from clast vesicularity. *Journal of Volcanology and Geothermal Research* 195, 31-47.
- Houghton BF, Carey RJ, Rosenberg MD 2015. The Taupo eruption: "Ill wind" blows the ultraplinian type event down to Plinian. *Geology* 42, 459-461.
- Huang, Y-T., Lowe, D.J., Churchman, G.J., Schipper, L., Rawlence, N.J., Cooper, A. 2014. Carbon storage and DNA adsorption in allophanic soils and paleosols. In: Hartemink, A.E., McSweeney, K. (eds). "Soil Carbon". *Progress in Soil Science Series*, Springer, New York, pp. 163-172.
- Huang, Y.-T., Lowe, D.J., Churchman, G.J., Schipper, L.A., Cursons, R., Zhang, H., Chen, T.-Y., Cooper, A. 2016a. DNA adsorption by nanocrystalline allophane spherules and nanoaggregates, and implications for carbon sequestration in Andisols. *Applied Clay Science* 120, 40-50.

- Huang, Y.-T., Lowe, D.J., Zhang, H., Cursons, R., Young, J.M., Churchman, G.J., Schipper, L.A., Rawlence, N.J., Wood, J.R., Cooper, A. 2016b. A new method to extract and purify DNA from allophanic soils and paleosols, and potential for paleoenvironmental reconstruction and other applications. *Geoderma* 247, 114-125.
- Hudspith, V.A., Scott, A.C., Wilson, C.J.N., Collinson, M.E. 2009. Charring of woods by volcanic processes: an example from the Taupo ignimbrite, New Zealand. *Palaeogeography Palaeoclimatology Palaeoecology*, 291, 40-51.
- Jara, I.A., Newnham, R.M., Alloway, B.V., Wilmshurst, J.M., Rees, A.B.H. 2017. Pollen-based temperature and precipitation records of the past 14,600 years in northern New Zealand (37°S) and their linkages with the Southern Hemisphere atmospheric circulation. *The Holocene* 27, 1756-1768.
- Johnson, D.L., Watson-Stegner, D. 1987. Evolution model of pedogenesis. *Soil Science* 143, 349-366.
- Johnson, D.L., Keller, E.A., Rockwell, T.K. 1990. Dynamic pedogenesis: new views on some key soil concepts, and a model for interpreting Quaternary soils. *Quaternary Research* 33, 306-319.
- Johnson-Maynard, J.L., McDaniel, P.A., Ferguson, D.E., Falen, A.L. 1997. Chemical and mineralogical conversion of Andisols following invasion by bracken fern. *Soil Science Society of America Journal* 61, 549-555.
- Keam, R.F. 1988. Tarawera. Published by R.F. Keam, Auckland. 472 pp.
- Kondo, R., Childs, C., Atkinson, I. 1994. *Opal Phytoliths of New Zealand*. Mannaki Whenua Press, Lincon. 85p
- Keam, R.F. 2016. The Tarawera eruption, Lake Rotomahana, and the origin of the Pink and White Terraces. *Journal of Volcanology and Geothermal Research* 314, 10–38.
- Kilgour, G.N., Smith, R.T. 2008. Stratigraphy, dynamics, and eruption impacts of the dual magma Rotorua eruptive episode, Okataina Volcanic Centre, New Zealand. *N.Z. J. Geol. Geophys.* 51, 367-378.
- Kim, N.D., Taylor, M.D., Drewry, J.J. 2016. Anthropogenic fluorine accumulation in the Waikato and Bay of Plenty regions of New Zealand: comparison of field data with projections. *Environ Earth Sci* 75: 147 DOI 10.1007/s12665-015-4897-2
- Laubscher, N. 2014. Improvement in soil water availability in pastures by excavating and mixing buried soil horizons from multilayered Pumice Soils (Vitrandis) at Galatea, central North Island, New Zealand. MSc thesis, University of Waikato.
- Leonard, G.S., Begg, J.G., Wilson, C.J.J. (Compilers) 2010. *Geology of the Rotorua area: scale 1: 250,000*. Institute of Geological and Nuclear Sciences 1: 250,000 geological map 5. 1 sheet and 99 pp. Institute of Geological and Nuclear Sciences, Lower Hutt.
- Loame, R.C., Lowe, D.J., Pittari, A., Davies, S.M., Magill, C.R., Vandergoes, M.J., Rees, A.B.H., Ross, N. 2018. Using CT scanning for reconnaissance and detection of cryptotephra in ~22,000-yr-old lake sediments, central Waikato region, New Zealand. In: Hambach, U., Veres, D. (editors), *Book of Abstracts, Crossing New Frontiers: INTAV International Field Conference on Tephrochronology, 'Tephra Hunt in Transylvania'*, Moieciu de Sus, Romania, 24 June-1 July, pp. 41-42
- Lorrey, A.M., Woolley, J.-M. 2018. Locating relict sinter terrace sites at Lake Rotomahana, New Zealand, with Ferdinand von Hochstetter's legacy cartography, historic maps, and LIDAR. *Frontiers in Earth Science* 6: 205 (doi: 10.3389/feart.2018.00205).
- Lowe, D.J. 1986. Controls on the rates of weathering and clay mineral genesis in airfall tephra: a review and New Zealand case study. In: Colman, S.M., Dethier, D.P. (Editors), *Rates of Chemical Weathering of Rocks and Minerals*. Academic Press, Orlando, pp. 265-330.
- Lowe, D.J. 1990. Tephra studies in New Zealand: an historical review. *Journal of the Royal Society of New Zealand* 20, 119-150.
- Lowe, D.J. (compiler) 2008. *Guidebook for Pre-conference North Island Field Trip A1 'Ashes to Issues'*. Australian and New Zealand 4th Joint Soils Conference, Massey University, Palmerston North. New Zealand Society of Soil Science, 194 pp.
- Lowe, D.J. 2011. Tephrochronology and its application: a review. *Quaternary Geochronology* 6, 107-153.
- Lowe, D.J. 2019. Using soil stratigraphy and tephrochronology to understand the origin, age, and classification of a unique Late Quaternary tephra-derived Ultisol in New Zealand. *Quaternary* (submitted 31 December 2018)
- Lowe, D.J., Percival, H.J. 1993. Clay mineralogy of tephra and associated paleosols and soils, and hydrothermal deposits, North Island [New Zealand]. *Review and Guide F1 Field Tour – New Zealand*. Tenth International Clay Conference, Adelaide, Australia, 1-110.
- Lowe, D.J., de Lange, W.P. 2000. Volcano-meteorological tsunamis, the c. AD 200 Taupo eruption (New Zealand) and the possibility of a global tsunami. *The Holocene* 10, 401-407.
- Lowe, D.J., Palmer, D.J. 2005. Andisols of New Zealand and Australia. *Journal of Integrated Field Science* 2, 39-65.
- Lowe, D.J., McDaniel, P.A. 2010. Impacts of deforestation and burning, and the role of bracken fern, on properties of surficial or buried soil A-horizons In: Lowe, D.J., Neall, V.E., Hedley, M., Clothier, B., Mackay, A. *Guidebook for Pre-conference North Island, New Zealand 'Volcanoes to Oceans' field tour (27-30 July)*. 19th World Soils Congress, International Union of Soil Sciences, Brisbane. *Soil and Earth Sciences Occasional Publication No. 3*, Massey University, Palmerston North, pp. 4.39-4.42.

- Lowe, D.J., Tonkin, P.J. 2010. Unravelling upbuilding pedogenesis in tephra and loess sequences in New Zealand using tephrochronology. Proceedings published at <http://www.iuss.org>, IUSS 19th World Soil Congress, Brisbane (1-6 Aug., 2010), Symposium 1.3.2 Geochronological techniques and soil formation, pp. 34-37.
- Lowe, D.J., Pittari, A. 2014. An ashy septingentenarian: the Kaharoa tephra turns 700 (with notes on its volcanological, archaeological, and historical importance). *Geoscience Society of New Zealand Newsletter* 13, 35-46.
- Lowe, D.J., King, C.M. 2015. A dramatic landscape [Chapter 1]. In: King, C.M., Gaukrodger, D.J., Ritchie, N.A. (eds), *The Drama of Conservation – The History of Pureora Forest, New Zealand*. Springer, Berlin, and New Zealand Department of Conservation, Wellington, pp. 1-17.
- Lowe, D.J., McFadgen, B.G., Higham, T.F.G., Hogg, A.G., Froggatt, P.C., Nairn, I.A. 1998. Radiocarbon age of the Kaharoa Tephra, a key marker for late-Holocene stratigraphy and archaeology in New Zealand. *The Holocene* 8, 487-495.
- Lowe, D.J., Newnham, R.M., McFadgen, B.G., Higham, T.F.G. 2000. Tephtras and New Zealand archaeology. *Journal of Archaeological Science* 27, 859-870.
- Lowe, D.J., Shane, P.A.R., Alloway, B.V., Newnham, R.M. 2008. Fingerprints and age models for widespread New Zealand tephra marker beds erupted since 30,000 years ago: a framework for NZ-INTIMATE. *Quaternary Science Reviews* 27, 95-126.
- Lowe, D.J., Neall, V.E., Hedley, M., Clothier, B., Mackay, A. 2010. Guidebook for pre-conference North Island, New Zealand 'Volcanoes to Ocean' field tour (27-30 July, 2010). 19th World Soils Congress, International Union of Soil Sciences, Brisbane. *Soil and Earth Sciences Occasional Publication No. 3*, Massey University, Palmerston North, 239 pp.
- Lowe, D.J., Lanigan, K.M., Palmer, D.J. 2012. Where geology meets pedology: Late Quaternary tephtras, loess, and paleosols in the Mamaku Plateau and Lake Rerewhakaaitu areas. In: Pittari, A. (compiler). *Field Trip Guides, Geosciences 2012 Conference*, Hamilton, New Zealand. *Geoscience Society of New Zealand Miscellaneous Publication 134B*, pp. 2.1–2.45.
- Lowe, D.J., Blaauw, M., Hogg, A.G., Newnham, R.M. 2013. Ages of 24 widespread tephtras erupted since 30,000 years ago in New Zealand, with re-evaluation of the timing and palaeoclimatic implications of the late-glacial cool episode recorded at Kaipo bog. *Quaternary Science Reviews* 74, 170-194.
- Lowe, D.J., Holt, K.A., Shane, P.A.R., Hogg, A.G., Lorrey, A.M., Vincent, K.A., Esler, W.R., Cronin, S.J., Newnham, R.M. 2015. Developing a tephrostratigraphic framework for 60 to 30 cal ka for SHAPE in New Zealand. 19th INQUA Congress, Nagoya. Abstract T00616, 1 p.
- Lowe, D.J., Pearce, N.J.G., Jorgensen, M.A., Kuehn, S.C., Tryon, C.A., Hayward, C.L. 2017. Correlating tephtras and cryptotephtras using glass compositional analyses and numerical and statistical methods: review and evaluation. *Quaternary Science Reviews* 175, 1-44.
- Manville, V. 2001a. Environmental impacts of large-scale explosive rhyolitic eruptions in the central North Island. In: Smith, R.T. (ed), *Fieldtrip Guides, Geological Society of New Zealand Annual Conference*, University of Waikato, Hamilton. *Geological Society of New Zealand Miscellaneous Publication 110B*, 19pp.
- Manville, V. 2001b. Sedimentology and history of Lake Reporoa: an ephemeral supra-ignimbrite lake, Taupo Volcanic Zone, New Zealand. *Special Publication of the International Association of Sedimentology* 30, 109-140.
- Manville, V. 2002. Sedimentary and geomorphic responses to ignimbrite emplacement: readjustment of the Waikato River after the A.D. 181 Taupo eruption, New Zealand. *The Journal of Geology* 110, 519-541.
- Manville, V., Wilson, C.J.N. 2004. The 26.5 ka Oruanui eruption, New Zealand: a review of the roles of volcanism and climate in the post-eruptive sedimentary response. *New Zealand Journal of Geology and Geophysics* 47, 525-547.
- Manville, V., White, J.D.L., Houghton, B.F., Wilson, C.J.N. 1999. Paleohydrology and sedimentology of a post-1.8 ka breakout flood from intracaldera Lake Taupo, North Island, New Zealand. *Geological Society of America Bulletin* 111, 1435-1447.
- Manville, V., Hodgson, K.A., Nairn, I.A. 2007. A review of break-out floods from volcanogenic lakes in New Zealand. *New Zealand Journal of Geology and Geophysics* 50, 131-150.
- Manville, V., Swegschneider, B., Newton, E., White, J. D. L., Houghton, B.F., Wilson, C.J.N. 2009. Environmental impact of the 1.8 ka Taupo eruption, New Zealand: landscape responses to a large-scale explosive rhyolite eruption. *Sedimentary Geology* 220, 318-336.
- McLay, C.D.A., Allbrook, R.F., Thompson, K. 1992. Effect of development and cultivation on physical properties of peat soils in New Zealand. *Geoderma* 54, 23-37.
- McClelland, E., Wilson, C.J.N., Bardot, L. 2004. Paleotemperature determinations for the 1.8-ka Taupo ignimbrite, New Zealand, and implications for the emplacement history of a high-velocity pyroclastic flow. *Bulletin of Volcanology* 66, 492-513.
- McDaniel, P.A., Lowe, D.J., Arnalds, O., Ping, C.-L. 2012. Andisols. In: Huang, P.M., Li, Y., Sumner, M.E. (Eds), *Handbook of Soil Sciences, 2nd edition. Vol. 1: Properties and Processes*. CRC Press (Taylor & Francis), Boca Raton, FL, pp. 33.29-33.48.

- McGlone, M.S., Wilmshurst, J.M. 1999. Dating initial Maori environmental impact in New Zealand. *Quaternary International* 59, 5-16.
- McGlone, M., Wilmshurst, J.M., Leach, H.M. 2005. An ecological and historical review of bracken (*Pteridium esculentum*) in New Zealand, and its cultural significance. *New Zealand Journal of Ecology* 29, 165-184.
- McWethy, D.B., Whitlock, C., Wilmshurst, J.M., McGlone, M.S., Li, X. 2009. Rapid deforestation of South Island, New Zealand, by early Polynesian fires. *The Holocene* 19, 883-897.
- McWethy D.B., Wilmshurst, J.M., Whitlock, C., Wood, J.R., McGlone, M.S. 2014. A high-resolution chronology of rapid forest transitions following Polynesian arrival in New Zealand. *PLoS ONE* 9(11): e111328.
- Milne, D., Clayden, B., Singleton, P., Wilson, A. 1995. Soil description handbook. Revised edition. Manaaki Whenua Press, Lincoln, 157 pp.
- Molloy, L., Christie, Q. 1998. Soils in the New Zealand Landscape, 2nd edition. New Zealand Society of Soil Science, 253pp.
- Nairn, I.A. 1992. The Te Rere and Okareka eruptive episodes — Okataina Volcanic Centre, Taupo Volcanic Zone, New Zealand. *N.Z. J. Geol. Geophys.* 35, 93-108.
- Nairn, I.A. 2002. Geology of the Okataina Volcanic Centre. Institute of Geological and Nuclear Sciences Geological Map 25, 156p + map.
- Nathan, S. 2007. Lakes. Te Ara – the Encyclopedia of New Zealand. New Zealand Ministry for Heritage and Culture. Updated 5-Nov-2007. <http://www.TeAra.govt.nz/TheBush/Landscapes/Lakes/en>
- Neall, V.E. 2006. Volcanic soils, pp. 1-24. In: W. Verheyne (ed.) Land use and land cover. Encyclopaedia of life support systems (EOLSS). EOLSS Publishers with UNESCO, Oxford, U.K. Available online at: <http://www.eolss.net>
- Newnham, R.M., Lowe, D.J., McGlone, M.S., Wilmshurst, J.M., Higham, T.F.G. 1998. The Kaharoa Tephra as a critical datum for earliest human impact in northern New Zealand. *Journal of Archaeological Science* 25, 533-544.
- Palmer, D.J., Höck, B.K., Kimberley, M.O., Lowe, D.J., Payn, T.W. 2009. Comparison of spatial prediction techniques for developing *Pinus radiata* productivity surfaces across New Zealand. *Forest Ecology and Management* 258, 2046-2055.
- Parfitt, R. L., 1990. Allophane in New Zealand – a review. *Australian Journal of Soil Research* 28, 343–360.
- Parfitt, R.L. 2009. Allophane and imogolite: role in soil biogeochemical processes. *Clay Minerals* 44, 135-155.
- Parfitt, R.L., Pollok, J., Furkert, R.J. 1981. Guide book for tour 1, North Island. International 'Soils With Variable Charge' conference, Palmerston North (Feb 1981), pp. 84-90.
- Parfitt, R.L., Russell, M., Orbell, G.E. 1983. Weathering sequence of soils from volcanic ash involving allophane and halloysite. *Geoderma* 29, 41-57.
- Parfitt, R.L., Saigusa, M., Cowie, J.D. 1984. Allophane and halloysite formation in a volcanic ash bed under different moisture conditions. *Soil Science* 138, 360-364.
- Pronger, J., Schipper, L.A., Hill, R.B., Campbell, D.I., McLeod, M. 2014. Subsidence rates of drained agricultural peatlands in New Zealand and the relationship with time since drainage. *Journal of Environmental Quality* 43, 1442-1449.
- Rijkse, W.C., Vucetich, C.G. 1980. Soils of Wairakei Research Station, North Island, New Zealand. New Zealand Soil Survey Report 57. 26pp.
- Rijkse, W.C., Guinto, D.F. 2010. Soils of the Bay of Plenty. Volume 2: Central Bay of Plenty. Environment BOP Environmental Publication 2010/11-2. 198pp.
- Sahetapy-Engel, S., Self, S., Carey, R.J., Nairn, I.A. 2014. Deposition and generation of multiple widespread fall units from the c. AD 1314 Kaharoa rhyolitic eruption, Tarawera, New Zealand. *Bulletin of Volcanology* 76, article 836: DOI 10.1007/s00445-014-0836-4.
- Sase, T., Hosono, M. 1996. Vegetation histories of Holocene volcanic ash soils in Japan and New Zealand – relationship between genesis of melanic volcanic ash soils and human impact. *Earth Science (Chikyū Kagaku)* 50, 466-482.
- Sase, T., Hosono, M., Utsugawa, T., Aoki, K. 1988. Opal phytolith analysis of present and buried volcanic ash soils at Te Ngae Road tephra section, Rotorua Basin, North Island, New Zealand. *The Quaternary Research (Japan)* 27, 153-163.
- Schaetzl, R., Anderson, S. 2005. Soils – Genesis and Geomorphology. Cambridge University Press, New York, 817 pp.
- Selby, M.J., Hoskins, P.J. 1973. The erodibility of pumice soils of the North Island, New Zealand. *Journal of Hydrology (NZ)* 12, 32-56.
- Shane, P.A.R., Smith, V.C., Lowe, D.J., Nairn, I.A. 2003. Re-identification of c. 15 700 cal yr BP tephra bed at Kaipo Bog, eastern North Island: implications for dispersal of Rotorua and Puketarata tephra beds. *N.Z. J. Geol. Geophys.* 46, 591-596.
- Shoji, S., Kurebayashi, T., Yamada, I. 1990. Growth and chemical composition of Japanese pampas grass (*Miscanthus sinensis*) with special reference to the formation of dark-coloured Andisols in northeastern Japan. *Soil Science and Plant Nutrition* 36, 105-120.
- Shoji, S., Nanzyo, M., Dahlgren, R.A. 1993. *Volcanic Ash Soils: Genesis, Properties and Utilization*. Developments in Soil Science 21, Elsevier, Amsterdam, 288 pp.

- Singleton, P.L., McLeod, M., Percival, H.J. 1989. Allophane and halloysite content and soil solution silicon in soils from rhyolitic volcanic material, New Zealand. *Australian Journal of Soil Research* 27, 67-77.
- Smith, R.T., Houghton, B.F. 1995. Vent migration and changing eruptive style during the 1800a Taupo eruption: new evidence from the Hatepe and Rotongaio phreatoplinian ashes. *Bulletin of Volcanology* 57, 432-439.
- Smith, V.C., Shane, P.A.R., Nairn I.A. 2004. Reactivation of a rhyolitic magma body by new rhyolitic intrusion before the 15.8 ka Rotorua eruptive episode: implications for magma storage in the Okataina Volcanic Centre, New Zealand. *Jl Geolog Soc, London*. 161, 757–772.
- Smith, R.T., Lowe, D.J., Wright, I.C. 2007. Volcanoes. *Te Ara – The Encyclopedia of New Zealand*. New Zealand Ministry for Culture and Heritage, Wellington. Updated 16 April, 2007. <http://www.TeAra.govt.nz/EarthSeaAndSky/NaturalHazardsAndDisasters/Volcanoes/en>
- Soil Survey Staff 1999. *Soil Taxonomy*, 2nd Edition. United States Department of Agriculture Natural Resources Conservation Service, Agriculture Handbook No. 436, Washington D.C., 869 pp.
- Soil Survey Staff 2014. *Keys to Soil Taxonomy*, 12th edition. United States Department of Agriculture Natural Resources Conservation Service, Washington D.C., 360pp.
- Taylor, M.D., Lowe, D.J., Hardi, P., Smidt, G., Schnug, E. 2016. Comparing volcanic glass shards in unfertilised and fertilised Andisols derived from rhyolitic tephra, New Zealand: evidence for accelerated weathering and implications for land management. *Geoderma* 271, 91-98.
- Telfer, E.I. 2013. Leaching and uptake of N and pasture production associated with irrigation of treated wastewater, Taupo, New Zealand. MSc thesis, University of Waikato.
- Theng, B.K.G. 2019. *Clay Mineral Catalysis of Organic Reactions*. CRC Press (Taylor and Francis), Boca Raton, 425 pp.
- Tonkin, P.J. 2012. Soil investigation - DSIR soil science: 1930–1950. *Te Ara – the Encyclopedia of New Zealand*. New Zealand Ministry for Heritage and Culture. Updated 13-Jul-12. <http://www.TeAra.govt.nz/en/soil-investigation/>
- Underwood, E.J., Filmer, J.F. 1935. The determination of the biologically potent element (cobalt) in limonite. *The Australian Veterinary Journal* 11, 84-92.
- Vandergoes, M.J., Hogg, A.G., Lowe, D.J., Newnham, R.M., Denton, G.H., Southon, J., Barrell, D.J.A., Wilson, C.J.N., McGlone, M.S., Allan, A.S.R., Almond, P.C., Petchey, F., Dalbell, K., Dieffenbacher-Krall, A.C., Blaauw, M. 2013. A revised age for the Kawakawa/Oruanui tephra, a key marker for the Last Glacial Maximum in New Zealand. *Quaternary Science Reviews* 74, 195-201.
- Vucetich, C.G., Pullar, W.A. 1964. Stratigraphy of Holocene ash in the Rotorua and Gisborne districts. *New Zealand Geological Survey Bulletin* 73, Part 2, 43-88.
- Vucetich, C.G., Wells, N. 1978. Soils, agriculture, and forestry of Waiotapu region, central North Island, New Zealand. *New Zealand Soil Bureau Bulletin* 31, 100 pp + 5 map sheets and extended legends.
- Vucetich, C.G., Leamy, M.L., Popplewell, M.A., Ure, J., Taylor, C.R., Will, G.M., Sutton, J.A., Blakemore, L.C. 1960. Soils, forestry and agriculture of the northern part, Kaingaroa State Forest and the Galatea Basin. *New Zealand Soil Bureau Bulletin* 18, 51 pp + 2 map sheets.
- Wada, K. (editor) 1986. *Ando Soils in Japan*. Kyushu University Press, Fukuoka, 276 pp.
- Wagstaff, S.J., Clarkson, B.R. 2012. Systematics and ecology of the Australasian genus *Empodisma* (Restionaceae) and description of a new species from peatlands in northern NZ. *PhytoKeys* 13, 39-79.
- Walker, G.P.L. 1980. The Taupo plinian pumice: product of the most powerful known (ultraplinian) eruption? *Journal of Volcanology and Geothermal Research* 8, 69-94.
- Walker, G.P.L., Wilson, C.J.N. 1983. Lateral variations in the Taupo ignimbrite. *Journal of Volcanology and Geothermal Research* 18, 117-133.
- Walker, M., Johnsen, S., Rasmussen, S.O., Popp, T., Steffensen, J.-P., Gibbard, P., Hoek, W., Lowe, J.J., Andrews, J., Björck, S., Cwynar, L., Hughen, K., Kershaw, P., Kromer, B., Litt, T., Lowe, D.J., Nakagawa, T., Newnham, R.M., Schwander, J. 2009. Formal definition and dating of the GSSP (Global Stratotype Section and Point) for the base of the Holocene using the Greenland NGRIP ice core, and selected auxiliary records. *Journal of Quaternary Science* 24, 3-17.
- Watanabe, M., Sakagami, K. 1999. Recent pedological approach to reconstruction of the Quaternary environment in Japan and east Asia. *The Quaternary Research (Japan)* 38, 244-252.
- Watt, M.J., Palmer, D.J., Kimberley, M.O., Höck, B.K., Payn, T., Lowe, D.J. 2010. Development of models to predict *Pinus radiata* productivity throughout New Zealand. *Canadian Journal of Forest Research* 40, 488-499.
- Webb, T.H., Lilburne, L.R. 2011. Criteria for defining the soil family and soil sibling, the fourth and fifth categories of the *New Zealand Soil Classification* (2nd ed). *Landcare Research Science Series* 3, 38pp.
- Wilmshurst, J.M., McGlone, M.S. 1996. Forest disturbance in the central North Island, New Zealand, following the 1850 BP Taupo eruption. *The Holocene* 6, 399-411.

- Wilmshurst, J.M., Anderson, A.J., Higham, T.F.G., Worthy, T.H. 2008. Dating the late prehistoric dispersal of Polynesians to New Zealand using the commensal Pacific rat. *Proceedings of the National Academy of Sciences of the United States* 105, 7676-7680.
- Wilson, C.J.N. 1985. The Taupo eruption, New Zealand II. The Taupo ignimbrite. *Philosophical Transactions of the Royal Society, London A* 314, 229-310.
- Wilson, C.J.N. 1993. Stratigraphy, chronology, styles, and dynamics of late Quaternary eruptions from Taupo volcano, New Zealand. *Philosophical Transactions of the Royal Society of London A* 343, 205-306.
- Wilson, C.J.N. 1994. Post-conference tour day 1: Hamilton-Tokaanu. In: Lowe, D.J. (ed) *Conference Tour Guides. International Inter-INQUA Field Conference on Tephrochronology, Loess and Paleopedology*. University of Waikato, Hamilton, 74-100.
- Wilson C.J.N. 2001. The 26.5 ka Oruanui eruption, New Zealand: an introduction and overview. *Journal of Volcanology and Geothermal Research* 112, 133-174.
- Wilson C, Leonard G 2015. Slumbering giants. In Graham IJ (ed.), *A continent on the move*, 2nd edition. Geoscience Society of New Zealand, *Miscellaneous Publication* 141, pp. 168-171.
- Wilson, C.J.N., Walker, G.P.L. 1985. The Taupo eruption, New Zealand. I. General aspects. *Philosophical Transactions of the Royal Society, London A* 314, 199-228.
- Wilson, C.J.N., Blake, S., Charlier, B.L.A., Sutton, A.N. 2006. The 26.5 ka Oruanui eruption, Taupo volcano, New Zealand: development, characteristics and evacuation of a large magma body. *Journal of Petrology* 47, 35-69.
- Wilson, C.J.N., Gravley, D.M., Leonard, G.S., Rowland, J.V. 2009. Volcanism in the central Taupo Volcanic Zone, New Zealand: tempo, styles and controls. In: Thordarson, T., Self, S., Larsen, G., Rowland, S.K., Hoskuldsson, A. (Eds), *Studies in Volcanology: the Legacy of George Walker*. Special Publications of IAVCEI (Geological Society, London) 2, pp. 225-247.
- Yamamoto, K., Tate, K.R., Churchman, G.J. 1989. A comparison of the humic substances from some volcanic ash soils in New Zealand and Japan. *Soil Science and Plant Nutrition* 35, 257-270.
- Yuan, G., Theng, B.K.G. 2012. Clay-organic interactions in soil environments. In: Huang, P.M., Li, Y., Sumner, M.E. (Eds), *Handbook of soil sciences*, 2nd edn, Vol. 2: Resource management and environmental impact. CRC Press, Boca Raton, pp. 2.1-2.20.
- Yuan, G., Wada, S.-I. 2012. Allophane and imogolite nanoparticles in soil and their environmental applications. In: Barnard, A.S., Guo, H. (Eds), *Nature's nanostructures*. Pan Stanford, Singapore, pp. 494-515.